

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

B15

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 5 :

C07D 233/64, 263/32, 277/30
A61K 31/415, 31/42, 31/425

A1

(11) International Publication Number:

WO 93/02057

(43) International Publication Date:

4 February 1993 (04.02.93)

(21) International Application Number: PCT/US92/06047

(22) International Filing Date: 17 July 1992 (17.07.92)

(30) Priority data:

731,563 17 July 1991 (17.07.91) US
870,975 20 April 1992 (20.04.92) US

(60) Parent Applications or Grants

(63) Related by Continuation

US 870,975 (CIP)
Filed on 19 April 1992 (19.04.92)
US 731,563 (CIP)
Filed on 17 July 1991 (17.07.91)

(71) Applicant (for all designated States except US): SMITH-KLINE BEECHAM CORPORATION [US/US]; Corporate Patents-U.S., UW2220, 709 Swedeland Road, P.O. Box 1539, King of Prussia, PA 19406-0939 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): CARR, Thomas, Joseph [US/US]; 1232 Harrison Avenue, Phoenixville, PA 19460 (US). DeMARSH, Peter, Lawrence [US/US]; 203 Armstead Court, Downingtown, PA 19335 (US). DREYER, Geoffrey, Bainbridge [US/US]; 2 Marlin Drive, Marlvern, PA 19355 (US). FENWICK, Ashley, Edward [GB/US]; 115 Pine Tree Road, Radnor, PA 19087 (US).

(74) Agents: KINZIG, Charles, M. et al.; SmithKline Beecham Corporation, Corporate Patents - U.S., UW2220, 709 Swedeland Road, P.O. Box 1538, King of Prussia, PA 19406-0939 (US).

(81) Designated States: AT, AU, BB, BG, BR, CA, CH, CS, DE, DK, ES, FI, GB, HU, JP, KP, KR, LK, LU, MG, MN, MW, NL, NO, PL, RO, RU, SD, SE, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LU, MC, NL, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, SN, TD, TG).

Published

With international search report.
Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.

(54) Title: RETROVIRAL PROTEASE INHIBITORS

(57) Abstract

The present invention provides compounds, more particularly dipeptide analogs, which bind to retroviral proteases. These compounds are inhibitors of retroviral proteases and are useful for treating diseases related to infection by retroviruses.

Chemical structure		Molecular weight		Molar ratio	
1	2	3	4	5	6
7	8	9	10	11	12
13	14	15	16	17	18
19	20	21	22	23	24
25	26	27	28	29	30
31	32	33	34	35	36
37	38	39	40	41	42
43	44	45	46	47	48
49	50	51	52	53	54
55	56	57	58	59	60
61	62	63	64	65	66
67	68	69	70	71	72
73	74	75	76	77	78
79	80	81	82	83	84
85	86	87	88	89	90
91	92	93	94	95	96
97	98	99	100	101	102
103	104	105	106	107	108
109	110	111	112	113	114
115	116	117	118	119	120
121	122	123	124	125	126
127	128	129	130	131	132
133	134	135	136	137	138
139	140	141	142	143	144
145	146	147	148	149	150
151	152	153	154	155	156
157	158	159	160	161	162
163	164	165	166	167	168
169	170	171	172	173	174
175	176	177	178	179	180
181	182	183	184	185	186
187	188	189	190	191	192
193	194	195	196	197	198
199	200	201	202	203	204
205	206	207	208	209	210
211	212	213	214	215	216
217	218	219	220	221	222
223	224	225	226	227	228
229	230	231	232	233	234
235	236	237	238	239	240
241	242	243	244	245	246
247	248	249	250	251	252
253	254	255	256	257	258
259	260	261	262	263	264
265	266	267	268	269	270
271	272	273	274	275	276
277	278	279	280	281	282
283	284	285	286	287	288
289	290	291	292	293	294
295	296	297	298	299	300
301	302	303	304	305	306
307	308	309	310	311	312
313	314	315	316	317	318
319	320	321	322	323	324
325	326	327	328	329	330
331	332	333	334	335	336
337	338	339	340	341	342
343	344	345	346	347	348
349	350	351	352	353	354
355	356	357	358	359	360
361	362	363	364	365	366
367	368	369	370	371	372
373	374	375	376	377	378
379	380	381	382	383	384
385	386	387	388	389	390
391	392	393	394	395	396
397	398	399	400	401	402
403	404	405	406	407	408
409	410	411	412	413	414
415	416	417	418	419	420
421	422	423	424	425	426
427	428	429	430	431	432
433	434	435	436	437	438
439	440	441	442	443	444
445	446	447	448	449	450
451	452	453	454	455	456
457	458	459	460	461	462
463	464	465	466	467	468
469	470	471	472	473	474
475	476	477	478	479	480
481	482	483	484	485	486
487	488	489	490	491	492
493	494	495	496	497	498
499	500	501	502	503	504
505	506	507	508	509	510
511	512	513	514	515	516
517	518	519	520	521	522
523	524	525	526	527	528
529	530	531	532	533	534
535	536	537	538	539	540
541	542	543	544	545	546
547	548	549	550	551	552
553	554	555	556	557	558
559	560	561	562	563	564
565	566	567	568	569	570
571	572	573	574	575	576
577	578	579	580	581	582
583	584	585	586	587	588
589	590	591	592	593	594
595	596	597	598	599	600
601	602	603	604	605	606
607	608	609	610	611	612
613	614	615	616	617	618
619	620	621	622	623	624
625	626	627	628	629	630
631	632	633	634	635	636
637	638	639	640	641	642
643	644	645	646	647	648
649	650	651	652	653	654
655	656	657	658	659	660
661	662	663	664	665	666
667	668	669	670	671	672
673	674	675	676	677	678
679	680	681	682	683	684
685	686	687	688	689	690
691	692	693	694	695	696
697	698	699	700	701	702
703	704	705	706	707	708
709	710	711	712	713	714
715	716	717	718	719	720
721	722	723	724	725	726
727	728	729	730	731	732
733	734	735	736	737	738
739	740	741	742	743	744
745	746	747	748	749	750
751	752	753	754	755	756
757	758	759	760	761	762
763	764	765	766	767	768
769	770	771	772	773	774
775	776	777	778	779	780
781	782	783	784	785	786
787	788	789	790	791	792
793	794	795	796	797	798
799	800	801	802	803	804
805	806	807	808	809	810
811	812	813	814	815	816
817	818	819	820	821	822
823	824	825	826	827	828
829	830	831	832	833	834
835	836	837	838	839	840
841	842	843	844	845	846
847	848	849	850	851	852
853	854	855	856	857	858
859	860	861	862	863	864
865	866	867	868	869	870
871	872	873	874	875	876
877	878	879	880	881	882
883	884	885	886	887	888
889	890	891	892	893	894
895	896	897	898	899	900
901	902	903	904	905	906
907	908	909	910	911	912
913	914	915	916	917	918
919	920	921	922	923	924
925	926	927	928	929	930
931	932	933	934	935	936
937	938	939	940	941	942
943	944	945	946	947	948
949	950	951	952	953	954
955	956	957	958	959	960
961	962	963	964	965	966
967	968	969	970	971	972
973	974	975	976	977	978
979	980	981	982	983	984
985	986	987	988	989	990
991	992	993	994	995	996
997	998	999	1000	1001	1002

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FI	Finland	ML	Mali
AU	Australia	FR	France	MN	Mongolia
BB	Barbados	GA	Gabon	MR	Mauritania
BE	Belgium	GB	United Kingdom	MW	Malawi
BF	Burkina Faso	GN	Guinea	NL	Netherlands
BG	Bulgaria	GR	Greece	NO	Norway
BJ	Benin	HU	Hungary	PL	Poland
BR	Brazil	IE	Ireland	RO	Romania
CA	Canada	IT	Italy	RU	Russian Federation
CF	Central African Republic	JP	Japan	SD	Sudan
CG	Congo	KP	Democratic People's Republic of Korea	SE	Sweden
CH	Switzerland	KR	Republic of Korea	SN	Senegal
CI	Côte d'Ivoire	LJ	Liechtenstein	SU	Soviet Union
CM	Cameroun	LK	Sri Lanka	TD	Chad
CS	Czechoslovakia	LU	Luxembourg	TG	Togo
DE	Germany	MC	Monaco	US	United States of America
DK	Denmark	MG	Madagascar		
ES	Spain				

- 1 -

RETROVIRAL PROTEASE INHIBITORS**Background of the Invention**

The present invention relates to retroviral protease inhibitor compounds, pharmaceutical compositions thereof, and a method of treating retroviral diseases therewith, including a method of treating disease states associated with human immunodeficiency virus (HIV-1, HIV-2).

Retroviruses, that is, viruses within the family of Retroviridae, are a class of viruses which transport their genetic material as ribonucleic acid rather than deoxyribonucleic acid. Also known as RNA-tumor viruses, their presence has been associated with a wide range of diseases in humans and animals. They are believed to be the causative agents in pathological states associated with infection by Rous sarcoma virus (RSV), murine leukemia virus (MLV), mouse mammary tumor virus (MMTV), feline leukemia virus (FeLV), bovine leukemia virus (BLV), Mason-Pfizer monkey virus (MPMV), simian sarcoma virus (SSV), simian acquired immunodeficiency syndrome (SAIDS), human T-lymphotropic virus (HTLV-I, -II) and human immunodeficiency virus (HIV-1, HIV-2), which is the etiologic agent of AIDS.

- 2 -

(acquired immunodeficiency syndrome) and AIDS related complexes, and many others. Although the pathogens have, in many of these cases, been isolated, no effective method for treating this type of infection has been developed.

5 Retroviral replication occurs only in host cells. Critical to this replication is the production of functional viral proteins. Protein synthesis is accomplished by translation of the appropriate open reading frames into polyprotein constructs, which are processed, at least in
10 part, by a viral protease into the functional proteins. The proteolytic activity provided by the viral protease in processing the polyproteins cannot be provided by the host and is essential to the life cycle of the retrovirus. In fact, it has been demonstrated that retroviruses which lack
15 the protease or contain a mutated form of it, lack infectivity. See Katoh et al., *Virology*, 145, 280-92 (1985), Crawford, et al., *J. Virol.*, 53, 899-907 (1985) and Debouk, et al., *Proc. Natl. Acad. Sci. USA*, 84, 8903-6 (1987). Inhibition of retroviral protease, therefore, presents a
20 method of therapy for retroviral disease.

The use of isosteric replacements has been disclosed as a strategy for the development of protease inhibitors for HIV-1. European Patent Applications EP-A 337 714, EP-A 357 332, EP-A 346 847, EP-A 342 541, EP-A 352 000, EP-A 393 445
25 and EP-A 434 365 are representative, and are incorporated herein by reference. These references disclose dipeptide analogs of the natural polyprotein substrates of retroviral proteases. As discussed therein, these dipeptide analogs bind selectively and competitively to retroviral proteases;
30 however, the protease is unable to cleave the carbon-carbon bond presented to it instead of the scissile amide bond of the natural substrate. Thus, such compounds are useful for inhibiting viral replication by inactivation of the protease. The incorporation of heterocyclic elements in the P3' and P4'
35 substrate positions of compounds containing a dipeptide isostere has been disclosed by deSolms et al., *J. Med. Chem.*, 34, 2852 (1991). However, these compounds can be less than desirable for obtaining optimal drug delivery in mammalian

- 3 -

organisms, particularly in humans. Some of these compounds can also have a less than desirable serum half-life, and therefore duration of action, because they contain amide bonds in relatively high proportion, and thus are prone to metabolic degradation, hepatic clearance, or other elimination mechanisms.

There exists a need for novel compounds which inhibit retroviral protease activity, and a need for compounds which possess desirable pharmacokinetic properties for good drug delivery and metabolic stability for good serum half-life and duration of action. Such pharmaceutical uses provide therapies for retroviral diseases in mammals, especially in humans, which have been heretofore difficult to treat.

15

SUMMARY OF THE INVENTION

The present invention provides compounds, hereinafter represented as formula (I), which bind to retroviral proteases. These compounds are inhibitors of retroviral proteases and are useful for treating diseases related to infection by retroviruses.

The present invention also provides a pharmaceutical composition comprising a compound of formula (I) and a pharmaceutically acceptable carrier.

The present invention additionally provides a method for treating retroviral disease, comprising administering to a mammal in need thereof an effective amount of a compound of formula (I).


30

DETAILED DESCRIPTION OF THE INVENTION

The compounds of the present invention are illustrated by formula (I):

35

- 4 -



wherein:


R¹ and R³ are each independently Q, Q-C₁₋₆alkyl,
 5 Q-C₂₋₆alkenyl, Q-C₂₋₆alkynyl or C₁₋₆alkyl substituted by one
 to five fluorine atoms, each optionally substituted by R²³;

Q is H, C₃₋₆cycloalkyl, C₅₋₆cycloalkenyl, Ar or Het

R² is H or OH;

R⁴ is R⁶-NR¹¹ or CONR¹¹CHR⁶R⁷;

10 R⁵ is R⁶-NR¹¹ or R¹⁰-NR¹¹;



R⁶ is

X is NR¹¹, O or S;

R⁷ is Q, Q-C₁₋₆alkyl or Q-C₂₋₆alkenyl;

R⁸ and R⁹ are each independently H, OH, halo, NO₂, COR¹²,
 15 CF₃, Ar, C₁₋₆alkyl-R¹⁵, or R¹⁷(R¹⁸R¹⁹)_m, or together form a
 fused C₂₋₄alkylene, aryl or heteroaryl moiety;

R¹⁰ is A-(B)_n;

R¹¹ is H or C₁₋₄alkyl;

R¹² is R⁷, OR⁷, NR⁷R¹¹ or an amino acid or amino alcohol;

20 B is an amino acid;

A is H, Ar, Het, R¹⁷(R¹⁸R¹⁹)_m, Ar-W, Het-W or

R¹⁷(R¹⁸R¹⁹)_m-W, or phthaloyl each optionally substituted by
 one to three groups chosen from R¹⁵ or C₁₋₆alkyl-R¹⁵;

W is C=O, OC(=O), NR¹¹C(=O), SC(=O), NR¹¹C(=S), SO₂,

25 NR¹¹SO₂ or P(=O)(OR²²);

R¹⁵ is H, nitro, C₁₋₆alkoxy, C₁₋₆alkylthio, O(C=O)R¹⁶,
 C=OR²², CO₂R²², CON(R¹⁶)₂, N(R²²)₂, NHC(=N)NH-A, I, Br, Cl, F,
 OR¹⁰, or OH, provided that when R¹⁵ is a substituent of the
 carbon adjacent to W, R¹⁵ is not halogen or OH when W is

30 OC(=O) or NHCO;

R¹⁶ is H or C₁₋₆alkyl;

R¹⁷, R¹⁸ and R¹⁹ are independently: i) H, R¹⁵ or
 C₁₋₄alkyl, C₂₋₆alkenyl, phenyl, naphthyl, C₃₋₆cycloalkyl or
 Het, each optionally substituted by one to three R¹⁵ or

- 5 -

or R^{15} - C_{1-6} alkyl groups, or ii) R^{17} is as above and $(R^{18}R^{19}C)$ are joined together to form a phenyl, naphthyl, C_{3-6} cycloalkyl or Heteroring, or iii) R^{17} is as above and R^{18} and R^{19} together are

$=O$; or R^{18} and R^{19} together are $-(CH_2)_q$ where q is 2-5;

R^{22} is H , C_{1-6} alkyl, phenyl or phenyl- C_{1-4} alkyl;

R^{23} is $-X''-(CH_2)_qNR^{24}R^{25}$, $X''=[(CH_2)_rO]_sR^{26}$,

or $CH_2X''[(CH_2)_rO]_sR^{26}$, or benzofuryl, indolyl, azacycloalkyl,

azabicyclo C_{7-11} cycloalkyl or benzopiperidinyl, optionally

substituted with C_{1-4} alkyl;

q is 2-5;

s is 1-6 and r is 1-3 within each repeating unit s ;

X'' is CH_2 , O , S or NH ;

(30) X'' is CH_2 , NR' , O , S , SO or SO_2 ;

R^{24} and R^{25} are i) C_{1-6} alkyl, optionally substituted by

15 OH , C_{1-3} alkoxy, or $N(R')_2$, ii) the same or different and

joined together to form a 5-7 member heterocycle containing

up to two additional heteroatoms selected from NR , O , S , SO ,

or SO_2 , said heterocycle optionally substituted with C_{1-4} alkyl,

iii) aromatic heterocycle, optionally substituted with

20 C_{1-4} alkyl or $N(R')_2$;

R' is H or C_{1-4} alkyl;

R^{26} is H , C_{1-4} alkyl, $C(=O)R^{27}$, $C(=O)U[(CH_2)_mO]NR'$,

$P(=O)(OM)_2$, CO_2R^{27} , $C(=O)NR^{27}R^{28}$, where M is a mono or

divalent metal ion, and U is NR' or O ;

25 R^{27} is C_{1-6} alkyl or Ar , optionally substituted with one

or more hydroxy, carboxy, halo, C_{1-3} alkoxy, $CONR'^2$, NR'^2 ,

CO_2R' , $SO_2NR'^2$, CH_2NR_2 , $NR'COR'$, $NR'SO_2R'$, $X''[(CH_2)_rO]_sR'$ or

$CH_2X''[(CH_2)_rO]_sR'$;

R^{28} is H , C_{1-6} alkyl or together with R^{27} forms a 5-7

30 membered heterocycle or a 6 membered heterocycle containing a

heteroatom selected from N , O and S ;

m is 1-4; and

n is 0 or 1;

or a pharmaceutically acceptable salt thereof.

35 Also included in this invention are pharmaceutically

acceptable addition salts, complexes or prodrugs of the

compounds of this invention. Prodrugs are considered to be

- 6 -

any covalently bonded carriers which release the active parent drug according to formula (I) *in vivo*.

Formula (I) is intended to encompass all unique nonracemic stereoisomers which may occur due to the presence of asymmetric carbon atoms in the molecule. Such compounds may occur as pure enantiomers or diastereomers or as a mixture of individual stereoisomers. The definition of any substituent moiety which may occur more than once in formula (I) is independent of any other occurrence. Combinations of substituents and/or variables are permissible only if such combinations result in stable compounds.

Compounds of this invention which include acyclic double bonds may be present in either the *cis* (Z) or *trans* (E) geometrical configuration with respect to any two substituents.

When X is NH, it will be appreciated that the heterocyclic ring is an imidazole which can undergo tautomerization. All tautomeric forms of the imidazole are within the scope of this invention.

Suitably R^1 and R^3 are C_1 -alkyl, Ar- C_1 -alkyl, Ar- C_2 -alkenyl, Ar- C_2 -alkynyl, C_1 -alkyl optionally substituted by one to five fluorine atoms or benzyl substituted in the 4-position by R^{23} . Preferably R^1 is benzyl and R^3 is benzyl, 4-hydroxybenzyl or phenylpropenyl.

Suitably R^2 is H.

Suitably X is S or N- R^{11} . Preferably X is NH.

Preferably R^4 is CONR¹¹CHR⁶R⁷.

Suitably R^5 is R^{10} -NR¹¹. Preferably R^5 is t-butyl, butyloxycarbonylamino or isopropylloxycarbonylamino.

Suitably R^7 is H, C_1 -alkyl, C_3 -cycloalkyl, phenyl or benzyl. Preferably R^7 is C_1 -alkyl. Isopropyl is most preferred.

Suitably R^8 is H, C_1 -alkyl, COR¹², NO₂ or Br. Preferably R^8 is H.

Suitably R^9 is H, NO₂, Br, COR¹², CF₃, Ar, C_1 -alkyl or C_1 -alkyl- R^{15} , wherein R^{12} is H, C_1 -alkyl, Ar, OC₁-alkyl, NH₂, and R^{15} is OH. Preferably R^9 is H or COR¹².

Suitably B is Ala or Val. Preferably m is 0 and B is absent.

Suitably A is Het, $R^{17}(R^{18}R^{19}C)_m-W$, Ar-W or Het-W.

Suitably R^{17} , R^{18} and R^{19} are H, or C_{1-4} alkyl, Het or Ar optionally substituted by one or two R^{15} or C_{1-6} alkyl- R^{15} , or ($R^{18}R^{19}C$) are joined together to form a phenyl, C_{3-6} cycloalkyl or Het ring.

Suitably W is C=O, OC(=O), NHC(=O), NHC(=S), or SC(C=O).

Suitably $R^{17}(R^{18}R^{19}C)_m-$ is Ar-CH₂, Ar, Het, Het-CH₂, C_{1-6} alkyl or C_{3-6} cycloalkyl optionally substituted by one to three groups selected from R^{15} . Suitably R^{15} is OH. When R^{17} or ($R^{18}R^{19}C$) are Het or Ar, Het is suitably quinolinyl, pyridyl, imidazolyl, thiazolyl, tetrahydrothiopyranyl or tetrahydropyranyl and Ar is phenyl.

Suitably R^{23} is hydroxy- C_{1-4} alkoxy, C_{1-4} alkoxy- C_{1-4} alkoxy, or $-O(CH_2)_2NR^{24}R^{25}$, wherein R^{24} and R^{25} are a 5- or 6-membered heterocycle, such as morpholino.

In one preferred embodiment W is C=O.

In another preferred embodiment W is OC(=O).

In a third preferred embodiment R^{10} is C_{1-6} alkylOC(=O) or C_{5-6} cycloalkylOC(=O) substituted by one or two OH or CH₂OH groups.

Representative compounds of this invention are:

(2R, 4S, 5S, 1'S)-2-phenylmethyl-4-hydroxy-5-(t-butoxycarbonyl)-amino-6-phenyl-N-[1'-isopropyl-1'-(4-aminocarbonyl-thiazo-2-yl)]methyl-hexanamide;

(2R, 4S, 5S, 1'S)-2-phenylmethyl-4-hydroxy-5-(t-butoxycarbonyl)-amino-6-phenyl-N-[1'-isopropyl-1'-(thiazo-2-yl)]methyl-hexanamide;

(2R, 4S, 5S, 1'S)-2-phenylmethyl-4-hydroxy-5-(t-butoxycarbonyl)-amino-6-phenyl-N-(1'-imidazo-2-yl)methyl-hexanamide hydrochloride;

(2R, 4S, 5S, 1'S)-2-phenylmethyl-4-hydroxy-5-(t-butoxycarbonyl)-amino-6-phenyl-N-[1'-methyl-1'-(imidazo-2-yl)]methyl-hexanamide hydrochloride;

(2R, 4S, 5S, 1'S)-2-phenylmethyl-4-hydroxy-5-(t-butoxycarbonyl)-amino-6-phenyl-N-[1'-benzyl-1'-(imidazo-2-yl)]methyl-hexanamide hydrochloride;

- (2R,4S,5S,1'S)-5-(carbobenzyloxy)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide;
- (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-[1'-isopropyl-1'-(4,5-dimethyl)imidazol-2-yl]methyl-6-phenyl-2-phenylmethyl-hexanamide;
- (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-[1'-isopropyl-1'-(N'-methyl)imidazol-2-yl]methyl-6-phenyl-2-phenylmethyl-hexanamide;
- 10 (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-(3-phenylpropargyl)hexanamide;
- (2R,4S,5S,1'S)-5-(benzyloxyethoxycarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide;
- 15 (2R,4S,5S,1'S)-5-(methoxycarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide;
- (2R,4S,5S,1'S)-5-(ethoxycarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide;
- 20 (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-(3-phenyl-2-propenyl)hexanamide;
- 25 (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-[1'-isopropyl-1'-(4-nitroimidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide;
- (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-(1'-ethyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide;
- 30 (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-(1'-propyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide;
- (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-[1'-isopropyl-1'-(4-bromoimidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide;
- 35

- (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-[1'-isopropyl-1'-(4,5-dibromoimidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide;
- (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-[1'-isopropyl-1'-(4-methylimidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide;
- (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-[1'-isopropyl-1'-(4-trifluoromethylimidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide;
- (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-methyl-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide;
- (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-[1'-isopropyl-1'-(4-carbomethoxyimidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide;
- (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-[1'-isopropyl-1'-(4-methylcarbonylimidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide;
- (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-[1'-isopropyl-1'-(4-phenylcarbonyl-imidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide;
- (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-[1'-isopropyl-1'-(4-formylimidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide;
- (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-[1'-isopropyl-1'-(4-(hydroxymethyl)-imidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide;
- (2R,4S,5S,1'S)-5-((tetrahydrothiopyran-4-yl)oxycarbonyl)-amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide;
- (2R,4S,5S,1'S)-5-((tetrahydro-4H-pyran-4-yl)oxycarbonyl)-amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide;
- (2R,4S,5S,1'S)-5-(4-picolinyloxy)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide;

- 10 -

- (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-(4,4,4-trifluorobut-1-yl)hexanamide;
- (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-[1'-isopropyl-1'-(4-((1R)-1-hydroxyethyl)-imidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide;
- (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-[1'-(1-methyl)propyl-1'-(imidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide;
- 10 (2R,4S,5S,1'S)-5-(propylaminocarbonyl)amino-4-hydroxy-N-[1'-isopropyl-1'-(imidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide;
- (2R,4S,5S,1'S)-5-(4-hydroxybutanoyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethylhexanamide;
- 15 (2R,4S,5S,1'S)-2-phenylmethyl-4-hydroxy-5-(benzyloxy-carbonyl)valylamino-6-phenyl-N-(1'-isobutyl-1'-imidazo-2-yl)methyl-hexanamide;
- (2R,4S,5S,1'S)-2-phenylmethyl-4-hydroxy-5-(N-acetylvalyl)-amino-6-phenyl-N-(1'-isobutyl-1'-imidazo-2-yl)methyl-hexanamide;
- 20 (2R,4S,5S,1'S)-5-[(imidazol-2-yl)methyloxycarbonyl]amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide;
- 25 (2R,4S,5S,1'S,1"RS)-5-[(1"-(imidazol-2-yl)-2"-methyl)-propyloxycarbonyl]amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide;
- (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-[1'-isopropyl-1'-(4-(imidazol-2-yl)imidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide;
- 30 (2R,4S,5S,1'S)-5-(1-oxo-thian-4-yl)oxycarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethylhexanamide;
- (2R,4S,5S,1'S)-5-((tetrahydrosulfonylpyran-4-yl)oxycarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethylhexanamide;
- 35 (2R,4S,5S,1'S)-5-((1,1-dimethyl-2-(benzyloxycarbonyl)glycyloxy)ethoxycarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-

- imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide
hydrochloride salt;
(2R,4S,5S,1'S)-5-((1,1-dimethyl-2-glycyloxy)ethoxycarbonyl)-
amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-
5 phenyl-2-phenylmethyl-hexanamidedihydrochloridesalt;
(2R,4S,5S,1'S)-5-((1-acetyl)amino-4-hydroxy-N-(1'-isopropyl-
1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethylhexanamide;
(2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-(1'-
isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-(4-
10 benzyloxyphenylmethyl)hexanamide;
(2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-(1'-
isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-(4-
hydroxyphenylmethyl)hexanamide;
(2R,4S,5S)-5-(t-butoxycarbonyl)amino-4-hydroxy-2-
15 phenylmethyl-6-phenyl-N-[1'-cyclopropyl-1'-imidazol-2-
yl)methyl-hexanamide;
(2R,4S,5S,1'S)-5-((isopropylthiol)carbonyl)-amino-4-hydroxy-
2-phenylmethyl-6-phenyl-N-[1-isopropyl-1'-imidazol-2-
yl)methyl-hexanamide;
20 (2R,4S,5S,1'S)-5-[3-(1H-imidazol-2-yl)-3-hydroxy-4-
methylpentylamido]-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-
yl)methyl-6-phenyl-2-phenylmethyl-hexanamide;
(2R,4S,5S,1'S)-5-[(4-methoxyphenoxy)carbonyl]amino-4-hydroxy-
N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-
25 phenylmethyl-hexanamide;
(2R,4S,5S,1'S)-5-(t-butylaminocarbonyl)amino-4-hydroxy-N-(1'-
isopropyl-1'-imidazol-2-yl)methyl-6-phenylmethyl-hexanamide;
(2R,4S,5S,1'S)-5-(methylaminocarbonyl)-
amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-
30 phenylmethyl-hexanamide;
(2R,4S,5S,1'S)-5-phenylaminocarbonyl)amino-4-hydroxy-N-(1'-
isopropyl-1'-imidazol-2-yl)methyl-6-phenylmethyl-hexamide;
(2R,4S,5S,1'S)-5-N-(propylaminocarbonyl)amino-4-hydroxy-N-
(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenylmethyl-
35 hexamide;
(2R,4S,5S,1'S)-5-(n-propylaminothiono)amino-4-hydroxy-N-
(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenylmethyl-hexamide;

- 2R, 4S, 5S, 1'S)-5-(isopropylaminocarbonyl)-amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenylmethyl-hexamide;
- (2R, 4S, 5S, 1'S)-5-(aminocarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenylmethyl-hexamide;
- (2R, 4S, 5S, 1'S)-5-(6-quinolinylmethyloxy-carbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenylmethyl-hexanamide;
- (2R, 4S, 5S, 1'S)-5-(benzoyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenylmethyl-hexanamide;
- (2R, 4S, 5S, 1'S)-5-(2-furylcarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenylmethyl-hexanamide;
- (2R, 4S, 5S, 1'S)-5-(4-methoxybenzoyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenylmethyl-hexanamide;
- (2R, 4S, 5S, 1'S)-5-benzylcarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenylmethyl-hexamide;
- (2R, 4S, 5S, 1'S)-5-(4-hydroxybenzoyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenylmethyl-hexanamide;
- (2R, 4S, 5S, 1'S)-5-(cinnamoyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenylmethyl-hexanamide;
- (2R, 4S, 5S, 1'S)-5-(2-hydroxybenzoyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenylmethyl-hexanamide;
- (2R, 4S, 5S, 1'S)-5-(imidazol-4-yl-acetyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenylmethyl-hexanamide;
- (2R, 4S, 5S, 1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-[1'-isopropyl-1'-(4-carbomethoxyethylimidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide;
- (2R, 4S, 5S, 1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-[1'-isopropyl-1'-(4-carboxamidoimidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide;
- (2R, 4S, 5S, 1'S)-2-phenylmethyl-4-hydroxy-5-(5-(1-oxopropyl)-2-thiazolyl)amino)-6-phenyl-N-(1'-isopropyl-1'-(imidazol-2-yl)methyl-hexanamide;
- (2R, 4S, 5S, 1'S)-2-phenylmethyl-4-hydroxy-5-(5-(1-oxopropyl)-2-thiazolyl)amino)-6-phenyl-N-(1'-isopropyl-1'-(imidazol-2-yl)methyl-hexanamide;

(2R,4S,5S,1'S)-2-phenylmethyl-4-hydroxy-5-(5-propyl-2-thiazolyl)amino)-6-phenyl-N-(1'-isopropyl-1'-(imidazo-2-yl)methyl-hexanamide; and

(2R,4S,5S,1'S)-5-(nicotiny) amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenylmethyl-hexanamide.

Another group of preferred representative compounds are:

(2R,4S,5S,1'S)-5-[di(hydroxymethyl)-methoxycarbonyl] amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide;

(2R,4S,5S,1'S)-5-(1,1-dimethyl-2-acetoxyethoxycarbonyl) amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide;

(2R,4S,5S,1'S)-5-((1,1-dimethyl-2-hydroxy) ethoxy-carbonyl) amino-4-hydroxy-N-(1'-isopropyl-1'-(4-

15 isopropylcarbonyl-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide dihydrochloride salt;

(2R,4S,5S,1'S)-5-((1S)-1-methyl-2-hydroxyethoxycarbonyl)-amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethylhexanamide;

(2R,4S,5S,1'S)-5-((1R)-1-methyl-2-hydroxyethoxycarbonyl)-amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethylhexanamide;

(2R,4S,5S,1'S)-5-(1-hydroxymethyl-cyclopentyloxycarbonyl)-amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide;

(2R,4S,5S,1'S)-5-(1,1-dimethyl-2-hydroxyethoxycarbonyl) amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide hydrochloride;

(2R,4S,5S,1'S)-5-(hydroxyethoxycarbonyl) amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethylhexanamide; and

(2R,4S,5S,1'S)-5-(2-hydroxy-1-methylethoxycarbonyl) amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethylhexanamide.

35 More preferred representative compounds are:

(2R,4S,5S,1'S)-2-phenylmethyl-4-hydroxy-5-(t-butoxycarbonyl)-amino-6-phenyl-N-(1'-isopropyl-1'-(imidazo-2-yl)methyl-hexanamid hydrochloride;

- (2R,4S,5S,1'S)-5-(isopropoxycarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide;
- (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-[1'-isopropyl-1'-(4-isopropylcarbonyl-imidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide;
- (2R,4S,5S,1'S)-5-(1,1-dimethyl-2-hydroxyethoxycarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide hydrochloride;
- (2R,4S,5S,1'S)-5-(hydroxyethoxycarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethylhexanamide; and
- (2R,4S,5S,1'S)-5-(2-hydroxy-1-methylethoxycarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethylhexanamide.

The term "alkyl" refers to a straight or branched chain alkyl radical of the indicated number of carbon atoms.

"C₁₋₄alkyl" as applied herein is meant to include methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl; "C₁₋₆alkyl" includes additionally pentyl, isopentyl, 2-methylbutyl, 1-methylbutyl, 2-ethylpropyl, neopentyl, n-hexyl 2,2-dimethylbutyl, 2-methylpentyl, and the like.

"Alkoxy" refers to an alkyl group of the indicated number of carbon atoms attached through a bridging oxygen atom.

"Alkylthio" refers to an alkyl group of the indicated number of carbon atoms attached through a bridging sulfur atom.

The term "substituted alkyl" as used herein is meant to include C₁₋₆alkyl, Ar-C₁₋₆alkyl, Het-C₁₋₆alkyl, C₂₋₆alkenyl, Ar-C₂₋₆alkenyl, Het-C₂₋₆alkenyl, C₃₋₆cycloalkyl-C₁₋₆alkyl, C₃₋₆cycloalkenyl-C₁₋₆alkyl or C₁₋₆alkyl substituted with acyl or hydroxyl.

"Alkenyl" refers to a straight or branched hydrocarbon chain of the indicated number of carbon atoms, which contains one or more carbon-carbon double bonds at any stable point along the chain, such as ethenyl, propenyl, butenyl, pentenyl, 2-methylpropenyl, hexenyl, and the like.

"Alkynyl" refers to a straight or branched hydrocarbon chain of the indicated number of carbon atoms which contains

- 15 -

...a carbon-carbon triple bond at any stable point along the chain, such as ethynyl, 2-propynyl, 2-butyne, 4-pentynyl, 2-methyl-3-propynyl, hexynyl and the like.

The term "acyl" means $R_{12}-CO$, wherein R_{12} is H,

5 C₁-6alkyl, Ar-C₁-6alkyl, Het-C₁-6alkyl, C₂-6alkenyl,

Ar-C₂-6alkenyl, Het-C₂-6alkenyl, C₃-6cycloalkyl-C₁-6alkyl,

10 C₅-6cycloalkenyl-C₁-6alkyl, OH, NHR_{13} , wherein R_{13} is H,

C₁-6alkyl, Ar-C₁-6alkyl, Het-C₁-6alkyl, C₂-6alkenyl,

Ar-C₂-6alkenyl, Het-C₂-6alkenyl, C₃-6cycloalkyl-C₁-6alkyl, or

15 C₃-6cycloalkyl, or C₅-6cycloalkenyl-C₁-6alkyl; or an α -amino acid or an α -amino alcohol bonded at the nitrogen.

"Cycloalkyl" refers to a saturated ring group of the indicated number of carbon atoms. "C₃-7cycloalkyl" includes

cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and

15 cycloheptyl. "Cycloalkenyl" refers to a saturated ring group

of the indicated number of carbon atoms, having at least one

endocyclic carbon-carbon double bond. "C₅-7cycloalkenyl"

includes cyclopentenyl, cyclohexenyl and cycloheptenyl.

"Aryl", abbreviated as Ar, refers to phenyl or naphthyl,

20 optionally substituted with one to three halo, OH, OR¹⁰,

C₁-6alkyl, C₁-6alkoxy, C₁-6alkylthio, C₁-6alkylamino, CF₃,

amino, NO₂, carboxy, C₁-4alkylcarbonyl, aminocarbonyl,

C₁-6alkyl-Het, C₁-6alkoxy-Het, C₁-6alkyl-phenyl, C₁-6alkoxy-

phenyl, C₁-6alkyl-, C₁-6alkoxy-, HetC₁-6alkyl-, HetC₁-6alkoxy-,

25 phenylC₁-6alkyl-, phenylC₁-6alkoxy- or phenyloxy.

As used herein except where noted, the term

"heterocycle", abbreviated as "Het", represents a stable 5- to 7-membered monocyclic or a stable 7- to 10-membered

bicyclic heterocyclic ring, which is either saturated or

30 unsaturated, and which consists of carbon atoms and from one

to three heteroatoms selected from the group consisting of N,

O and S, and wherein the nitrogen and sulfur heteroatoms may

optionally be oxidized, and the nitrogen heteroatom may

optionally be quaternized, and including any bicyclic group

35 in which any of the above-defined heterocyclic rings is fused

to a benzene ring. The heterocyclic ring may be attached at

any heteroatom or carbon atom which results in the creation

of a stable structure, and may optionally be substituted with

one to three halo, OH, alkyl, alkoxy, alkyl-Het, alkoxy-Het, alkyl-phenyl, alkoxy-phenyl. Examples of such heterocyclic elements include piperidinyl, piperazinyl, 2-oxopiperazinyl, 2-oxopiperidinyl, 2-oxopyrrolodiny, 2-oxoazepinyl, azepinyl, 5 pyrrolyl, 4-piperidonyl, pyrrolidinyl, pyrazolyl, pyrazolidinyl, imidazolyl, pyridyl, pyrazinyl, oxazolidinyl, oxazolinyl, oxazolyl, isoxazolyl, morpholinyl, thiazolidinyl, thiazolinyl, thiazolyl, quinuclidinyl, indolyl, quinolinyl, isoquinolinyl, benzimidazolyl, benzopyranyl, benzoxazolyl, 10 furyl, pyranyl, tetrahydrofuryl, tetrahydropyranyl, thienyl, benzoxazolyl, thiamorpholinyl sulfoxide, thiamorpholinyl sulfone, and oxadiazolyl. Heteroaryl refers to a heterocycle which has aromatic character (eg., characterized by delocalized electron resonance and the ability to sustain a 15 ring current). Pyridine, imidazole, thiazole, furan and oxazole are examples of heteroaryl rings.


"Amino acid" means the D- or L- isomer of alanine, arginine, asparagine, aspartic acid, cysteine, glutamine, glutamic acid, glycine, histidine, isoleucine, leucine, 20 lysine, methionine, phenylalanine, proline, serine, threonine, tryptophan, tyrosine, valine or trifluoroalanine. In general, the amino acid abbreviations follow the IUPAC-IUB Joint Commission on Biochemical Nomenclature as described in *Eur. J. Biochem.*, 158, 9 (1984). Usually lipophilic amino 25 acids are preferred for the moiety B, for instance, Val, Ala, Leu and Ile. It will be understood that a linkage B-O refers to an oxygen atom bonded to the carboxyl group of an amino acid, and that a B-N linkage indicates a nitrogen atom bonded to the carboxyl group of an amino acid, as in an amide bond. 30 "Amino alcohol" refers to an amino acid in which the carboxyl group has been reduced to a methylene hydroxy group.

Certain chemical names are abbreviated herein for the sake of convenience. Boc refers to the *t*-butoxycarbonyl radical. Cbz refers to the carbobenzyloxy radical. Bzl 35 refers to the benzyl radical. Ac refers to acetyl. Ph refers to phenyl. BOP refers to benzotriazol-1-yloxy-tris(dimethylamino)phosphonium hexafluorophosphate. DCC refers to dicyclohexylcarbodiimide. DMAP refers to

- 17 -


dimethylamin-opyridine. DMSO refers to dimethylsulfoxide. HOBT refers to 1-hydroxybenzotriazole. NMM is N-methylmorpholine. DTT is dithiothreitol. EDTA is ethylenediamine tetraacetic acid. DIEA is diisopropyl ethylamine. DBU is 1,8 diazobicyclo[5.4.0]undec-7-ene. DMSO is dimethylsulfoxide. DMF is dimethyl formamide; Lawesson's reagent is 2,4-bis(4-methoxyphenyl)-1,3-dithia-2,4-diphosphetane-2,4-disulfide and THF is tetrahydrofuran. HF refers to hydrofluoric acid and TFA refers to trifluoroacetic acid.

The compounds of formula (I):




wherein R^4 is $\text{CO-NR}^1\text{CHR}^6\text{R}^7$, R^5 is $\text{R}^{10}\text{R}^{11}\text{N-}$, and R^1 , R^2 , R^3 and R^6 are as defined in formula (I), are prepared by:

1) (a) coupling a compound of the formula (II):



with a compound of formula (III):




where R^1 , R^2 , R^3 , R^5 , R^6 and R^7 are as defined for

formula (I) with any reactive groups protected, Pr^1 is H or a hydroxyl protecting group, and L' is OH or a leaving group;

or

(b) coupling a compound of the formula (IV):



with a compound of the formula (V):


- 18 -

A'-(B')_n-L' (v)

(v)

wherein A' and B' are as defined in formula (I) with any reactive groups protected; or

5 (c) coupling a compound of the formula (VI):



(VI)

with a compound of the formula (VII):

A'-L'

(VII)

and,

2) if appropriate, a coupling agent; and

3) removing any protecting groups and

15 4) forming a pharmaceutically acceptable salt thereof.

The coupling reactions may be accomplished by activating the substrate with a reactive functional group *in situ* or prior to the coupling reaction, such that it is reactive with an amino group. For instance, acids may be converted to acid chlorides, bromides, activated esters or anhydrides, or by

20 adding a coupling reagent. Coupling agents are well known in the art for activating a functional group *in situ*.

Exemplary of such agents are DCC and other carbodiimides, DMAPEC, BOP and PPA. These coupling agents may optionally be

25 used with other reagents, such as HOBT, NMM and DMAP, which may facilitate the reaction.

Suitable leaving groups, L', are those which are displaceable by an amino group, such as bromo, chloro, a substituted acyl (eg. trifluoroacetyl, bromobenzoyl,

30 nitrobenzoyl) or a substituted phenol (eg. 4-nitrophenol) and the like. If L' is OH, so that A-OH is an acid, it will be appropriate to use a coupling agent as hereinbefore described.

For instance:

When A is a substituted alkyl group, such as $R^{17}(R^{18}R^{19}C)_m$, L' may be a bromo, chloro, iodo or an alkyl or aryl sulfonate.

When A is $R^{17}(R^{18}R^{19}C)_m-W$, Ar-W or Het-W, and W is C=O, A-L' may be a carboxylic acid halide, activated ester or anhydride, or a carboxylic acid in the presence of a coupling agent. Methods for preparing such compounds are well known.

When W is OC=O, A-L' may be a chloro- or bromo-formate, or an activated carbonate. Haloformates may be prepared by reacting the appropriate alcohol with phosgene or carbonyldibromide.

Activated carbonates may be prepared by reacting the appropriate alcohol with a suitable carbonate such as bis(4-nitrophenyl)carbonate.

When W is SO₂, A-L' may be a sulfonyl halide which may be prepared from the corresponding sulfonic acid.


When W is SC=O, A-L' may be a halothioformate, which may be prepared from a carbonyldihalide and an appropriate mercaptan.

When W is PO(OR²²), A-L' may be a phosphonyl halide, which may be prepared from the corresponding phosphonic acid.

Compounds wherein A is $R^{17}(R^{18}R^{19}C)_m-W$, Ar-W or Het-W, and W is NR'C=O are ureas, and may be prepared by reacting a compound of formula (VII) with an isocyanate of the formula $R^{17}(R^{18}R^{19}C)_m-NCO$, Ar-NCO or Het-NCO, in a suitable solvent such as methylene chloride, optionally with heating.

Compounds of formula (III), wherein X is nitrogen, are imidazoles and may be prepared according to Scheme 1, wherein Pr² is a removeable amino protecting group, and R^{7'}, R^{8'} and R^{9'} correspond to R⁷, R⁸ and R⁹ as defined for formula (I), or a group which may be converted into R⁷, R⁸ or R⁹, with any reactive groups protected.

Scheme 1




- 20 -

The amino aldehydes are generally known or are prepared by methods well known in the art, for instance, by reduction of a suitable α -amino acid ester with diisobutylaluminum hydride. Further reaction of the aldehyde with a gem dialdehyde, or diketone, and ammonia yields the desired imidazole. Alkylation and further modification of the substituent groups of the imidazole are within the skill of the art. Such a method and other methods for preparing imidazoles are disclosed, for instance, by Baldwin et al., *J. Med. Chem.*, 29, 1065 (1986), *Angew. Chem. Int.*, 22, 560 (1983); and Hughey et al., *Synthesis*, 1489 (1980). Alternately, acyl imidazoles may be prepared by coupling an α -amino acid to a substituted 4-amino-isoxazole, and subsequent reduction and base catalyzed rearrangement as disclosed generally by Reiter, L.A., *J. Org. Chem.*, 52, 2714 (1987). Intermediate compounds of formula (VIII) are a part of this invention. Preferably, $R^{7'}$ is C_1 -alkyl and more preferably C_3 -alkyl. Suitably, $R^{8'}$ and $R^{9'}$ are H, NO_2 , Br, COR^{12} , CF_3 , Ar, C_1 -alkyl or C_1 -alkyl- R^{15} , wherein R^{12} is H, C_1 -alkyl, Ar, OC_1 -alkyl, NH_2 , and R^{15} is OH or a protected hydroxyl group. Preferably R^9 is H or COR^{12} .

Compounds of formula (III), wherein X is sulfur, are thiazoles and may be prepared according to Scheme 2, wherein L'' is a suitable displaceable group.

Scheme 2




Accordingly, a thioamide is reacted with a ketone or aldehyde. Thioamides are commonly prepared from carboxamides by reacting the corresponding carboxamides with a reagent such as Lawessons reagent, as disclosed, for instance, by Hamada et al., *Tet. Lett.*, 931 (1991). Suitable displaceable groups are those which are displaced by a sulfur nucleophile,

such as chloride, bromide, iodide, mesylate, p-toluenesulfonate groups, and the like.

Compounds of formula (III), wherein X is oxygen, are oxazoles and may be prepared according to Scheme 3 from common amino acids.

Scheme 3




10 Typically the acid may be coupled to an appropriately substituted amino alcohol by common techniques, as described above, and cyclized by treatment with thionyl chloride to yield an oxazoline, as described by Meyers et al., J. Org. Chem., 43, 1372 (1978). Oxidation of the oxazoline, such as
15 described by Evans et al., J. Org. Chem., 44, 497 (1979), yields an oxazole.

The compounds of formula (II), (IV) and (VI), wherein R² is H, are prepared, for instance, according to Scheme 4.

- 22 -

Scheme 4





Other methods for preparing protected 5-amino-4-hydroxy-2,5-disubstituted-pentanoate esters and acids, and the corresponding γ -lactones, are well known and are disclosed, for instance, in Szelke et al., U.S. Patent 4,713,455, Boger et al., U.S. Patent 4,661,473, EP-A 0 352 000, Evans et al., J. Org. Chem., 50, 4615 (1985), Kempf, J. Org. Chem., 51, 3921 (1986), Fray et al., J. Org. Chem., 51, 4828 (1986), Halladay et al., Tett. Lett., 24, 4401 (1983), Wuts et al., J. Org. Chem., 53, 4503 (1988), DeCamp et al., Tett. Lett., 32,1867 (1991), and Szelke et al., WO 84/03044, all of which are incorporated herein by reference.


The compounds of formula (II), (IV) and (VI), wherein R^2 is OH, are also prepared by methods common in the art such as those disclosed in U.S. Patent 4,864,017, and Thaisrivongs et al., J. Med. Chem., 30, 976 (1987).

Compounds of formula (I), wherein R^5 is $R^6\text{-NR}^{11}$, are prepared according to Scheme 5, Scheme 6 or Scheme 7:


Scheme 5



Scheme 6



Scheme 7




wherein R^{1'}-R^{4'}, R^{7'} and R^{8'} are as defined in formula (I)

with any reactive groups protected, L' is a leaving group,

15 such as halogen, and Pr¹ is a hydroxy-protecting group

- 24 -

Compounds wherein R^4 is R^6NR^{11} are prepared in an analogous manner from a compound of formula (IX):



5

Suitable protecting groups for the amino, hydroxyl, carboxylic acid, mercaptan group, and reagents for deprotecting these functional groups are disclosed in Greene et al., PROTECTIVE GROUPS IN ORGANIC SYNTHESIS, Second Edition, John Wiley and Sons, New York, 1991. Deprotection indicates the removal of the protecting group and replacement with an hydrogen atom. In particular, suitably substituted acetyl, benzyl and silyl groups are useful for protecting the hydroxyl group. The acetyl group is commonly removed by reacting the compound with a base, such as an alkali metal hydroxide, in a mixture of an alcohol and water. The silyl group, such as trimethyl silyl, dimethyl-t-butyl silyl, and t-butyl-diphenyl silyl may be removed by a fluoride reagent, such as a tetra-alkyl ammonium fluoride, or by acid hydrolysis. The benzyl group may be removed by catalytic hydrogenation.

Suitable protecting groups for the amino group are those disclosed by Greene et al., as indicated previously. The benzyloxycarbonyl and t-butoxycarbonyl groups are especially useful amino protecting groups.

The present invention includes pharmaceutically acceptable acid addition salts. Acid addition salts of the present compounds are prepared in a standard manner in a suitable solvent from the parent compound and an excess of an acid, such as hydrochloric, hydrobromic, sulfuric, phosphoric, acetic, maleic, succinic or methanesulfonic. The acetate salt form is especially useful. If the final compound contains an acidic group, cationic salts may be prepared. Typically the parent compound is treated with an excess of an alkaline reagent, such as a hydroxide, carbonate or alkoxide, containing the appropriate cation. Cations such

- 25 -

as Na^+ , K^+ , Ca^{++} and NH_4^+ are examples of cations present in pharmaceutically acceptable salts. Certain of the compounds form inner salts or zwitterions which may also be acceptable. The compounds of the present invention selectively bind to retroviral proteases in the same manner as the virally coded natural substrates of the proteases and compete with these substrates for protease, thereby serving to inhibit viral replication by blocking the formation of crucial viral proteins from polyprotein precursors by the protease, and hence, to inhibit disease progression in vivo. The present compounds achieve such beneficial therapeutic effect because they contain unique structural features which impart desirable pharmacokinetic properties to the compounds. One such property is long duration of action. We have found that substitution of a heterocycle, especially imidazole, in the putative P2' position of the present compounds affords compounds which retain good enzyme binding affinity, good antiviral activity, a favorable duration of action and water solubility for good drug delivery.

When a compound of the present invention is administered to an animal infected or potentially infected with a retrovirus, viral replication is inhibited and hence disease progression is retarded. Inasmuch as the amino acid sequences of the protease binding and peptide bond cleavage sites of various retroviruses appear to be highly conserved, an inhibitor is likely to be broadly active against more than one retrovirus. Also, DNA viruses which are dependant upon virally encoded proteases, such as the hepatitis virus, may also be susceptible to such treatment.

The compounds of formula (I) are used to inhibit retroviral replication, and are useful in treating mammals, particularly human patients, who are infected with susceptible retroviruses and require such treatment. The method of treating a retroviral disease in a mammal, particularly a human, comprises internally administering (e.g. orally, parent rally, buccally, trans-dermally, rectally or by insufflation) to said mammal an effective amount of a compound of formula (I), preferably dispersed in

a pharmaceutical carrier. Dosage units of the active ingredient may be selected by procedures routine to one skilled in the art, and are generally in the range of 0.01-50 mg/kg. These dosage units may be administered one to ten times daily for acute or chronic infection. Preferably the compound is administered at a level of 1-10 mg/kg, two to four times daily. No unacceptable toxicological effects are indicated when compounds of this invention are administered in the above noted dosage range.

10. The present invention also provides a method of treating disease states associated with HIV infection or Acquired Immune Deficiency Syndrome (AIDS), comprising administering an effective amount of a compound of formula (I), preferably dispersed in a pharmaceutical carrier.

15. Beneficial effects may be realized by co-administering, individually or in combination, other anti-viral agents with the protease inhibiting compounds of the present invention. Examples of anti-viral agents include nucleoside analogues, phosphonoformate, rifabutin, ribavirin, phosphonothioate oligodeoxynucleotides, castanospermine, dextran sulfate, 20 alpha interferon and amplitagen. Nucleoside analogues, which include 2',3'-dideoxycytidine (ddC), 2',3'-dideoxyadenine (ddA) and 3'-azido-2',3'-dideoxythymide (AZT), are especially useful. AZT is a preferred agent. Suitably, pharmaceutical compositions comprise an anti-viral agent, a protease 25 inhibiting compound of the present invention, and a pharmaceutically acceptable carrier.

This invention is also a pharmaceutical formulation which comprises a compound of formula (I) and a pharmaceutically acceptable carrier. Pharmaceutical acceptable carrier are well known in the art and are disclosed, for instance, in SPROWL'S AMERICAN PHARMACY, Dittert, L. (ed.), J.B. Lippincott Co., Philadelphia, 1974, and REMINGTON'S PHARMACEUTICAL SCIENCES, Gennaro, A. (ed.), 35 Mack Publishing Co., Easton, Pennsylvania, 1985.

Pharmaceutical compositions of the compounds of the present invention, or derivatives thereof, may be formulated as solutions or lyophilized powders for parenteral

- 27 -

administration. Powders may be reconstituted by addition of a suitable diluent or other pharmaceutically acceptable carrier prior to use. The liquid formulation is generally a buffered, isotonic, aqueous solution. Examples of suitable diluents are normal isotonic saline solution, standard 5% dextrose in water or buffered sodium or ammonium acetate solution. Such formulation is especially suitable for parenteral administration, but may also be used for oral administration or contained in a metered dose inhaler or nebulizer for insufflation. It may be desirable to add excipients such as ethanol, polyvinylpyrrolidone, gelatin, hydroxy cellulose, acacia, polyethylene glycol, mannitol, sodium chloride or sodium citrate.

Alternately, these compounds may be encapsulated, tableted or prepared in a emulsion or syrup for oral administration. Pharmaceutically acceptable solid or liquid carriers may be added to enhance or stabilize the composition, or to facilitate preparation of the composition. Liquid carriers include syrup, soy bean oil, peanut oil, olive oil, glycerin, saline, ethanol, and water.

Solubilizing agents, such as dimethylsulfoxide, ethanol or formamide, may also be added. Carriers, such as oils, optionally with solubilizing excipients, are especially suitable. Oils include any natural or synthetic non-ionic water-immiscible liquid, or low melting solid, which is capable of dissolving lipophilic compounds. Natural oils, such as triglycerides are representative. In fact, another aspect of this invention is a pharmaceutical composition comprising a compound of formula (I) and an oil.

Solid carriers include starch, lactose, calcium sulfate dihydrate, terra alba, magnesium stearate or stearic acid, talc, pectin, acacia, agar or gelatin. Solubilizing agents, such as dimethylsulfoxide or formamide, may also be added.

The carrier may also include a sustained release material such as glyceryl monostearate or glyceryl distearate, alone or with a wax. The amount of solid carrier varies but, preferably, will be between about 20 mg to about 1 g per dosage unit. The pharmaceutical preparations are made

following the conventional techniques of pharmacy involving milling, mixing, granulating, and compressing, when necessary, for tablet forms; or milling, mixing and filling for hard gelatin capsule forms. When a liquid carrier is used, the preparation will be in the form of a syrup, elixir, emulsion or an aqueous or non-aqueous suspension. Such a liquid formulation may be administered directly p.o. or filled into a soft gelatin capsule.

A suitable dosage form for oral administration has been prepared by dissolving the peptide of Example 1 (312.5 mg) in dimethyl sulfoxide (1 mL) and diluting to a concentration of 12.5 mg/mL with soybean oil. A suitable dosage form for intravenous administration has been prepared by dissolving the compound of Example 1 (0.02 g) in dimethyl sulfoxide (1 mL) and diluting to 20 mL with a 70% propylene glycol/30% ethanol solution.

For rectal administration, a pulverized powder of the compounds of this invention may be combined with excipients such as cocoa butter, glycerin, gelatin or polyethylene glycols and molded into a suppository. The pulverized powders may also be compounded with an oily preparation, gel, cream or emulsion, buffered or unbuffered, and administered through a transdermal patch.

The pharmacological activity of the compounds of this invention may be demonstrated by enzyme assays to determine the inhibitory activity of the retroviral protease, by *in vitro* cellular-based assays to determine the ability of the compounds to penetrate cells and inhibit viral replication, and by pharmacokinetic assays to determine oral bioavailability, drug half-life and clearance. These assays are well known in the art.

ENZYME ACTIVITY

The ability of the compounds of this invention to inhibit the HIV-1 protease enzyme may be demonstrated by using the assay disclosed by Dryer et al., *Proc. Natl. Acad. Sci., U.S.A.*, 86, 9752 (1989), Grant et al., *Biochemistry*, 30 8441 (1992), and EP-A 352 000. The K_i for the compounds of

this invention are in the range of 1 nM to 5 μ M. Preferred compounds have K_i 's of less than 100 nM.

INFECTIVITY

The ability of the compounds of this invention to gain entry to cells infected with the human immunodeficiency virus, and to inhibit viral replication *in vitro* may be demonstrated using the assay described by Meek *et al.*, *Nature*, 343, 90 (1990), and Petteway *et al.*, *Trends Pharmacol. Sci.*, 12, 28 (1991). The IC_{50} for the compounds of this invention are in the range of 0.1 to 10 μ M.

CYTOTOXICITY

Cytotoxicity is assessed by both direct microscopic examination of trypan blue stained cells (T-lymphocytes) and by the treated culture's ability to metabolize the tetrazolium salt XTT (2,3-bis[2-methoxy-4-nitro-5-sulfophenyl]-2H-tetrazolium-5-carboxanilide sodium salt), to its formazan dye. The XTT assay allows determination of the 50% toxic concentration of compounds for the cell/virus system used.

PHARMACOKINETICS

Dual jugular cannulated Sprague Dawley rats weighing 200 to 250 g were utilized in all studies. All dosing and sample collection was done from conscious rats. Before dosing, a time 0 blood sample, 300 μ L, was drawn using one of the catheters. Utilizing the second catheter the rats were dosed intravenously. At 1, 10, 30, 60, 90, 120, 150, 180 and 210 min after dosing, 300 μ L blood samples were drawn. The rats dosed orally were administered the compound by utilizing a 22 gauge gastric gavage needle and samples were drawn at 30, 60, 90, 120, 240, 360, 480, 600, 720 and 1440 min. The blood samples were placed in precooled tubes containing 30 mL of sodium citrate and centrifuged in a microfuge. The plasma was transferred then snap frozen on dry ice, and stored at -70°C until analyzed.

- 30 -

Standard stock solutions (1 mg/mL) of inhibitor was prepared in 100% DMSO. A dilution series of the stock solutions were prepared in a total volume of 0.1 mL (pooled normal rat plasma/DMSO) to yield final concentrations of 0 and 0.5-120X the K_i of the inhibitor. All dilutions were performed in triplicate. These spiked plasma solutions were extracted with 0.5 mL acetonitrile by vigorous vortexing, followed by centrifugation for 10 min. An aliquot (0.4 mL) of the supernatant was removed and dried in Eppendorf tubes using a Speed-vac. The resulting residue was redissolved in DMSO. The inhibition of the HIV-1 protease activity was assayed as follows. An aliquot of the extracted sample was added to a 50 μ L mixture containing 1X MENDT buffer, 1 mM substrate and incubated at 37°C < 10 min. The reaction was then initiated by the addition of HIV-1 protease and continued at 37°C for an additional 15 min, then quenched by the addition of TFA (0.5% final concentration). Initial rates were determined for each standard curve as the fraction of remaining enzymatic activity (v_i/v_0) at each inhibitor concentration, in which v_0 is the velocity of the I_0 (inhibitor concentration)=0 sample. Assuming that all of the original inhibitor in the spiked samples was extracted, the values of v_i/v_0 were plotted versus inhibitor concentration of the original extracted sample and fitted to the equation:

$$v_i/v_0 = [AE_t - I_t - K_i + (K_i - AE_t - I_t)^{0.5}] / (2AE_t)$$

in which E_t is the total enzyme concentration at time zero, K_i is the apparent inhibition constant and A is the fraction of active enzyme.

Ex vivo animal plasma samples containing unknown levels of protease inhibitor were prepared and analyzed as described for the standard curve described above. The concentration of inhibitor in these samples was then determined using the K_i and A parameters from the fitted standard curve according to the following equation:

$$I_t = AE_t [1 - (v_i/v_0)] + K_i (v_0/v_i)$$

The data was plotted as the natural log (ln) of the plasma concentration versus time on semilogarithmic paper to generate the plasma concentration-vs-time curves. Using the IV data, the apparent terminal rate constant was determined

of the linear regression analysis of the plasma concentration-vs-time curve. The elimination half-life ($t_{1/2}$) was derived by dividing $\ln 0.5$ ($=0.693$) by the terminal rate constant. The area under the plasma concentration-vs-time curve (AUC) was determined by using the ln/log trapezoidal rule. C_{max} represents the maximal plasma concentration and t_{max} , the time following drug administration at which C_{max} was observed. Both values were estimated by inspection of the plasma concentration-vs-time curve. Total plasma clearance (CL) was calculated by dividing the dose by the AUC. The fraction of the oral dose available to the systemic circulation (the bioavailable fraction, F) was determined by the equation: $F = [AUC_{po}/DOSE_{po}] \times [DOSE_{iv}/AUC_{iv}]$.

The Examples which follow serve to illustrate this invention. The Examples are not intended to limit the scope of this invention, but are provided to show how to make and use the compounds of this invention. In the Examples, all temperatures are in degrees Centigrade. Mass spectra were performed using fast atom bombardment (FAB) or electro-spray (ES) ionization. Melting points were taken on a Thomas-Hoover capillary melting point apparatus and are uncorrected.

NMR were recorded at 250 MHz using a Bruker AM 250 spectrometer, unless otherwise indicated. Chemical shifts are reported in ppm (δ) downfield from tetramethylsilane. Multiplicities for NMR spectra are indicated as: s=singlet, d=doublet, t=triplet, q=quartet, m=multiplet, dd=doublet of doublets, dt=doublet of triplets etc. and br indicates a broad signal. J indicates the NMR coupling constant in Hertz.

Celite® is filter aid composed of acid washed diatomaceous silica manufactured by Mansville Corp., Denver, Colorado. Florisil® is an activated magnesium silicate chromatographic support and is a registered trademark of Floridon Co., Pittsburgh, Pennsylvania. Sat. indicates a

- 32 -

saturated solution, eq indicates the proportion of a molar equivalent of reagent relative to the principal reactant.

Example 1

5

Preparation of (2R,4S,5S,1'S)-2-phenylmethyl-4-hydroxy-5-(t-butoxycarbonyl)amino-6-phenyl-N-(1'-isopropyl-1'-(imidazo-2-yl)methyl-hexanamide hydrochloride

10 a) (1'S)-1'-carbobenzyloxyamino-1'-isopropyl-1'-(imidazo-2-yl)methane

Cbz-valinal (4.6 g, 1 eq) and glyoxal trimeric dihydrate (1.33 g, 1eq) were stirred in MeOH at -10°C. Ammonia was bubbled through the solution for several min and the mixture
15 was allowed to stir for 4 h at -10°C. The mixture was allowed to warm to room temperature over 14 h, then was poured into 250 mL water. The suspension was filtered and the filter cake washed twice with water to give the title compound as a white solid (1.9 g, 36%). NMR(CD₃OD) δ 7.28

20 (5H, m), 6.89 (2H, s), 5.04 (2H, dd), 4.46 (1H, d), 2.10 (1H, m), 0.91 (3H, d), 0.70 (3H, d); MS(CI/CH₄) m/e 274.2, [M+H]⁺, 230.1, 166.1, 123.1, 91.1.

b) (1'S)-1'-amino-1'-isopropyl-1'-(imidazo-2-yl)methane

25 (1'S)-1'-carbobenzyloxyamino-1'-isopropyl-1'-(imidazo-2-yl)methane (1.9 g) was stirred in methanol over 10% Pd/C (200 mg). Hydrogen was bubbled through the solution for 1 h and the solution was maintained under a positive hydrogen atmosphere overnight. The mixture was filtered through
30 Celite® and was evaporated to a tacky solid (720 mg, 75%). NMR(CDC₃Cl) δ 6.87 (2H, s), 3.88 (1H, d), 2.04 (1H, m), 0.81 (6H, dd); MS(DCI/NH₃) m/e 190.2 [M+H]⁺.

c) (2R,4S,5S,1'S)-2-phenylmethyl-4-(t-butyl dimethyl)siloxy-5-(t-butoxycarbonyl)amino-6-phenyl-N-[1'-isopropyl-1'-(imidazo-2-yl)methyl-hexanamide

To a solution of (2R,4S,5S)-2-phenylmethyl-4-(t-butylmethyl)siloxy-5-(t-butoxycarbonyl)amino-6-phenyl-

- 33 -

hexanoic acid (200 mg, 0.38 mmol) in dichloromethane, (1'S)-1'-amino-1'-isopropyl-1'-(imidazo-2-yl)methane (48 mg, 0.35 mmol), BOP reagent (168 mg, 0.38 mmol), and triethylamine (0.053 mL, 0.38 mmol) were added. The mixture was stirred under argon overnight, and washed successively with water, 5% aqueous sodium bicarbonate, and saturated aqueous sodium chloride. The solution was dried over MgSO_4 , filtered, and evaporated to a white solid. The solid was chromatographed (silica, 4% methanol/dichloromethane) to afford the title compound as a white solid (0.154 g, 68%). NMR(CDCl_3) δ 7.18 (10H, m), 6.91 (2H, d), 6.32 (1H, d), 4.69 (1H, d), 4.40 (1H, t), 3.92 (1H, q), 3.63 (1H, m), 2.84 - 2.31 (6H, m), 1.67 (4H, m), 1.24 (9H, s), 0.89 (9H, s), 0.74 (6H, dd), 0.05 (6H, d); MS(DCI/ NH_3) m/e 649.6 $[\text{M}+\text{H}]^+$.

(2R,4S,5S,1'S)-2-phenylmethyl-4-hydroxy-5-(t-butoxycarbonyl)amino-6-phenyl-N-[1'-isopropyl-1'-(imidazo-2-yl)]methyl-hexanamide hydrochloride

The compound of Example 1(c) (0.140 g) was stirred in THF at room temperature under an argon atmosphere. Tetrabutyl ammonium fluoride (0.38 mL, 6 eq) was added and the solution was stirred overnight. The mixture was diluted with water and extracted with dichloromethane (3X). The combined organic extracts were washed with water and evaporated. The residue was treated with 1 eq of methanolic HCl, concentrated, and triturated with diethyl ether and diethyl acetate to give the title compound as a white solid (95 mg, 83%). NMR($\text{DMSO}-d_6$) δ 7.78 (1H, d), 7.16 (10H, m), 6.71 (2H, s), 6.39 (1H, d), 4.68 (1H, m), 4.52 (1H, d), 2.71 (3H, m), 2.48 (3H, m), 1.97 (1H, m), 1.61 (1H, m), 1.30 (9H, s), 0.78 (3H, d), 0.61 (3H, d); MS(DCI/ NH_3) m/e 535.4 $[\text{M}+\text{H}]^+$.

Example 2

Preparation of (2R,4S,5S,1'S)-2-phenylmethyl-4-hydroxy-5-(t-butoxycarbonyl)amino-6-phenyl-N-[1'-isopropyl-1'-(4-aminocarbonyl-thiazo-2-yl)]methyl-hexanamide

- 34 -

a) Boc-valin amide

To a solution of di-*t*-butyl-dicarbonate (7.15 g, 1 eq) in dry dichloromethane was added valinamide hydrochloride (5.0 g, 1 eq) and triethylamine (9.14 mL, 2 eq). The mixture was heated to reflux for 4 h, and cooled to room temperature. The organic layer was washed twice with water and evaporated to give the title compound (6.03 g, 85%). NMR(CDCl₃) δ 6.00 (1H, br), 5.54 (1H, br), 5.01 (1H, br), 3.93 (1H, dd), 2.12 (1H, m), 1.44 (9H, s), 0.92 (6H, dd).

b) Boc-valinethioamide

Boc-valineamide (0.5 g) was stirred in dry THF at room temperature under argon. Lawesson's reagent (1.56 g, 0.6 eq) was added and the mixture was stirred overnight. The solvent was evaporated and the residue chromatographed (silica, 2.5% methanol/dichloromethane) to give the title compound as a white solid (0.373 g, 70%). NMR(CDCl₃) δ 8.59 (1H, br s), 8.09 (1H, br s), 5.41 (1H, d (br)), 4.20 (1H, dd), 1.99 (1H, m), 1.39 (9H, s), 0.90 (6H, m).

c) (1'S)-1'-(*t*-butoxycarbonyl)amino-1'-isopropyl-1'-(4-carboethoxythiazo-2-yl)methane

Boc-valinethioamide (0.265 g) was stirred in dry acetone under argon at -10°C. Ethylbromopyruvate (0.16 mL, 1.1 eq) was added and stirred for 1 h at -10°C. The solution was poured into a well-stirred mixture of chloroform and water and then saturated with sodium bicarbonate. The organic phase was separated and the aqueous layer extracted with chloroform. The combined organic extracts were dried over MgSO₄, filtered, and evaporated to an oil. The oily residue was treated with trifluoroacetic anhydride (0.16 g) and pyridine (0.2 g) in dichloromethane for 1 h at -20°C. Excess solvent was removed in vacuo and the residue dissolved in dichloromethane. The solution was washed with sat. aqueous sodium bicarbonate and 1.0N KHSO₄ until pH 7. The solution was dried over sodium sulfate, filtered, and evaporated to an oil which was chromatographed (silica, 4% methanol/dichloromethane) to give the title compound as a tan solid.

- 35 -

¹H NMR(CDCl₃) δ 8.04 (1H, s), 5.26 (1H, br d), 4.85 (1H, m), 4.37 (2H, q), 2.40 (1H, m), 1.41 (9H, s), 1.34 (3H, t), 0.93 (3H, d), 0.84 (3H, d).

d) (1'S)-1'-(t-butoxycarbonyl)amino-1'-isopropyl-1'-(4-carboxythiazo-2-yl)methane

The compound of Example 2(c) (50 mg) was stirred in THF at 0°C. Excess 1.0N NaOH was added and the mixture was stirred for 12 h at 0°C. The mixture was diluted with 1.0N citric acid and extracted with dichloromethane (3X). The combined organic extracts were evaporated and dried in vacuo to give the title compound (0.045 g, 98%). ¹H NMR(CDCl₃) δ 8.08 (1H, s), 5.19 (1H, m), 4.80 (1H, m), 2.31 (1H, m), 1.38 (9H, s), 0.86 (6H, dd).

e) (1'S)-1'-(t-butoxycarbonyl)amino-1'-isopropyl-1'-(4-aminocarbonylthiazo-2-yl)methane

(1'S)-1'-(t-butoxycarbonyl)amino-1'-isopropyl-1'-(4-carboxythiazo-2-yl)methane (0.078 g, 0.26 mmol) was stirred under argon in dry THF at -40°C. NMM (0.06 mL; 0.55 mM) and isobutyl chloroformate (0.034 mL; 0.26 mmol) were added. After stirring 15 min, ammonia was bubbled through the mixture for several min. The solution was warmed to room temperature and the THF evaporated. The residue was diluted with ethyl acetate and washed successively with 1.0N citric acid, 5% aqueous sodium bicarbonate, and sat. aqueous sodium chloride. The organic layer was dried over MgSO₄, filtered, and evaporated to a solid which was chromatographed (silica, 3% methanol/dichloromethane) to give the title compound as a white solid (0.052 g, 67%). ¹H NMR(CDCl₃) δ 8.02 (1H, s), 7.14 (1H, s(br)), 6.28 (1H, s(br)), 5.24 (1H, d(br)), 4.82 (1H, m), 2.30 (1H, m), 1.39 (9H, s), 0.92 (6H, dd).

f) (2R,4S,5S,1'S)-2-phenylmethyl-4-(t-butyldimethylsiloxy)-5-(t-butoxycarbonyl)amino-6-phenyl-N-[1'-isopropyl-1'-(4-aminocarbonylthiazo-2-yl)]methyl-hexanamide

The compound of Example 2(e) (52 mg) was stirred in neat trifluoroacetic acid for 10 min and evaporated. The residue

- 36 -

was diluted with methanol and treated with 2 eq of conc. HCl. The solvents were evaporated and dried in vacuo to give a white solid. This solid (40 mg) was added to a solution of (2R,4S,5S)-2-phenylmethyl-4-(t-butyldimethyl)siloxy-5-(t-butoxycarbonyl)amino-6-phenyl-hexanoic acid (97 mg, 1.1 eq), DCC (38 mg, 1.1 eq), and HOBT (0.05 g, 2.2 eq) in DMF at room temperature under argon. N-methylmorpholine (0.04 mL; 2.2 eq) was added and the mixture was stirred overnight. The mixture was filtered through Celite®, evaporated, and diluted with ethyl acetate. The solution was washed successively with 1.0N citric acid, 5% aqueous sodium bicarbonate, and sat. aqueous sodium chloride. The organic layer was chromatographed (silica, 2.5% methanol/dichloromethane) to yield the title compound (60 mg, 55%). NMR(CDCl₃) δ 7.89 (1H, s), 7.60 (1H, d), 7.24 (10H, m), 6.82 (1H, m), 5.12 (1H, m), 4.89 (1H, m), 3.92 (1H, q), 3.81 (1H, dd), 2.73 (4H, m), 2.21 (1H, m), 1.73 (2H, m), 1.40 (9H, s), 1.23 (1H, m), 0.93 (9H, s), 0.84 (6H, dd), 0.11 (6H, d).

(20 g) (2R,4S,5S,1'S)-2-phenylmethyl-4-hydroxy-5-(t-butoxycarbonyl)amino-6-phenyl-N-[1'-isopropyl-1'-(4-aminocarbonyl(thiazo-2-yl)]methyl-hexanamide

The compound of Example 2(f) (60 mg) was stirred in dry THF under argon and tetrabutylammonium fluoride (0.50 mL, 6 eq) was added. The solution was stirred at room temperature overnight. After diluting with water, the aqueous layer was extracted with dichloromethane (3X). The combined organic extracts were washed with water, evaporated, and triturated with diethyl ether and ethyl acetate to give a tan solid. The solid was chromatographed (silica gel, 4% methanol/dichloromethane) to give the title compound as a white solid (0.022 g). NMR(CDCl₃) δ 7.90 (1H, s), 7.15 (10H, m), 6.39 (1H, d), 5.93 (1H, br s), 5.06 (1H, dd), 4.91 (1H, d), 3.90 (1H, d), 3.67 (2H, m), 2.91 (4H, m), 2.64 (1H, d), 2.13 (1H, m), 1.87 (3H, m), 1.36 (9H, s), 0.83 (6H, dd); MS(DCI/NH₃) m/e 612 [M+NH₄]⁺, 595 [M+H]⁺, 495, 413.1, 391, 374, 356, 239.1, 202, 185.

Example 3Preparation of (2R,4S,5S,1'S)-2-phenylmethyl-4-hydroxy-5-(t-butoxycarbonyl)amino-6-phenyl-N-[1'-isopropyl-1'-(thiazo-2-yl)]methyl-hexanamide

a) (1'S)-1'-(t-butoxycarbonyl)amino-1'-isopropyl-1'-(thiazo-2-yl)methane

The compound of Example 2(c) was stirred in neat quinoline. Cu powder (0.50 g) was added and the suspension was heated to 160°C for 2 h. After cooling to room temperature, the solution was diluted with ethyl acetate and washed with 2.0N citric acid (4X). The organic layers were combined and dried over MgSO₄, filtered, and evaporated to a dark oil. The oil was chromatographed (silica, 4% methanol/dichloromethane) to give the title compound as an orange oil.

NMR(CDCl₃) δ 7.68 (1H, d), 7.19 (1H, d), 5.26 (1H, d), 4.88 (1H, m), 2.31 (1H, m), 1.43 (9H, s), 0.92 (3H, d), 0.84 (3H, d).

20

b) (2R,4S,5S,1'S)-2-phenylmethyl-4-hydroxy-5-(t-butoxycarbonyl)amino-6-phenyl-N-[1'-isopropyl-1'-(thiazo-2-yl)]methyl-hexanamide

Following the procedure of Example 2(f)-2(g), except using the compound of Example 3(a) in place of (1'S)-1'-(t-butoxycarbonyl)amino-1'-isopropyl-1'-(4-aminocarbonylthiazo-2-yl)methane, the title compound was prepared (88%).

NMR(DMSO-d₆) δ 8.31 (1H, d), 7.62 (1H, d), 7.49 (1H, d), 7.16 (10H, m), 2.61 (6H, m), 1.28 (9H, s), 0.89 (3H, dd);

MS(DCI/NH₃) m/e 552.3 [M+H]⁺, 413.2, 331.1, 183.1, 157.1,

142.0, 120.1.

Example 4Preparation of (2R,4S,5S,1'S)-2-phenylmethyl-4-hydroxy-5-(t-butoxycarbonyl)amino-6-phenyl-N-[1'-isopropyl-1'-benzimidazo-2-yl]]methyl-hexanamide

- 38 -

a) (1'S)-1'-carbobenzyloxyamino-1'-isopropyl-1'-(benzimidazo-2-yl)methane

Cbz-valine (2.0 g, 1 eq) was stirred at -10°C in dry THF under argon. Triethylamine (1.11 mL, 1.0 eq) was added, followed by isobutyl chloroformate (1.03 mL, 1 eq). The reaction mixture was stirred for 10 min. Phenylene diamine (0.944 g, 1.1 eq) was added slowly in 10 mL dry THF. The mixture was warmed to room temperature and stirred for 1 h. The solvents were evaporated and the residue partitioned between water and ethyl acetate. The ethyl acetate layer was washed with 5% aqueous sodium bicarbonate and brine. The organic layer was dried over MgSO₄, filtered, and evaporated. The residue was dissolved in glacial acetic acid and heated to 65°C for 16 h. The solvents were evaporated and the residue diluted with water. After neutralizing with saturated aqueous sodium bicarbonate, the solid was filtered (and the filter cake was washed with hexane). The solid was recrystallized from ethyl acetate and hexane. NMR(CD₃OD) δ 7.48-7.11 (9H, m), 5.06 (2H, q), 4.62 (1H, m), 2.27 (1H, m), 1.23 (1H, m), 1.02 (3H, d), 0.84 (3H, d).

b) (1'S)-1'-amino-1'-isopropyl-1'-(benzimidazo-2-yl)methane

The compound of Example 4(a) (2.76 g) was stirred in methanol. 10% palladium on activated carbon (Pd/C) (250 mg) was added and hydrogen gas was bubbled through the solution for 1 h. The reaction was maintained under an hydrogen atmosphere overnight. The mixture was filtered through Celite® and the solvents evaporated to give the title compound as a white solid (1.58 g, 98%). NMR(CDCl₃) δ 7.48-7.10 (4H, m), 4.02 (1H, d), 2.24 (1H, m), 0.96 (3H, d), 0.83 (3H, d); MS(DCI/NH₃) m/e 190.2 [M+H]⁺.

c) (2R,4S,5S,1'S)-2-phenylmethyl-4-(t-butyldimethyl)siloxy-5-(t-butoxycarbonyl)amino-6-phenyl-N-[1'-isopropyl-1'-benzimidazo-2-yl]methyl-hexanamide

To a solution of (2R,4S,5S)-2-phenylmethyl-4-(t-butyldimethyl)siloxy-5-(t-butoxycarbonyl)amino-6-phenyl-hexanoic acid (75 mg, 1.1 eq) in dimethyl formamide under

- 39 -

argon, the compound of Example 4(b) (25 mg, 1.0 eq), DCC (30 mg, 1.1 eq) and HOBT (44 mg, 2.2 eq) were added. The mixture was stirred overnight, then filtered through Celite®. The solvents were evaporated and the residue was chromatographed (silica gel, 4% methanol/dichloromethane) to give the title compound (0.070 g, 78%). NMR(CDCl₃) δ 7.88 (1H, d), 7.30 (14H, m), 6.80 (1H, d), 4.93 (2H, m), 4.26 (1H, q), 4.00 (1H, m), 2.92 (7H, m), 2.01 (2H, m), 1.53 (9H, s), 1.20 (9H, s), 1.14 (6H, d), 0.41 (6H, d); MS(DCI/NH₃) m/e 699.6 [M+H]⁺.

d) (2R,4S,5S,1'S)-2-phenylmethyl-4-hydroxy-5-(t-butoxycarbonyl)amino-6-phenyl-N-[1'-isopropyl-1'-benzimidazo-2-yl]methyl-hexanamide

15 The compound of Example 4(c) was stirred in dry THF and tetrabutyl ammonium fluoride (0.6 mL, 6 eq) was added. The mixture was stirred under argon overnight at room temperature. The solution was diluted with water and extracted with dichloromethane (3X). The combined organic layers were washed with water and evaporated to a residue which was chromatographed (silica, 2% methanol/CH₂Cl₂) to give the title compound (0.029 g, 50%). NMR(CDCl₃) δ 7.54 (1H, m), 7.11 (11H, m), 6.69 (4H, s), 4.98 (1H, d), 4.69 (2H, m), 3.66 (2H, m), 2.74 (5H, m), 2.31 (1H, m), 1.73 (2H, m), 1.32 (9H, s), 0.70 (6H, d); MS(DCI/NH₃) m/e 585.4 [M+H]⁺, 413.3, 364.3, 296.2, 190.2, 173.1, 120.1.

Example 5

30 Preparation of (2R,4S,5S,1'S)-2-phenylmethyl-4-hydroxy-5-(t-butoxycarbonyl)amino-6-phenyl-N-(1'-imidazo-2-yl)methyl-hexanamide hydrochloride

a) 2-(carbobenzyloxyamino)methyl-imidazole

35 Following the procedure of Example 1(a), except substituting Cbz-glycinal for Cbz-valinal, the title compound was prepared. NMR(CDCl₃) δ .33 (5H, s), 6.95 (2H, s), 5.95

- 40 -

(1H, s(br)), 5.12 (2H, s), 4.42 (2H, d); MS(DCI/NH₃) m/e 232.2 [M+H]⁺, 188, 171.

b) (2R,4S,5S,1'S)-2-phenylmethyl-4-hydroxy-5-(t-butoxycarbonyl)amino-6-phenyl-N-(1'-imidazo-2-yl) methyl-hexanamide hydrochloride

Following the procedure of Example 1(b)-1(d), except substituting the compound of Example 5(a) for (1'S)-1'-carbobenzyloxyamino-1'-isopropyl-1'-(imidazo-2-yl)methane, the title compound was prepared. NMR(CD₃OD) δ 7.20 (10H,m), 6.94 (2H,s), 6.11 (1H,d), 4.24 (2H,dd), 3.61 (1H,m), 3.52 (1H,m), 2.69 (4H,m), 1.66 (2H,m), 1.28 (9H,s); MS (DCI/NH₃) m/e 493.7 [M+H]⁺, 475.7, 120.2, 98.2, 83.1, 69.1.

15

Example 6

Preparation of (2R,4S,5S,1'S)-2-phenylmethyl-4-hydroxy-5-(t-butoxycarbonyl)amino-6-phenyl-N-[1'-methyl-1'-(imidazo-2-yl)] methyl-hexanamide hydrochloride

20

a) (1'S)-1'-carbobenzyloxyamino-1'-methyl-1'-(imidazo-2-yl)methane

Following the procedure of Example 1(a), except substituting Cbz-alanal for Cbz-valinal, the title compound was prepared. NMR(CDCl₃) δ .35 (5H,s), 6.92 (2H,s), 5.52 (1H,d), 5.12 (2H,q), 4.90 (1H,q); MS(DCI/NH₃) m/e 246 [M+H]⁺, 202, 185.

b) (2R,4S,5S,1'S)-2-phenylmethyl-4-hydroxy-5-(t-butoxycarbonyl)amino-6-phenyl-N-[1'-methyl-1'-(imidazo-2-yl)] methyl-hexanamide hydrochloride

Following the procedure of Example 1(b)-1(d), except substituting the compound of Example 6(a) for (1'S)-1'-carbobenzyloxyamino-1'-isopropyl-1'-(imidazo-2-yl)methane, the title compound was prepared. NMR(CD₃OD) δ 7.11 (10H,m), 6.86 (2H, s), 4.69 (1H, d), 3.62 (1H, d), 3.51 (1H, m), 2.68 (6H, m), 1.59 (2H, m), 1.30 (9H, s), 1.14 (3H, d); MS(DCI/NH₃) m/e 507.5 [M+H]⁺, 489.4, 112.1.

- 41 -

Example 7

Preparation of (2R,4S,5S,1'S)-2-phenylmethyl-4-hydroxy-5-(t-butoxycarbonyl)amino-6-phenyl-N-(1'-benzyl-1'-(imidazo-2-yl)methyl-hexanamide hydrochloride.

a) (1'S)-1'-carbobenzyloxyamino-1'-benzyl-1'-(imidazo-2-yl)methane

10 Following the procedure of Example 1(a), except substituting Cbz-phenylalaninal for Cbz-valinal, the title compound was prepared. NMR(CDCl₃) δ 7.37-7.05 (10H, m), 6.95 (2H, s br), 5.52 (1H, d), 5.05 (2M, s), 4.95 (1H, q), 3.32 (2H, d); MS(DCI/NH₃) m/e 322, 261, 171.

15

b) (2R,4S,5S,1'S)-2-phenylmethyl-4-hydroxy-5-(t-butoxycarbonyl)amino-6-phenyl-N-(1'-benzyl-1'-(imidazo-2-yl)methyl-hexanamide hydrochloride

Following the procedure of Example 1(a)-1(d), except substituting the compound of Example 7(a) for (1'S)-1'-carbobenzyloxyamino-1'-isopropyl-1'-(imidazo-2-yl)methane, the title compound was prepared. NMR(CD₃OD) δ 7.15 (15H, m), 6.79 (2H, s), 5.78 (1H, d), 5.04 (1H, d), 3.58 (1H, m), 3.47 (1H, m), 2.68 (8H, m), 1.59 (2H, m), 1.31 (9H, s).

25 and the usual workup, oil, b.p. 100-105°C/0.5 mm Hg.

Example 8

Preparation of (2R,4S,5S,1'S)-5-(carbobenzyloxy)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide

A solution of (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide (0.086 g) in trifluoroacetic acid was stirred for 10 min, then was evaporated in vacuo. To the residue were added dimethylformamide, benzylchloroformate (1 eq) and triethylamine (5 eq), and the resulting mixture was stirred at room temperature for 16 h. The reaction mixture

was poured into H₂O and extracted with dichloromethane. The combined organic extracts were evaporated, and the residue was triturated with diethyl ether to afford the title compound as a white solid. NMR(CD₃OD) δ 7.36-6.94 (15H, m), 6.84 (2H, s), 4.99 (2H, s), 4.54 (2H, d), 3.76 (1H, m), 3.52 (1H, dd), 2.77 (5H, m), 2.04 (1H, m), 1.76 (1H, m), 1.58 (1H, m), 0.82 (3H, d), 0.66 (3H, d).

Example 9

10

Preparation of (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-[1'-isopropyl-1'-(4,5-dimethyl)imidazol-2-yl]methyl-6-phenyl-2-phenylmethyl-hexanamide

15 a) (1S)-1-carbobenzyloxyamino-1-isopropyl-1-(4,5-dimethylimidazol-2-yl)methane

Cbz-Valinal (4.14 g) was stirred in methanol with 2,3-butanedione (1.54 mL, 1.0 eq). Ammonia was bubbled through the solution at -25°C for 5 min. The cooling bath was removed and the mixture allowed to warm to 20°C. The solution was stirred for 16 h under Ar. The solvents were removed by rotary evaporation, and the residue was diluted with dichloromethane and extracted with dilute aqueous HCl. The organic layer was concentrated to afford unreacted Cbz-valinal (4.02 g). The acidic aqueous layer was basified with 1N NaOH and extracted with dichloromethane, the organic extract was concentrated and the residue purified by flash chromatography (4% methanol in dichloromethane) to provide the title compound as a white solid (50 mg). NMR(CD₃OD) δ 7.29 (5H, m), 5.04 (2H, dd), 4.38 (1H, d), 2.06 (6H, s), 2.01 (1H, m), 0.93 (3H, d), 0.77 (3H, d).

b) (1S)-1-(4,5-dimethylimidazol-2-yl)-2-methylpropylamine

The benzyloxycarbonyl group was cleaved by hydrogenolysis using the same procedure as described previously in Example 1(b), except using the product of 1(a) (50 mg), to afford the title compound as a white solid

(24 mg, 87%). NMR(CDCl₃) δ 4.11 (2H, s(br)), 3.71 (1H, d), 2.06 (6H, s), 2.00 (1H, m), 0.71 (6H, dd).

c) (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-t-butyl-
5 butyldimethylsiloxy-N-[1'-isopropyl-1'-(4,5-dimethyl)
imidazol-2-yl]methyl-6-phenyl-2-phenylmethyl-hexanamide

Using the procedure of Example 1(c), except substituting (2R,4S,5S)-5-(t-butoxycarbonyl)amino-4-t-butyl-
6-phenyl-2-phenylmethylhexanoic acid and (1S)-1-(4,5-
10 dimethylimidazol-2-yl)-2-methylpropylamine (24 mg), the title compound was prepared (55 mg, 57%). NMR(CDCl₃) δ 7.26-6.80
(10H, m), 4.65 (1H, d), 4.24 (1H, dd), 3.87 (1H, q), 3.61
(1H, m), 2.77-2.39 (5H, m), 2.22 (1H, m), 1.98 (6H, s), 1.79
(1H, m), 1.58 (1H, m), 1.24 (9H, s), 0.85 (9H, s), 0.69 (6H,
15 d), 0.06 (6H, d).

d) (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-[1'-
isopropyl-1'-(4,5-dimethyl)imidazol-2-yl]methyl-6-phenyl-2-
phenylmethyl-hexanamide

20 By following the deprotection procedure described in
Example 1(d), except using (2R,4S,5S,1'S)-5-(t-
butoxycarbonyl)amino-4-t-butyl-
butyldimethylsiloxy-N-[1'-isopropyl-
1'-(4,5-dimethyl)imidazol-2-yl]methyl-6-phenyl-2-
phenylmethyl-hexanamide (55 mg) and omitting the final
25 treatment with methanolic HCl, the title compound was
prepared (25 mg, 62%). NMR(CDCl₃) δ 7.29-6.88 (10H, m), 4.98
(1H, br d), 4.47 (1H, m), 4.29 (1H, m), 3.58 (2H, m), 2.84-
2.51 (5H, m), 2.20 (1H, m), 2.04 (6H, s), 1.71 (2H, m), 1.38
(9H, s), 0.69 (6H, dd).

30

Example 10

Preparation of (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-
hydroxy-N-[1'-isopropyl-1'-(4,5-dimethyl)imidazol-2-
35 yl]methyl-6-phenyl-2-phenylmethyl-hexanamide

a) (1S)-1-carbobenzyloxyamino-1-isopropyl-1-(4-
phenylimidazol-2-yl)methane

- 44 -

Using the procedure of Example 1(a), except using Cbz-(L)-valine (2.19 g) and α -ketophenylacetaldehyde instead of glyoxal, the title compound was prepared (1.54 g, 48%).
NMR(CDCl₃) δ 7.62 (1H, (br)), 7.24 (10H, m), 5.79 (1H, d),
5 5.04 (2H, dd), 4.32 (1H, dd), 2.31 (1H, m), 0.96 (3H, d),
0.79 (3H, d); MS m/e 350.4 [M+H]⁺, 199.0.

b) (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-t-butyl-
butyldimethylsiloxyl-N-[1'-isopropyl-1'-(4-phenyl)imidazol-2-
10 yl]methyl-6-phenyl-2-phenylmethyl-hexanamide

Using the procedure of Example 1(b)-1(c), except using
the compound of 10(a) (72 mg), the title compound was
prepared (67 mg, 44%). NMR(CDCl₃) δ 7.70 (1H, d), 7.40-6.71
(16H, m), 4.73 (1H, d), 4.54 (1H, dd), 3.96 (1H, q), 3.69
15 (1H, m), 2.88-2.36 (5H, m), 1.73 (2H, m), 1.33 (9H, s), 0.91
(9H, s), 0.84 (6H, dd), 0.11 (6H, d); MS m/e 725.4 [M+H]⁺.

c) (2R, 4S, 5S, 1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-
[1'-isopropyl-1'-(4-phenyl)imidazol-2-yl]methyl-6-phenyl-2-
20 phenylmethyl-hexanamide

Using the procedure of Example 9(d), except starting
from (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-t-
butyldimethylsiloxyl-N-[1'-isopropyl-1'-(4-phenyl)imidazol-2-
yl]methyl-6-phenyl-2-phenylmethyl-hexanamide (67 mg), the
25 title compound was prepared (30 mg, 54%). NMR(CDCl₃) δ 7.52-
6.67 (16H, m), 5.48 (1H, d), 3.60 (1H, q), 3.44 (1H, d), 2.60
(4H, m), 1.96 (1H, m), 1.62 (2H, m), 1.23 (9H, s), 0.73 (3H,
d), 0.62 (3H, d); MS m/e 611.4 [M+H]⁺, 242.2, 195.0, 150.2.

30

Example 11

Preparation of (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-
hydroxy-N-[1'-isopropyl-1'-(N'-methyl)imidazol-2-yl]methyl-6-
35 phenyl-2-phenylmethyl-hexanamide

35

- 45 -

a) (1S)-carb benzyl xyamino-1-isopropyl-1-(N'-methylimidazol-2-yl)methane

The product of Example 1(a) (273 mg, 1 mmol) was heated at 40°C for 2 h in methyl iodide (5 mL). The reaction mixture was evaporated, and the residue was suspended in aqueous Na₂CO₃. The mixture was extracted with dichloromethane, dried (Na₂CO₃) and concentrated. The crude product was purified by flash chromatography (silica, 2% methanol/dichloromethane) to yield the title compound (200 mg, 70%). NMR(CDCl₃) δ 7.29 (5H, s), 6.92 (1H, s), 6.69 (1H, s), 5.94 (1H, d), 5.03 (2H, q), 4.55 (1H, dd), 3.64 (3H, s), 2.20 (1H, m), 1.01 (3H, d), 0.82 (3H, d).

b) (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-t-butylbutyldimethylsiloxy-N-[1'-isopropyl-1'-(N'-methyl)imidazol-2-yl]methyl-6-phenyl-2-phenylmethyl-hexanamide

Following the procedure of Example 1(b)-1(c), except using the compound of 11(a) (90 mg), the title compound was prepared (104 mg, 50%). NMR(CDCl₃) δ 7.32-6.89 (10H, m), 6.81 (1H, s), 6.59 (1H, s), 6.08 (1H, d), 4.71 (2H, m), 3.94 (1H, q), 3.70 (1H, m), 3.25 (3H, s), 2.80-2.36 (5H, m), 2.21 (1H, m), 1.73 (2H, m), 1.31 (9H, s), 0.94 (9H, s), 0.85 (6H, dd), 0.11 (6H, s).

c) (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-[1'-isopropyl-1'-(N'-methyl)imidazol-2-yl]methyl-6-phenyl-2-phenylmethyl-hexanamide

Following the procedure of Example 9(d), except using (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-t-butylbutyldimethylsiloxy-N-[1'-isopropyl-1'-(N'-methyl)imidazol-2-yl]methyl-6-phenyl-2-phenylmethyl-hexanamide (100 mg), the title compound was prepared (74 mg, 89%). NMR(CDCl₃) δ 7.21-6.74 (11H, m), 6.70 (1H, s), 6.59 (1H, s), 4.95 (1H, d), 4.61 (1H, dd), 3.60 (3H, m), 3.48 (3H, s), 2.71 (5H, m), 2.06 (1H, m), 1.64 (2H, m), 1.32 (9H, s), 0.82 (3H, d), 0.63 (3H, d);

MS m/e 549.3 [M+H]⁺.

Example 12**Preparation of (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-(3-phenylpropargyl)hexanamide****a) (3R,5S,1'S)-(1'-t-butoxycarbonylamino-2'-phenyl)ethyl-3-(3-phenylpropargyl)-tetrahydrofuran-2-one**

To a solution of lithium diisopropylamide (3.61 mL, 2.0 M in THF, 2.2 eq) in THF at -78°C under an argon atmosphere, (5S,1'S)-(1'-t-butoxycarbonylamino-2'-phenyl)ethyl-tetrahydrofuran-2-one (1.0 g, 1.0 eq) was added. After stirring at -78°C for 15 min, hexamethylphosphoramide (1.14 mL, 2 eq) was added, and stirring was continued an additional 10 min. Phenylpropargyl bromide (1.28 g, 2.0 eq), was added and the resulting mixture was stirred at -78°C for 2 h, then poured into dilute aqueous HCl and extracted with dichloromethane. The combined organic extracts were evaporated under reduced pressure to an oil, which was chromatographed (silica, 20% ethyl acetate/hexanes) to afford the title compound as a white solid (0.455 g, 33%).

NMR(CDCl₃) δ 7.18 (10H, m), 4.50 (2H, m), 3.93 (1H, q), 2.79 (5H, m), 2.23 (2H, m), 1.24 (9H, s).

b) (2R,4S,5S)-5-(t-butoxycarbonyl)amino-4-t-butyldimethylsiloxy-6-phenyl-2-(3-phenylpropargyl)hexanoic acid

The title compound (496 mg, 84%) was prepared by the procedure of Evans et al., *J. Org. Chem.* 50, 14615 (1985) from the product of 12(a) (450 mg). NMR(CDCl₃) δ 7.49-7.10 (10H, m), 4.71 (1H, d), 3.94 (3H, m), 2.69 (4H, m), 1.90 (2H, m), 1.31 (9H, s), 0.89 (9H, s), 0.11 (6H, d).

c) (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-t-butyldimethylsiloxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-(3-phenylpropargyl)hexanamide

Following the procedure of Example 1(c), except using (2R,4S,5S)-5-(t-butoxycarbonyl)amino-4-t-butyldimethylsiloxy-6-phenyl-2-(3-phenylpropargyl)hexanoic acid (240 mg) and

(1S)-1-imidazol-2-yl-2-methylpropylamine, the title compound was prepared (244 mg, 84%). NMR(CDCl₃) δ 7.14 (12H, m), 6.72 (1H, d), 4.58 (1H, d), 4.49 (1H, dd), 3.92 (1H, q), 3.80 (1H, m), 2.54 (5H, m), 1.65 (2H, m), 1.20 (9H, s), 0.81 (9H, s), 0.80 (6H, dd), 0.05 (6H, d).

d) (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-(3-phenylpropargyl)hexanamide

10 Following the procedure of Example 9(d), except using (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-t-

butyldimethylsiloxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-(3-phenylpropargyl)hexanamide, the title compound was prepared (161 mg, 79%). NMR(CDCl₃) δ 7.24-6.98 (10H, m), 6.68 (2H, s), 5.20 (1H, m), 4.52 (1H, d), 3.49 (2H, m), 3.06 (1H, m), 2.56 (5H, m), 2.04 (1H, m), 1.61 (2H, m), 1.26 (9H, s), 0.68 (6H, dd); MS m/e 581.2 (M+Na)⁺, 559.2 [M+H]⁺, 541.4, 503.2, 485.2, 459.2, 441.2.

20

Example 13

Preparation of (2R,4S,5S,1'S)-5-(isopropoxycarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide

a) (2R,4S,5S,1'S)-5-amino-4-t-butyldimethylsiloxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide
The product of Example 1(c) (0.20 g, 0.31 mmol) was dissolved in trifluoroacetic acid and stirred at room temperature for 5 min, and partitioned between dichloromethane and saturated aqueous Na₂CO₃. The organic extract was dried over Na₂CO₃, filtered and evaporated to afford the title compound (0.17 g, 100%) which was used without further purification.

b) (2R,4S,5S,1'S)-5-(isopropoxycarbonyl)amino-4-t-butyldimethylsiloxy-N-[1'-isopropyl-1'-(N'-

isopropoxycarbonyl)imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide

A mixture containing the compound of 13(a) (0.17 g, 0.31 mmol), isopropyl chloroformate (0.62 mL, 1M in dichloromethane, 2 eq) and 4-dimethylaminopyridine (0.75 g, 2 eq) in dichloromethane (40 mL) was allowed to stir at room temperature overnight under an argon atmosphere. The mixture was then partitioned between dichloromethane and saturated aqueous Na₂CO₃, and the organic extract was dried over Na₂CO₃.

The solvent was removed in vacuo, and the residue was purified by flash chromatography (silica, 4% methanol/dichloromethane) to afford the title compound (0.214 g, 96%). NMR(CDCl₃) δ 7.35-6.78 (12H, m), 6.57 (1H, d), 5.61 (1H, dd), 5.19 (1H, m), 4.86 (1H, m), 4.77 (1H, d), 3.97 (1H, q), 3.63 (1H, t), 2.88 (1H, dd), 2.70-2.48 (4H, m), 2.06 (1H, m), 2.00-1.85 (1H, m), 1.79-1.64 (1H, m), 1.45 (6H, dd), 0.94 (9H, s), 0.85 (6H, d), 0.12 (6H, d).

c) (2R,4S,5S,1'S)-5-(isopropoxycarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide

To a solution of the compound of 13(b) (0.214 g) in methanol, excess aqueous HCl (~5 eq) was added. The resulting solution was allowed to stir at room temperature overnight, and was concentrated under reduced pressure. The residue was diluted with H₂O, and basified with aqueous Na₂CO₃. The mixture was extracted with dichloromethane, and the combined organic extracts were dried over Na₂CO₃. The solvent was removed in vacuo, and the residue was purified by flash chromatography (silica, 4% methanol/dichloromethane) to afford the title compound (0.150 g, 97%). NMR(CDCl₃) δ 7.32-6.96 (13H, m), 5.48 (1H, d), 5.08 (1H, m), 5.00 (1H, s(br)), 4.87 (1H, m), 3.78 (1H, m), 3.62 (1H, m), 3.25 (1H, m), 2.96-2.67 (4H, m), 2.29 (1H, m), 1.95-1.65 (2H, m), 1.25-1.12 (6H, dd), 0.80-0.60 (6H, dd); MS m/e 521 [M+H]⁺, 519 (M-H)⁻.

d) (2R,4S,5S,1'S)-5-(isopropoxycarbonyl)amino-4-hydroxy-N-
(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-
phenylmethyl-hexanamide hydrochloride

The product of 13(c) (100 mg, 0.192 mmol) was dissolved
in methanol (10 mL) and a 1M solution of HCl in ether (0.192
mL) was added. The solution was concentrated by rotary
evaporation without heating, and the residue was triturated
with ether and dried under vacuum to afford the title
compound (104 mg, 98%). NMR(CD₃OD) δ 7.30 (2H, s), 7.21-
6.88 (10H, m), 4.61 (2H, m), 3.65 (1H, m), 3.48 (1H, d), 2.99
(1H, m), 2.87 (1H, m), 2.74-2.56 (2H, m), 2.12 (1H, m), 1.75-
1.50 (2H, m), 1.17-1.00 (6H, dd), 0.90 (3H, d), 0.64 (3H, d).

Example 14

15

Preparation of (2R,4S,5S,1'S)-5-(benzyloxyethoxycarbonyl)
amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-
phenyl-2-phenylmethyl-hexanamide

In (20 mg) benzyloxyethyl-(4-nitro)phenylcarbonate
To a solution of 2-benzyloxyethanol (2.5 g, 16.4 mmol)
and bis(4-nitrophenyl)carbonate (5.0 g, 1 eq) in
dichloromethane (200 mL), N-methylmorpholine (1.81 mL, 1 eq)
was added. The resulting mixture was allowed to stir at room
temperature for 3 d. The reaction mixture was washed
successively with H₂O and saturated aqueous NaCl and dried
over Na₂SO₄. The solvent was removed in vacuo, and the
residue was purified by flash chromatography (silica, 20%
ethyl acetate/hexanes) to afford the title compound (4.38 g,
84%). NMR(CDCl₃) δ 8.26 (2H, m), 7.34 (7H, m), 4.62 (2H, s),
4.49 (2H, t), 3.70 (2H, t).

b) (2R,4S,5S,1'S)-5-(benzyloxyethoxycarbonyl)amino-4-t-
butyldimethylsiloxy-N-[1'-isopropyl-1'-(N'-benzyloxyethoxy-
carbonyl)imidazol-2-yl]methyl-6-phenyl-2-phenylmethyl-
hexanamide

To a solution of the compound of Example 14(a) (134.5
mg, 0.24 mmol) in dichloromethane (40 mL) under an argon

- 50 -

atmosphere, benzyloxyethyl 4-nitrophenyl carbonate (160 mg, 2 eq) and 4-dimethylaminopyridine (60 mg, 2 eq) were added. The resulting mixture was allowed to stir at room temperature overnight, and was diluted with dichloromethane. The organic extract was washed successively with aqueous Na₂CO₃, H₂O, aqueous Na₂CO₃ and H₂O, and dried over Na₂CO₃. The solvent was removed in vacuo, and the residue was purified by flash chromatography (silica, 4% methanol/dichloromethane) to afford the title compound (180 mg, 82%). NMR(CDCl₃) δ 7.45-6.80 (22H, m), 6.62 (1H, d), 5.60 (1H, t), 5.06 (1H, d), 4.60 (2H, s), 4.52 (2H, s), 4.50 (2H, m), 4.31 (1H, m), 4.07 (2H, m), 3.80 (2H, t), 3.68 (1H, q), 3.57 (1H, q), 2.85 (1H, m), 2.77-2.41 (4H, m), 2.09 (1H, m), 1.90 (1H, m), 1.73 (1H, m), 0.95 (9H, s), 0.81 (6H, dd), 0.11 (6H, d).

15

(c) (2R,4S,5S,1'S)-5-(benzyloxyethoxycarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide

Following the procedure of Example 13(c), except using the compound of Example 14(b) (160 mg), the title compound was prepared (100 mg, 81%). NMR(CDCl₃, CD₃OD) δ 7.40-6.79 (17H, m), 4.55 (2H, s), 4.45 (1H, d), 4.20 (2H, m), 3.80-3.45 (5H, m), 2.95-2.66 (4H, m), 2.59 (1H, dd), 2.07 (1H, m), 1.71 (2H, m), 0.80 (3H, d), 0.68 (3H, d).

25

Example 15

Preparation of (2R,4S,5S,1'S)-5-(methoxycarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide

a) (2R,4S,5S,1'S)-5-(methoxycarbonyl)amino-4-t-butylldimethylsiloxy-N-[1'-isopropyl-1'-(N'-methoxycarbonyl)imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide

Following the procedure of Example 13(b), except using (2R,4S,5S,1'S)-5-amino-4-t-butylldimethylsiloxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-

- 51 -

hexanamide, the title compound was prepared (89%).

NMR(CDCl₃) δ 7.40-6.79 (12H, m), 6.52 (1H, d), 5.58 (1H, dd), 4.91 (1H, d), 3.96 (3H, s), 3.95 (1H, d), 3.66 (1H, t), 3.60 (3H, s), 2.85 (1H, m), 2.73-2.40 (4H, m), 2.08 (1H, m), 1.90 (1H, m), 1.69 (1H, m), 0.95 (9H, s), 0.85 (6H, dd), 0.14 (6H, d).

b) (2R,4S,5S,1'S)-5-(methoxycarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide

Following the procedure of Example 13(c), except using the compound of Example 15(a), the title compound was prepared (70%). NMR(CDCl₃, CD₃OD) δ 7.23-6.60 (12H, m), 4.38 (1H, d), 3.65 (1H, t), 3.54 (3H, s), 3.33 (1H, m), 2.95 (1H, m), 2.82-2.40 (4H, m), 1.95 (1H, m), 1.64 (2H, m), 0.69 (6H, dd).

Example 16

20 Preparation of (2R,4S,5S,1'S)-5-(ethoxycarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide

a) (2R,4S,5S,1'S)-5-(ethoxycarbonyl)amino-4-t-butyl-25 dimethylsiloxy-N-[1'-isopropyl-1'-(N'-ethoxycarbonyl)imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide

Following the procedure of Example 13(b), except using (2R,4S,5S,1'S)-5-amino-4-t-butyl-2-dimethylsiloxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide and ethylchloroformate, the title compound was

prepared (90%). NMR(CDCl₃) δ 7.35-6.77 (12H, m), 6.55 (1H, d), 5.60 (1H, dd), 4.86 (1H, d), 4.41 (2H, m), 4.15-3.90 (3H, m), 3.66 (1H, t), 2.87 (1H, m), 2.75-2.45 (4H, m), 2.08 (1H, m), 1.92 (1H, m), 1.70 (1H, m), 1.45 (3H, t), 1.18 (3H, t), 0.98 (9H, s), 0.85 (6H, dd), 0.13 (6H, d).

- 52 -

b) (2R,4S,5S,1'S)-5-(ethoxycarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide

Following the procedure of Example 13(c), except using
 5 the compound of Example 16(a), the title compound was prepared (95%). NMR(CDCl₃, CD₃OD) δ 7.25-6.75 (12H, m), 4.43 (1H, d), 3.95 (2H, q), 3.61 (1H, q), 3.40 (1H, m), 2.85 (1H, m), 2.80-2.40 (4H, m), 2.05 (1H, m), 1.61 (2H, t), 1.11 (3H, t), 0.72 (3H, d), 0.55 (3H, t).

10

Example 17

Preparation of (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-(3-phenyl-2-propenyl)hexanamide

15

a) (3R,5S,1'S)-(1'-t-butoxycarbonylamino-2'-phenyl)ethyl-3-(3-phenylprop-2-enyl)-tetrahydrofuran-2-one

Following the procedure of Example 12(a), except using
 20 cinnamyl bromide (0.485 mL) as the alkylating agent, the title compound was prepared (0.51 g, 75%). NMR(CDCl₃) δ 7.35-7.10 (10H, m), 6.43 (1H, d), 6.09 (1H, m), 4.60 (1H, m), 4.48 (1H, q), 4.00 (1H, t (br)), 2.96-2.55 (4H, m), 2.53-2.21 (2H, m), 2.05 (1H, m), 1.35 (9H, s).

25

b) (2R,4S,5S)-5-(t-butoxycarbonyl)amino-4-t-butyl-dimethyl-siloxy-6-phenyl-2-(3-phenyl-2-propenyl)hexanoic acid

Following the procedure of Example 12(b), except using
 30 the compound of Example 17(a), the title compound was prepared (77%). NMR(CDCl₃) δ 7.40-7.05 (10H, m), 6.48-6.00 (4H, m), 4.78 (1H, d), 3.94 (1H, q), 3.80 (1H, m), 2.89 (1H, m), 2.83-2.26 (4H, m), 1.90 (1H, m), 1.59 (1H, m), 1.28 (9H, s), 0.90 (9H, s), 0.08 (6H, d).

(c) (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-(t-butyldimethylsiloxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-(3-phenyl-2-propenyl)hexanamide

Following the procedure of Example 1(c), except using the compound of 17(b), the title compound was prepared (82%).
NMR(CDCl₃) δ 7.35-7.15 (10H, m), 7.14-6.85 (2H, m), 6.73 (1H, s), 6.20 (1H, d), 6.10-5.88 (1H, m), 4.78 (1H, d), 4.65 (1H, s), 3.97 (1H, q), 3.76 (1H, m), 2.77 (2H, d), 2.50-2.25 (2H, m), 2.12 (1H, m), 1.70 (1H, m), 1.63 (1H, m), 1.36 (9H, s), 1.00-0.92 (9H, s), 0.81 (6H, d), 0.09 (6H, d).

d) (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-(3-phenyl-2-propenyl)hexanamide

Following the procedure of Example 9(d), except using the compound of 17(c), the title compound was prepared (90%).
NMR(CDCl₃, CD₃OD) δ 7.30-7.00 (10H, m), 6.71 (2H, s), 6.26 (1H, d), 6.41 (1H, m), 3.66 (1H, d), 3.50 (1H, d), 2.88-2.45 (4H, m), 2.36 (1H, m), 2.23 (1H, m), 2.06 (1H, m), 1.70 (2H, m), 1.34 (9H, s), 0.88 (3H, d), 0.74 (3H, d).

MS m/e 561 [M+H]⁺.

Example 18

Preparation of (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-(4-nitroimidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide

a) (1S)-N-(1-(imidazol-2-yl)-2-methyl)propylacetamide

To a solution of the compound of Example 1(b) (175 mg) in dichloromethane (10 mL) at 10°C was added diisopropylethylamine (355 mg, 2.75 mmol) followed by acetyl chloride (215 mg, 2.75 mmol). The resulting mixture was stirred overnight, washed with saturated aqueous Na₂CO₃, and

concentrated. The residue was treated with methanol, stirred overnight and concentrated under reduced pressure to afford

the title compound (181 mg, 78%) as a white solid.

NMR(CD₃OD) δ 6.95 (2H, s), 4.72 (1H, d, J=6 Hz), 2.35-2.10

- 54 -

(1H, m), 1.98 (3H, s), 0.98 (3H, d, J=5, 3 Hz), 0.82 (3H, d, J=5 Hz).

b) (1S)-N-(1-(4-nitroimidazol-2-yl)-2-methyl)propylacetamide

5 The compound of Example 18(a) (290 mg, 1.60 mmol) was dissolved in cold concentrated H_2SO_4 (2 mL), and after stirring for 15 min, 90% HNO_3 (0.4 mL) was added dropwise.

The resulting mixture was slowly warmed to 40°C and stirred for 2 h. The mixture was then poured onto ice, and the pH was adjusted to 4 by the addition of solid $NaHCO_3$. The

mixture was extracted with ethyl acetate (6x), and the combined organic extracts were dried over $MgSO_4$ and concentrated under reduced pressure to afford the title compound (153 mg, 42%). NMR(CD_3OD) δ 7.98 (1H, s), 4.70 (1H, d, J=6 Hz), 2.35-2.15 (1H, m), 1.98 (3H, s), 0.95 (3H, d, J=5 Hz), 0.85 (3H, d, J=5 Hz); MS m/e 475.2 (2M+Na)⁺, 249.2 (M+Na)⁺, 227.2 [M+H]⁺, 185.2, 168.0.

c) (1S)-1-(4-nitroimidazol-2-yl)-2-methylpropylamine, dihydrochloride salt

A mixture of the compound of Example 18(b) (153 mg, 0.68 mmol) in 6N HCl (2 mL) was heated at 90°C for 12 h, cooled and concentrated under reduced pressure. The title compound was obtained (138 mg, 80%) and used without further purification. NMR(CD_3OD) δ 8.12 (1H, s), 4.30 (1H, d, J=4 Hz), 2.45-2.30 (1H, m), 1.12 (3H, d, J=4 Hz), 0.90 (3H, d, J=4 Hz).

d) (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-t-butyl-2-butyl-1-dimethylsiloxy-N-[1'-isopropyl-1'-(4-nitroimidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide

Following the procedure of Example 1(c), except using (2R,4S,5S)-5-(t-butoxycarbonyl)amino-4-t-butyl-2-butyl-1-dimethylsiloxy-6-phenyl-2-phenylmethylhexanoic acid and (1S)-1-(4-nitroimidazol-2-yl)-2-methylpropylamine, the title compound was prepared. NMR($CDCl_3$) δ 7.30-6.90 (10 H, m), 6.60 (1H, d, J=4 Hz), 4.70 (1H, d, J=5 Hz), 4.40 (1H, t, J=4 Hz), 3.90 (1H, q, J=4 Hz), 3.75 (1H, dd, J=8, 3 Hz), 2.75-2.30 (6H, m),

- 55 -

1.80-1.50 (2H, m), 1.25 (9H, s), 0.85 (9H, s), 0.70 (6H, m), 0.05 (6H, d, J=4 Hz).

(e) (2R,4S,5S,1'S)-5-(*t*-butoxycarbonyl)amino-4-hydroxy-N-[1'-isopropyl-1'-(4-nitroimidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide

Following the deprotection procedure of Example 1(d), except using the compound of Example 18(d), the title compound was prepared. NMR(CD₃OD) δ 7.90 (1H, s), 7.40-6.90 (10H, m), 4.53 (1H, d, J=6 Hz), 3.70 (1H, m), 3.50 (1H, m), 2.90-2.60 (5H, m), 2.00 (1H, m), 1.90-1.55 (2H, m), 1.49 (9H, s), 0.85 (3H, d, J=4 Hz), 0.70 (d, 3H, J=4 Hz); MS m/e 602.4 (M+Na)⁺, 580.4 [M+H]⁺, 524.4, 480.4.

15

Example 19

Preparation of (2R,4S,5S,1'S)-5-(*t*-butoxycarbonyl)amino-4-

hydroxy-N-(1'-ethyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide

20

a) (1S)-1-carbobenzyloxyamino-1-ethyl-1-(imidazol-2-yl)methane

Following the procedure of Example 1(a), except using Cbz-(L)-α-ethylglycinal in place of valinal, the title compound was prepared. NMR(CDCl₃) δ 7.45-7.10 (5H, m), 6.90 (2H, s), 5.65 (1H, d, J=6 Hz), 5.10-4.95 (2H, m), 4.40 (1H, q, J=5 Hz), 2.00-1.70 (2H, m), 1.00-0.80 (3H, m).

b) (1S)-(1-imidazol-2-yl)propylamine

30 Following the procedure of Example 1(b), except using the compound of Example 19(a), the title compound was prepared. NMR(CDCl₃) δ 6.90 (2H, s), 5.00-4.50 (2H, br s), 4.00 (1H, t, J=5 Hz), 2.00-1.70 (2H, m), 1.00-0.80 (3H, m).

- 56 -

c) (2R,4S,5S,1'S)-5-(*t*-butoxycarbonyl)amino-4-*t*-butyldimethylsiloxy-N-[1'-ethyl-1'-imidazol-2-yl]methyl-6-phenyl-2-phenylmethyl-hexanamide

Following the procedure of Example 1(c), except using (2R,4S,5S)-5-(*t*-butoxycarbonyl)amino-4-*t*-butyldimethylsiloxy-6-phenyl-2-phenylmethylhexanoic acid and the compound of Example 19(c), the title compound was prepared. NMR(CDCl₃) δ 7.35-6.90 (10H, m), 6.78 (2H, s), 6.20 (d, J=5 Hz), 4.80-4.65 (2H, m), 4.05 (1H, q, J=5 Hz), 3.72 (1H, dd, J=10, 3 Hz), 2.90-2.50 (5H, m), 2.10-2.05 (1H, m), 1.90-1.65 (3H, m), 1.40 (9H, s), 0.95 (9H, s), 0.90-0.85 (3H, m), 0.50 (6H, s).

d) (2R,4S,5S,1'S)-5-(*t*-butoxycarbonyl)amino-4-hydroxy-N-[1'-ethyl-1'-imidazol-2-yl]methyl-6-phenyl-2-phenylmethyl-hexanamide

Following the procedure of Example 9(d), except using the compound of Example 19(c), the title compound was prepared. NMR(CD₃OD) δ 7.40-7.00 (10H, m), 6.85 (2H, s), 3.60-3.50 (2H, m), 2.95-2.60 (5H, m), 1.95-1.52 (4H, m), 1.48-1.26 (9H, m), 0.8-0.9 (3H, m). MS m/e 521.2 [M+H]⁺; 503.4, 447.4.

Example 20

25 Preparation of (2R,4S,5S,1'S)-5-(*t*-butoxycarbonyl)amino-4-hydroxy-N-(1'-propyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide

Following the procedure of Example 19(a)-19(d), except substituting Cbz-(L)- α -propylglycinal for Cbz-(L)- α -ethylglycinal, the title compound was prepared. Data for the intermediates of this synthesis were:

a) (1S)-1-carbobenzyloxyamino-1-propyl-1-(imidazol-2-yl)methane. NMR(CDCl₃) δ 7.40-7.10 (10H, m), 6.65 (2H, s), 5.55 (1H, d, J=6 Hz), 5.10-4.90 (2H, m), 4.65 (1H, q, J=5 Hz), 2.05-1.93 (1H, m), 1.90-1.75 (1H, m), 1.45-1.20 (4H, m), 0.95-0.85 (3H, m).

b) (1S)-1-(imidazol-2-yl)butylamine. NMR(CDCl₃) δ 6.90 (2H, s), 5.10-4.40 (2H, s(br)), 4.05 (1H, t, J=5 Hz), 1.90-1.55 (2H, m), 1.45-1.20 (4H, m), 0.95-0.80 (3H, m).

c) (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-t-butylldimethylsiloxyl-N-[1'-propyl-1'-imidazol-2-yl]methyl-6-phenyl-2-phenylmethyl-hexanamide. NMR(CDCl₃) δ 7.35-7.00 (10H, m), 6.78 (2H, s), 6.22 (1H, d, J=5 Hz), 4.85-4.68 (2H, m), 4.00 (1H, q, J=3 Hz), 3.75 (1H, dd, J=10, 3 Hz), 2.80-2.50 (5H, m), 2.12-1.95 (1H, m), 1.90-1.60 (3H, m), 1.40-1.20 (13H, m), 0.90 (9H, s), 0.87-0.80 (3H, m), 0.07 (6H, s).

d) (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-[1'-propyl-1'-imidazol-2-yl]methyl-6-phenyl-2-phenylmethyl-hexanamide. NMR(CD₃OD) δ 7.40-7.00 (10H, m), 6.90 (2H, s), 3.78-3.50 (2H, m), 2.90-2.60 (5H, m), 1.90-1.55 (4H, m), 1.45-1.20 (13H, m); MS m/e 535.4 [M+H]⁺.

Example 21

Preparation of (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-[1'-isopropyl-1'-(4-bromoimidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide

a) (1S)-N-1-(4-bromoimidazol-2-yl)-2-methylpropylacetamide and (1S)-N-1-(4,5-dibromoimidazol-2-yl)-2-methylpropylacetamide

To a solution of (1S)-N-1-imidazol-2-yl-2-methylpropylacetamide (1.58 g, 8.73 mmol) in 95% ethanol (30 mL), 2,4,4,6-tetrabromocyclohexadienone (3.93 g, 9.60 mmol) was added. The resulting mixture was stirred at room temperature for 30 min, and was concentrated in vacuo. The

- 58 -

residue was dissolved in dichloromethane, washed with aqueous NaHCO₃ and dried over Na₂SO₄. The solvent was removed in vacuo, and the residue was purified by flash chromatography to afford the title compound (650 mg, 29%). NMR(CDCl₃) δ 7.70 (1H, d, J=7 Hz), 6.85 (1H, s), 4.67 (1H, t, J=7 Hz), 2.35-2.25 (1H, m), 1.95 (3H, s), 1.05 (3H, d, J=5 Hz), 0.80 (3H, d, J=5 Hz).

Also isolated was (1S)-N-1-(4,5-dibromoimidazol-2-yl)-2-methylpropylacetamide (50 mg, 8%): NMR(CDCl₃) δ 4.68 (1H, t, J=7 Hz), 2.38-2.25 (1H, m), 2.05 (3H, s), 1.05 (3H, d, J=5 Hz), 0.85 (3H, d, J=5 Hz); MS m/e 340.0 [M+H]⁺, 280.8.

b) (1S)-1-(4-bromoimidazol-2-yl)-2-methylpropylamine, dihydrochloride
Following the procedure of Example 18(c), except using (1S)-N-1-(4-bromoimidazol-2-yl)-2-methylpropylacetamide, the title compound was prepared. NMR(CD₃OD) δ 7.60 (1H, s), 4.35 (1H, d, J=7 Hz), 2.50-2.38 (1H, m), 1.10 (3H, d, J=5 Hz), 0.82 (3H, d, J=5 Hz).

20

c) (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-t-butyl-
butyldimethylsiloxyl-N-[1'-isopropyl-1'-(4-bromoimidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide

Following the procedure of Example 1(c), except using (2R,4S,5S)-5-(t-butoxycarbonyl)amino-4-t-butyl-
butyldimethylsiloxyl-N-[1'-isopropyl-1'-(4-bromoimidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide, the title compound was prepared. NMR(CDCl₃) δ 7.40-7.00 (10H, m), 6.70 (1H, s), 6.45 (1H, d, J=5 Hz), 4.80 (1H, d, J=6 Hz), 4.40 (1H, t, J=5 Hz), 4.02 (1H, q, J=4 Hz), 3.78 (1H, dd, J=7, 2 Hz), 2.90-2.30 (9H, m), 1.85-1.60 (2H, m), 1.45 (9H, s), 1.00 (9H, s), 0.85 (6H, t, J=4 Hz), 0.10 (6H, d, J=6 Hz).

d) (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-[1'-isopropyl-1'-(4-bromoimidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide

Following the procedure of Example 9(d), except using (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-t-butyl-
butyldimethylsiloxyl-N-[1'-isopropyl-1'-(4-bromoimidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide, the title compound was prepared.

siloxo-N-[1'-isopropyl-1'-(4-bromoimidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide, the title compound was prepared. NMR(CDCl₃) δ 7.40-7.00 (10H, m), 6.70 (1H, s), 6.55 (1H, m), 4.90 (1H, d, J=5 Hz), 4.50 (1H, t, J=5 Hz), 3.75-3.55 (2H, m), 2.95-2.65 (5H, m), 2.40-2.25 (1H, m), 1.90-1.60 (2H, m), 1.48 (9H, s), 0.80 (6H, t, J=6 Hz).

MS m/e 613.2 [M+H]⁺; 535.2.

Example 22

10

Preparation of (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-[1'-isopropyl-1'-(4,5-dibromoimidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide

Following the procedures of Examples 18(c)-18(d) and 19(d), except substituting (1S)-N-1-(4,5-dibromoimidazol-2-yl)-2-methylpropylacetamide for (1S)-N-(1-4-nitroimidazol-2-yl)-2-methylpropylacetamide, the title compound was prepared.

Analytical data for the intermediates of this synthesis were:

a) (1S)-1-(4,5-dibromoimidazol-2-yl)-2-methylpropylamine, dihydrochloride: NMR(CD₃OD) δ 4.10-3.90 (1H, br s), 2.30-2.10 (2.10 (1H, s(br)), 1.10 (3H, d, J=5 Hz), 0.85 (3H, d, J=5 Hz).

b) (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-t-butyl-dimethylsiloxo-N-[1'-isopropyl-1'-(4,5-dibromoimidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide. NMR(CDCl₃) δ 7.40-6.90 (10H, m), 6.38 (1H, d, J=5 Hz), 4.80-4.50 (3H, m), 4.00 (1H, q, J=5 Hz), 3.72 (1H, dd, J=7, 2 Hz), 2.85-2.50 (5H, m), 2.30 (1H, br s), 2.20-2.05 (1H, m), 1.85-1.65 (2H, m), 1.38 (9H, s), 0.90 (9H, s), 0.80-0.60 (6H, m), 0.10 (6H, d, J=3 Hz).

c) (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-[1'-isopropyl-1'-(4,5-dibromoimidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide. NMR(CDCl₃) δ 7.35-6.85 (10H, m), 6.65 (1H, br s), 4.92 (1H, d, J=4 Hz), 4.50 (1H, m), 3.72-3.50 (2H, m), 2.98-2.63 (5H, m), 2.15-2.02 (1H, m), 1.90-1.70

- 60 -

(2H, m), 1.40 (9H, s); MS m/e 693.03 [M+H]⁺; 637, 619, 593, 575, 291.

Example 23

Preparation of (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-[1'-isopropyl-1'-(4-methylimidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide

10

a) (1S)-1-carbobenzyloxyamino-1-isopropyl-1'-(4-methylimidazol-2-yl)methane.

Cbz-(L)-valinal (1.0 g, 3.9 mmol) and pyruvaldehyde (4.3 mmol, 40% in H₂O) were dissolved in methanol (10 mL) and

15 chilled in an ice bath. Concentrated aqueous ammonia (2 mL) was added and the reaction mixture was stirred at 20°C

overnight. The solvent was removed in vacuo and the residue dissolved in 5% HCl (50 mL) and extracted with ethyl acetate

(3x20 mL). The aqueous layer was basified to pH 10 with

20 solid Na₂CO₃. A tan solid (463 mg) precipitated. The solid was purified by flash chromatography (silica, 2%-3%

methanol/dichloromethane) to yield the title compound as a white solid (180 mg, 16%). mp 163-164°C; NMR(CDCl₃) δ 7.45-

7.35 (5H, m), 6.60 (1H, s), 6.00 (1H, d, J=4 Hz), 5.05 (2H,

25 q, J=4 Hz), 4.40 (1H, t, J=4 Hz), 2.45-2.30 (1H, m), 2.20

(3H, s), 0.95 (3H, d, J=4 Hz), 0.80 (3H, d, J=4 Hz); MS m/e 575.4 (2M+H)⁺, 288.0 [M+H]⁺.

b) (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-t-

30 butyldimethylsiloxy-N-[1'-isopropyl-1'-(4-methylimidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide

Following the procedure of Example 1(b)-1(c), except using (2R,4S,5S)-5-(t-butoxycarbonyl)amino-4-t-

butyldimethylsiloxy-6-phenyl-2-phenylmethylhexanoic acid and

35 the compound of Example 23(a), the title compound was

prepared. NMR(CDCl₃) δ 7.37-6.90 (10H, m), 6.45 (1H, s),

6.38 (1H, d, J=3 Hz), 4.75 (1H, d, J=5 Hz), 4.40 (1H, t, J=5

Hz), 3.95 (1H, q, J=4 Hz), 3.72-3.68 (1H, m), 2.90-2.70 (4H,

m), 2.60-2.48 (1H, m), 2.45-2.30 (1H, m), 2.17 (3H, s), 1.90-1.80 (1H, m), 1.75-1.62 (1H, m), 1.40 (9H, s), 0.95 (9H, s), 0.75 (6H, t, J=3 Hz), 0.10 (6H, d, J=2 Hz).

5 c) (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-[1'-isopropyl-1'-(4-methylimidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide

Following the procedure of Example 9(d), except using the compound of Example 23(b), the title compound was

10 prepared. NMR(CDCl₃) δ 7.38-7.00 (10H, m), 6.52 (1H, s), 4.92 (1H, d, J=5 Hz), 4.42 (1H, t, J=4 Hz), 3.72-3.55 (2H, m), 2.95-2.65 (5H, m), 2.35-2.20 (1H, m), 2.18 (3H, s), 1.75 (2H, br s), 1.42 (9H, s), 0.75 (6H, d, J=3 Hz); MS m/e 549.2 [M+H]⁺.

15

Example 24

Preparation of (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-[1'-isopropyl-1'-(4-trifluoromethylimidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide

20

a) (1S)-1-carbobenzyloxyamino-1-isopropyl-1-(4-trifluoromethylimidazol-2-yl)methane.

Sodium acetate trihydrate (5.35 g, 2.2 eq) was dissolved in water (16 mL) and 1,1 dibromotrifluoroacetone (5.31 g, 1.1 eq) was added. The solution was stirred for 30 min at 90°C. The solution was cooled to 0°C and poured into a 0°C solution of Cbz-Valinal (4.22 g, 1.0 eq) in anhydrous methanol (80 mL). Concentrated ammonium hydroxide (22 mL) was added and the mixture stirred overnight at room temperature. The solvents were evaporated to give a white precipitate which was covered with 150 mL of water. The suspension was filtered and the solid washed twice with water. The white solid was dissolved in ethyl acetate, dried over sodium sulfate, filtered, and evaporated to a white solid (5.24 g, 86%).

35

¹HNMR(CD₃OD) δ 7.45 (1H, s), 7.40-7.20 (5H, m), 5.05 (2H, q, J=4 Hz), 4.50 (1H, d, J=4 Hz), 2.38-2.10 (1H, m), 1.00 (3H, d, J=4 Hz), 0.80 (3H, d, J=4 Hz), ¹³CNMR(CD₃OD, ¹H-decoupled)

- 62 -

δ 18.9, 19.4, 67, 117(q, J=3 Hz), 123.2(q, J=266 Hz), 128.7, 129.3, 133(q, J=39 Hz), 138.0, 151.7; MS m/e 342.0 [M+H]⁺.

b) (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-t-butyl-dimethylsiloxo-N-[1'-isopropyl-1'-(4-trifluoromethylimidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide

Following the procedure of Example 1(b)-1(c), except using the compound of Example 24(a) and (2R,4S,5S)-5-(t-butoxycarbonyl)amino-4-t-butyl-dimethylsiloxo-6-phenyl-2-phenylmethylhexanoic acid, the title compound was prepared. NMR(CDCl₃) δ 7.35-6.95 (11 H, m), 6.50 (1H, d, J=4 Hz), 4.75 (1H, d, J=6 Hz), 4.25 (1H, t, J=4 Hz), 3.95 (1H, q, J=4 Hz), 3.80-3.68 (1H, m), 2.90-2.40 (5H, m), 1.80-1.60 (2H, m), 1.35 (9H, s), 0.90 (9H, s), 0.80 (3H, d, J=3 Hz), 0.70 (3H, d, J=3 Hz), 0.05 (6H, d, J=2 Hz).

c) (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-[1'-isopropyl-1'-(4-trifluoromethylimidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide

Following the procedure of Example 9(d), except using the compound of Example 24(b), the title compound was prepared. NMR(CDCl₃) δ 7.35 (1H, s), 7.25-6.90 (10H, m), 4.53 (1H, d, J=5 Hz), 3.68 (1H, t, J=4 Hz), 3.52 (1H, d, J=6 Hz), 2.90-2.55 (5H, m), 2.10-1.95 (1H, m), 1.85-1.70 (1H, m), 1.65-1.50 (1H, m), 1.40-1.25 (9H, m), 0.90 (3H, d, J=4 Hz), 0.65 (3H, d, J=4 Hz); MS m/e 603.2 [M+H]⁺, 529.2, 503.2.

Example 25

Preparation of (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-methyl-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide

a) (1S)-1-carbobenzyloxyamino-1-isopropyl-1-(imidazol-2-yl)methane

Following the procedure of Example 1(a), except substituting N-methyl-Cbz-(L)-valinal for Cbz-(L)-valinal, the title compound was prepared. NMR(CDCl₃) δ 7.45-7.30 (5H,

m), 6.90 (2H, s), 5.12 (2H, s), 4.60 (1H, d, J=6 Hz), 2.95 (3H, s), 2.70-2.53 (1H, m), 1.02 (3H, d, J=3 Hz), 0.85 (3H, d, J=3 Hz).

5. b) (1S)-1-methylamino-1-isopropyl-1-(imidazol-2-yl)methane

Following the procedure of Example 1(b), except using the compound of Example 25(a), the title compound was prepared. NMR(CDCl₃) δ 6.95 (2H, s), 3.52 (1H, d, J=3 Hz), 2.30 (3H, s), 2.10-1.90 (1H, m), 0.98 (3H, d, J=3 Hz), 0.82 (3H, d, J=3 Hz).

15 c) (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-t-butylbutyldimethylsiloxyl-N-methyl-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide

Following the procedure of Example 1(c), except using the compound of Example 25(b), the title compound was prepared. NMR(CDCl₃) δ 7.40-6.72 (12H, m), 4.82 (1H, d, J=5 Hz), 3.95 (1H, q, J=4 Hz), 3.82-3.75 (1H, m), 2.95-2.70 (5H, m), 2.51 (2H, s), 2.50-2.38 (1H, m), 2.08 (1H, s), 1.87-1.68 (2H, m), 1.38 (9H, s), 0.95 (9H, s), 0.88 (3H, d, J=3 Hz), 0.75 (3H, d, J=3 Hz), 0.05 (6H, d, J=7 Hz).

25 d) (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-methyl-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide

Following the procedure of Example 9(d), except using the compound of Example 28(c), the title compound was prepared. NMR(CDCl₃) δ 7.35-6.82 (12H, m), 4.90-4.72 (1H, m), 3.70-3.00 (2H, m), 2.92-2.50 (8H, m), 1.90-1.60 (2H, m), 1.40-1.30 (9H, m), 0.95-0.70 (6H, m). MS m/e 549.2 [M+H]⁺.

Example 26

35 Preparation of (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-[1'-isopropyl-1'-(4-carbomethoxyimidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide

a) (1S)-1-carbobenzyloxyamino-1-isopropyl-1-(4-trimethoxymethylimidazol-2-yl)methane

Sodium methoxide (8 mL, 25% in methanol, 37.5 mmol) was added to a solution of the compound of Example 27(a) (640 mg, 1.88 mmol) in anhydrous methanol (10 mL). The resulting mixture was heated at 55°C overnight, cooled, and concentrated under reduced pressure. The residue was partitioned between ethyl acetate and H₂O, and the organic extract was dried over Na₂CO₃. The solvent was removed in vacuo, and the residue was purified by flash chromatography (silica, 2% methanol/dichloromethane) to afford the title compound (545 mg, 77%). NMR(CDCl₃) δ 7.40-7.20 (5H, m), 6.98 (1H, br s), 5.90 (1H, br s), 5.08 (2H, s), 4.50 (1H, br s), 3.15 (9H, s), 2.00 (1H, m (br)), 1.00-0.80 (6H, m); MS m/e 378.2 [M+H]⁺, 346, 332, 271, 195.

b) (1S)-1-carbobenzyloxyamino-1-isopropyl-1-(4-carbomethoxyimidazol-2-yl)methane

A solution of the compound of Example 26(a) (540 mg) in 1:1 methanol/aqueous HCl (10 mL) was stirred at room temperature for 2 h, and concentrated under reduced pressure. The residue was partitioned between aqueous Na₂CO₃ and dichloromethane, and the organic extract was dried over Na₂CO₃ and concentrated in vacuo to afford the title compound (470 mg, 75%). NMR(CDCl₃) δ 7.55 (1H, br s), 7.35 (5H, s), 5.90-5.65 (1H, m), 5.10 (2H, t, J=4 Hz), 4.60-4.42 (1H, m), 3.88 (3H, s), 2.40 (1H, br s), 1.00-0.80 (6H, m); MS m/e 332.2 [M+H]⁺.

c) (1S)-1-amino-1-isopropyl-1-(4-carbomethoxyimidazol-2-yl)methane

Following the procedure of Example 1(b), except using the compound of Example 26(b), the title compound was prepared. NMR(CDCl₃) δ 7.62 (1H, s), 3.97 (1H, d, J=4 Hz), 3.82 (3H, s), 2.27-2.05 (1H, m), 0.95-0.75 (6H, m).

d) (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-t-butyl-dimethylsiloxy-N-[1'-isopropyl-1'-(4-carbomethoxyimidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide

Following the procedure of Example 1(c), except using the compound of Example 26(c), the title compound was prepared. NMR(CDC1₃) δ 7.45-6.90 (12H, m), 6.48 (1H, d, J=4 Hz), 4.72 (1H, d, J=6 Hz), 4.35 (1H, s br), 4.02-3.87 (1H, m), 3.85 (3H, s), 3.75-3.60 (1H, m), 2.90-2.40 (5H, m), 1.90-1.60 (2H, m), 1.42 (9H, s), 0.90 (9H, s), 0.72 (6H, d, J=4 Hz), 0.10 (6H, d, J=3 Hz).

e) (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-[1'-isopropyl-1'-(4-carbomethoxyimidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide

Following the procedure of Example 9(d), except using the compound of Example 26(d), the title compound was prepared. NMR(CDC1₃) δ 7.40-6.80 (12H, m), 4.90 (1H, d, J=5 Hz), 4.50 (1H, br s), 3.90 (3H, s), 3.80-3.60 (2H, m), 2.95-2.68 (5H, m), 2.45-2.30 (1H, m), 1.80-1.60 (2H, m), 1.40 (9H, s), 0.72 (6H, d, J=4 Hz); MS m/e 593.2 [M+H]⁺, 537.2, 519.2, 493.2, 475.2.

Example 27

25) Preparation of (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-[1'-isopropyl-1'-(4-methylcarbonylimidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide

a) (1S)-1-carbobenzyloxyamino-1-isopropyl-1-(4-hydroxymethylimidazol-2-yl)methane. The compound of Example 26(b) (0.314 g, 1.0 eq) was stirred in anhydrous toluene at -78°C under an argon atmosphere. Diisobutylaluminum hydride (3.8 mL, 1.0M in hexanes, 4.0 eq) was added and the solution stirred at -78°C for 1 h. The reaction was quenched with methanol (0.2 mL, 1.0 eq). The solution was then diluted with Rochelles salt solution (sat.) and stirred for 1 h. The solution was extracted with dichloromethane twice and the combined organic

- 66 -

extracts were washed successively with saturated aqueous Rochelles salt and brine. The organic layer was dried over magnesium sulfate, filtered, and evaporated to give the title compound as a white solid (0.27 g, 94%). NMR(CDCl₃) δ 7.25 (5H, s), 6.69 (1H, s), 6.14 (1H, d), 5.01 (2H, dd), 4.52 (2H, s), 4.37 (1H, t), 2.19 (1H, m), 0.92 (3H, d), 0.73 (3H, d); MS m/e 304.0 [M+H]⁺.

b) (1S)-1-carbobenzyloxyamino-1-isopropyl-1-(4-formylimidazol-2-yl)methane.

The compound of Example 27(a) (0.11 g, 1.0 eq) was stirred in anhydrous dichloromethane at room temperature under an inert argon atmosphere. Manganese dioxide (0.126 g, 4.0 eq) was added and the mixture was stirred at room temperature overnight. After 16 h and additional 2.0 eq of manganese dioxide was added. The reaction was complete by TLC after 2 h. The mixture was filtered through a pad of Celite® and the filter cake was washed with dichloromethane. The organic solvent was removed in vacuo to give the title compound as a white solid (0.075 g, 69%). NMR(CDCl₃) δ 9.57 (1H, s), 7.54 (1H, s), 7.12 (5H, s), 6.43 (1H, d), 4.96 (2H, dd), 4.43 (1H, t), 2.08 (1H, m), 0.91 (3H, d), 0.62 (3H, t); MS m/e 302.0 [M+H]⁺.

c) (1S,1'RS)-1-carbobenzyloxyamino-1-isopropyl-1-(4-(1'-hydroxyethyl)imidazol-2-yl)methane.

The compound of Example 27(b) (0.1 g, 1.0 eq) was stirred in a 3:1 ether/THF mixture at 0°C under an argon atmosphere. Methyl magnesium bromide (0.47 mL, 3.0M in THF, 4.0 eq) was added and allowed to stir at 0°C for 1.5 h. The solution was diluted with 5% aqueous HCl and made basic with solid sodium carbonate. The solution was extracted with ethyl acetate three times and the combined organic extracts were dried over sodium carbonate, filtered, and evaporated to a white solid (0.1 g, 95%). NMR(CDCl₃) δ 7.19 (5H, s), 6.59 (1H, s), 6.42 (1H, d), 4.92 (2H, dd), 4.73 (1H, m), 2.09 (1H, m), 1.37 (3H, d), 0.82 (3H, d), 0.66 (3H, d).

- 67 -

d) (1S, 1'RS)-1-amino-1-isopropyl-1-(4-(1'-hydroxyethyl)imidazol-2-yl)methane.

The compound of Example 27(c) (0.1 g, 1.0 eq) was stirred in anhydrous methanol with 10% Pd on activated carbon 5 (0.020 g). Hydrogen gas was bubbled through the solution via balloon for 1 h and the reaction was maintained under a hydrogen atmosphere for 3 h. The mixture was filtered through a pad of Celite® and the filter cake washed with methanol. The methanol was evaporated to give the title 10 compound as a white solid (0.05 g, 87%). NMR(CDCl₃) δ 6.63

(1H, s), 4.72 (1H, dd), 3.61 (1H, d), 1.92 (1H, m), 1.49 (3H, dd), 0.84 (3H, d), 0.67 (3H, d).

e) (2R, 4S, 5S, 1'S, 1'RS)-5-(t-butoxycarbonyl)amino-4-t-15 butyldimethylsiloxy-N-[1'-isopropyl-1'-(4-(1''-hydroxyethyl)-imidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide

To a solution of (2R, 4S, 5S)-5-(t-butoxycarbonyl)amino-4-t-20 butyldimethylsiloxy-6-phenyl-2-phenylmethylhexanoic acid (0.131 g, 1.0 eq) in anhydrous dimethylformamide, the compound of Example 27(d) (50 mg, 1.1 eq), BOP reagent (0.11 g, 1.0 eq), and triethylamine (0.04 mL, 1.0 eq) were added. The solution was stirred at room temperature for 16 h. The solution was diluted with water and extracted three 25 times with dichloromethane. The combined organic extracts were washed with water, then brine. The solution was dried over magnesium sulfate, filtered, and evaporated to give a white foam. The foam was chromatographed (silica, 4% 30 methanol/dichloromethane) to afford the title compound as a white foam (0.11 g, 65%). NMR(CDCl₃) δ 7.31-6.54 (12H, m), 4.72 (1H, d), 4.48 (2H, d), 3.82 (1H, q), 3.61 (1H, m), 2.81-2.31 (6H, m), 1.65 (3H, m), 1.48 (3H, d), 1.22 (9H, s), 0.89 (9H, s), 0.70 (3H, d), 0.61 (3H, d), 0.06 (6H, s); MS m/e 693.4 [M+H]⁺.

- 68 -

f) (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-t-butyl-
dimethylsiloxy-N-[1'-isopropyl-1'-(4-methylcarbonyl-
imidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide

The compound of Example 27(e) (45 mg, 1.0 eq) was
5 stirred in dry dichloromethane under an inert argon
atmosphere. Manganese dioxide (23 mg, 4.0 eq) was added and
the mixture was stirred at room temperature for 16 h. An
additional 2.0 eq of manganese dioxide was added and the
reaction was complete by TLC after 2.5 h. The mixture was
10 filtered through a pad of Celite® and the filter cake was
washed with dichloromethane. The organic solvent was
evaporated to give the title compound as a white solid
(0.038 g, 85%). NMR(CDCl₃) δ 7.49-6.76 (11H, m), 6.30 (1H,
br d), 4.71 (2H, m), 3.86 (1H, q), 3.61 (1H, dd), 2.77-2.41
15 (5H, m), 2.31 (3H, s), 1.58 (2H, m), 1.20 (9H, s), 0.83 (9H,
s), 0.69 (6H, dd), 0.04 (6H, d); MS m/e 691.4 [M+H]⁺.

g) (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-[1'-
isopropyl-1'-(4-methylcarbonylimidazol-2-yl)]methyl-6-phenyl-
20 2-phenylmethyl-hexanamide

The compound of Example 27(f) (38 mg, 1.0 eq) was
stirred in anhydrous THF under an argon atmosphere at room
temperature. Tetrabutyl ammonium fluoride (0.33 mL, 1.0M in
THF, 6.0 eq) was added and the solution stirred for 16 h.
25 The solution was diluted with water and extracted three times
with dichloromethane. The combined organic extracts were
washed with water and evaporated to a white solid. The solid
was covered with diethyl ether, decanted twice, and dried to
give the title compound as a white solid (25 mg, 79%).
30 NMR(CDCl₃) δ 7.14 (5H, m), 6.86 (5H, m), 5.14 (1H, d), 4.42
(1H, d), 3.58 (1H, q), 3.45 (1H, d), 2.80-2.50 (5H, m), 1.91
(1H, m), 1.63 (2H, m), 1.26 (9H, s) (rotamer observed), 0.70
(3H, d), 0.57 (3H, d); MS m/e 577.2 [M+H]⁺.

Example 28Preparation of (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-[1'-isopropyl-1'-(4-isopropylcarbonylimidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide

a) (1S, 1'RS)-1-carbobenzyloxyamino-1-isopropyl-1-(4-(1'-hydroxy-2'-methyl)propylimidazol-2-yl)methane.

Following the procedure of Example 27(c), except using isopropyl magnesium bromide (1.024 mL, 2.0M solution, 4.0 eq) in place of methyl magnesium bromide, to yield a crude product. The crude product was chromatographed (silica, 4% methanol/dichloromethane) to yield the title compound as a white solid (0.155 g, 88%). NMR(CDCl₃) δ 7.19 (5H, m), 6.58 (1H, s), 4.91 (2H, m), 4.38 (1H, q), 4.20 (1H, dd), 2.11 (1H, m), 1.83 (1H, m), 0.72 (12H, m); MS m/e 346.2 [M+H]⁺; 328.2, 279.0, 254.0, 205.0, 177.0, 149.0, 118.0.

b) (1S, 1'RS)-1-amino-1-isopropyl-1-(4-(1'-hydroxy-2'-methyl)propylimidazol-2-yl)methane. Following the procedure of Example 27(d), using the compound of Example 31(a), the title compound was prepared as a white foam (96 mg, 100%). NMR(CDCl₃) δ 6.65 (1H, s), 4.21 (1H, dd), 3.90 (1H, s), 2.22 (1H, m), 1.94 (1H, m), 0.93 (6H, m), 0.64 (6H, m); MS m/e 302.0 [M+H]⁺.

c) (2R,4S,5S,1'S,1''RS)-5-(t-butoxycarbonyl)amino-4-t-butyldimethylsiloxy-N-[1'-isopropyl-1'-(4-(1''-hydroxy-2''-methyl)propylimidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide

Following the procedure of Example 27(e), except using the compound of Example 31(b) (96 mg, 1.1 eq), substituting dimethyl formamide as the solvent instead of dichloromethane, and purifying the product by chromatography, the title compound was prepared (168 g, 57%). NMR(CDCl₃) δ 7.22-6.81 (11H, m), 6.62 (1H, dd), 4.71 (1H, dd), 4.53 (1H, t), 4.19 (1H, d), 3.82 (1H, q), 3.58 (1H, dd), 2.71-2.30 (5H, m), 2.03 (1H, m), 1.70 (1H, m), 1.57 (1H, m), 1.14 (9H, s), 0.91 (3H,

- 70 -

d), 0.88 (9H, s), 0.78 (3H, d), 0.67 (3H, d), 0.59 (3H, d), 0.03 (6H, d); MS m/e 721.4 [M+H]⁺.

d) (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-t-butyl-dimethylsiloxy-N-[1'-isopropyl-1'-(4-isopropylcarbonylimidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide

Following the procedure of Example 27(f), except using the compound of 31(c) (168 mg, 1.0 eq) and chromatographing the crude product (silica, 3% methanol/dichloromethane) the title compound was prepared as a white solid (132 mg, 79%). NMR(CDCl₃) δ 7.20-6.76 (11H, m), 5.05 (1H, br, m), 3.88 (1H, q), 3.61 (1H, m), 3.19 (1H, m), 2.80-2.46 (5H, m), 2.22 (1H, m), 2.07 (1H, m), 1.63 (1H, m), 1.15 (16H, m), 0.89 (9H, s), 0.74 (6H, m), 0.08 (6H, d); MS m/e 719.4 [M+H]⁺.

e) (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-[1'-isopropyl-1'-(4-isopropylcarbonylimidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide

Following the procedure of Example 27(g), except using the compound of Example 31(d) (132 mg), the title compound was prepared as a white foam (90 mg, 81%). NMR(CDCl₃) δ 7.48 (1H, s), 7.11 (5H, m), 6.82 (5H, m), 5.29 (1H, d), 4.46 (1H, m), 3.54 (1H, q), 3.48 (1H, m), 3.14 (1H, m), 2.74-2.44 (5H, m), 1.90 (1H, m), 1.61 (2H, m), 1.28 (9H, s) (rotamers observed), 1.13 (6H, m), 0.69 (3H, d), 0.48 (3H, d); MS m/e 605.2 [M+H]⁺.

Example 29

30 Preparation of (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-[1'-isopropyl-1'-(4-phenylcarbonylimidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide

a) (1S,1'RS)-1-carbobenzyloxyamino-1-isopropyl-1-(4-(1'-hydroxy)benzylimidazol-2-yl)methane

Following the procedure of Example 27(c), except substituting phenylmagnesium bromide (0.45 mL, 3.0M solution, 4.0 eq) for methyl magnesium bromide, and chromatographing

the crude product (silica, 3% methanol/dichloromethane) the title compound was prepared as a white solid (175 mg, 96%).
 NMR(CDCl₃) δ 7.26 (1H, d), 7.11 (10H, m), 6.39 (1H, dd), 6.08 (1H, d), 5.63 (1H, d), 4.82 (2H, m), 4.29 (1H, m), 2.01 (1H, m), 0.76 (3H, m), 0.59 (3H, d).

b) (1S,1'RS)-1-amino-1-isopropyl-1-(4-(1'-hydroxy)benzylimidazol-2-yl)methane

Following the procedure of Example 27(d), except using the compound of Example 29(a) (98 mg) the title compound was prepared as a tacky white foam (65 mg, 98%).

c) (2R,4S,5S,1'S,1''RS)-5-(t-butoxycarbonyl)amino-4-t-butyl-
 dimethylsiloxy-N-[1'-isopropyl-1'-(4-(1'-hydroxy)benzylimidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide

Following the procedure of Example 27(e), except using the compound of Example 29(b) (0.065 g, 1.1 eq), and chromatographing the crude product (2% methanol/dichloromethane) the title compound was prepared as a white solid (109 mg, 55%).
 NMR(CDCl₃) δ 7.48-6.79 (16H, m), 4.77 (1H, m), 3.88 (1H, m), 3.61 (1H, m), 2.65 (4H, m), 2.39 (1H, m), 2.15 (1H, m), 1.94 (1H, m), 1.75 (1H, m), 1.56 (1H, m), 1.21 (9H, s) (rotamers observed), 0.86 (9H, s), 0.68 (6H, dd), 0.07 (6H, s); MS m/e 755.4 [M+H]⁺.

d) (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-t-butyl-dimethylsiloxy-N-[1'-isopropyl-1'-(4-phenylcarbonylimidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide

Following the procedure Example 27(f), except using the compound of Example 29(c) (109 mg, 1.0 eq), the title compound was prepared as a white solid (80 mg, 74%).
 NMR(CDCl₃) δ 7.49-6.84 (17H, m), 3.88 (1H, q), 3.63 (1H, t), 2.87-2.49 (6H, m), 2.11 (2H, m), 1.64 (1H, m), 1.11 (9H, s), 0.82 (9H, s), 0.71 (6H, dd), 0.06 (6H, d); MS m/e 753.4 [M+H]⁺.

e) (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-[1'-isopropyl-1'-(4-phenylcarbonylimidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide

Following the procedure of Example 27(g), except using the compound of Example 29(d) (80 mg, 1.0 eq), the title compound was prepared as a white solid (45 mg, 74%).

NMR(CDCl₃) δ 7.84-6.77 (16H, m), 4.48 (1H, d), 3.59 (1H, m), 3.42 (1H, m), 2.80-2.54 (5H, m), 1.99 (1H, m), 1.63 (2H, m), 1.26 (9H, s) (rotamers observed), 0.73 (3H, d), 0.59 (3H, d);

MS m/e 639.2 [M+H]⁺.

Example 30

Preparation of (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-[1'-isopropyl-1'-(4-formylimidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide

a) (1S,1'RS)-1-amino-1-isopropyl-1'-(4-(hydroxy)methylimidazol-2-yl)methane.

Following the procedure of Example 27(d), except using the compound of Example 27(a) (90 mg), the titled compound was prepared (50 mg, 100%). NMR(CDCl₃) δ 6.85 (1H, s), 4.62 (2H, s), 3.85 (1H, d, J=4 Hz), 2.20-2.05 (1H, m), 0.88 (6H, d, J=5 Hz).

b) (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-t-butyldimethylsiloxy-N-[1'-isopropyl-1'-(4-(hydroxy)methylimidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide

Following the procedure of Example 27(e), except using the compound of Example 30(a) (50 mg), and chromatographing the crude product (silica, 2% methanol/dichloromethane) the title compound was prepared (130 mg, 65%). NMR(CDCl₃) δ

7.30-6.95 (11H, m), 4.82 (1H, d), 4.50-4.60 (1H, m), 4.40 (1H, d), 3.90-4.00 (1H, m), 3.60-3.68 (1H, m), 2.45-2.80 (5H, m), 2.20-2.30 (1H, m), 1.75-1.85 (1H, m), 1.60-1.70 (1H, m), 1.30 (9H, s), 0.95 (9H, s), 0.75 (3H, d), 0.62 (3H, d), 0.05 (6H, d).

- 73 -

c) (2R,4S,5S,1'S)-5-(*t*-butoxycarbonyl)amino-4-*t*-butyldimethylsiloxy-N-[1'-isopropyl-1'-(4-formylimidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide

Following the procedure of Example 27(f), except using the compound of Example 30(b) (50 mg), the title compound was prepared (20 mg, 40%). NMR(CDCl₃) δ 9.80(0.5H, s), 9.64

(0.5H, s), 7.50-6.90 (11H, m), 6.52-6.42 (1H, m), 4.88-4.70 (2H, m), 4.42-4.32 (1H, m), 4.02-3.93 (1H, m), 3.78-3.71 (1H, m), 2.90-2.40 (5H, m), 2.30-2.19 (1H, m), 1.87-1.62 (2H, m), 1.45 (9H, s), 0.95 (9H, s), 0.87-0.72 (6H, m), 0.05 (6H, m) (rotamers).

d) (2R,4S,5S,1'S)-5-(*t*-butoxycarbonyl)amino-4-hydroxy-N-[1'-isopropyl-1'-(4-formylimidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide

Following the procedure of Example 27(g), except using the compound of Example 30(c) (20 mg), the title compound was prepared (12 mg, 71%). NMR(CD₃OD) δ 9.60 (1H, s), 7.65 (1H, s), 7.20-6.90 (10H, m), 4.52 (1H, d), 3.60 (1H, m), 3.45 (1H, m), 2.80-2.45 (5H, m), 2.00-1.88 (1H, m), 1.75-1.65 (1H, m), 1.62-1.45 (1H, m), 1.27 (9H, s), 0.82 (3H, d), 0.62 (3H, d); MS m/e 563.4, 242.2, 204.8.

Example 31

25

Preparation of (2R,4S,5S,1'S)-5-(*t*-butoxycarbonyl)amino-4-hydroxy-N-[1'-isopropyl-1'-(4-(hydroxymethyl)-imidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide

Following the procedure of Example 27(g), except using the compound of Example 30(b) (40 mg), the title compound was prepared (20 mg). NMR(CD₃OD) δ 7.27-6.92 (10H, s), 6.72 (1H, s), 4.52 (1H, d), 3.64-3.60 (1H, m), 3.48 (1H, d), 2.82-2.50 (5H, m), 2.03-1.92 (1H, m), 1.78-1.67 (1H, m), 1.63-1.49 (1H, m), 1.28 (9H, s), 0.80 (3H, d), 0.65 (3H, d); MS m/e 565.4.

Example 32

Preparation of (2R,4S,5S,1'S)-5-((tetrahydrothiopyran-4-yl)oxycarbonylamino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide

Following the procedures of Example 14 (a)-14 (c), except using 4-hydroxytetrahydrothiopyran in place of 2-benzyloxyethanol, the title compound was prepared.

10 Analytical data for the intermediates of this synthesis were:

a) (tetrahydrothiopyran-4-yl)-(4-nitro)phenylcarbonate.

NMR(CDCl₃) δ 8.26 (1H, s), 8.22 (1H, s), 7.38 (1H, s), 7.33 (1H, s), 4.79 (1H, m), 2.90-2.75 (2H, m), 2.70-2.52 (2H, m),
15 2.31-2.16 (2H, m), 2.10-1.90 (2H, m).

b) (2R,4S,5S,1'S)-5-((tetrahydrothiopyran-4-yl)oxycarbonyl)-amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide. NMR(CD₃OD) δ 7.12-6.65 (10H, m), 6.64 (2H, s), 5.60 (1H, d), 4.36 (2H, m), 3.58 (1H, q), 3.49 (1H, d), 2.68-2.48 (6H, m), 2.44-2.30 (3H, m), 1.93-1.74 (3H, m), 1.70-1.40 (4H, m), 0.61 (3H, d), 0.50 (3H, d).

Example 33

25

Preparation of (2R,4S,5S,1'S)-5-((tetrahydro-4H-pyran-4-yl)oxycarbonylamino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide

30 Following the procedures of Example 14 (a)-14 (c), except using 4-hydroxytetrahydro-4H-pyran in place of 2-benzyloxyethanol, the title compound was prepared.

Analytical data for the intermediates of this synthesis were:

35 a) (tetrahydro-4H-pyran-4-yl)-(4-nitro)phenylcarbonate.
NMR(CDCl₃) δ 8.32 (1H, s), 8.28 (1H, s), 7.41 (1H, s), 7.38 (1H, s), 5.00 (1H, m), 4.05-2.90 (2H, m), 3.68-3.49 (2H, m), 2.17-2.00 (2H, m), 1.95-1.75 (2H, m).

b) (2R,4S,5S,1'S)-5-((tetrahydro-4H-pyran-4-yl)oxycarbonyl)-amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide. NMR(CD₃OD) δ 7.16-6.89 (10H, m), 6.79 (2H, s), 4.54 (2H, m), 3.82-3.70 (2H, m), 3.69-3.62 (1H, m), 3.50-3.46 (1H, m), 3.45-3.35 (2H, m), 2.79-2.65 (4H, m), 2.64-2.45 (3H, m), 2.00 (1H, m), 1.82-1.62 (3H, m), 1.55-1.45 (2H, m), 1.37 (1H, m), 0.79 (3H, d), 0.63 (3H, d).

10. The compound of Example 1(d) was dissolved in neat TFA.

Example 34

Preparation of (2R,4S,5S,1'S)-5-(4-picolinyloxy)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide

The compound of Example 1(d) was dissolved in neat TFA. After 10 min the solution was concentrated to provide the amine salt, (2R,4S,5S,1'S)-5-amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide trifluoroacetate. This amine salt (25 mg, 1 eq) was dissolved in DME, and 4-picolinium-(p-nitro)phenyl carbonate (23 mg, 1 eq) and triethylamine (0.04 mL, 5 eq) were added. The mixture was stirred under Ar for 17 h. Water was added and the mixture was extracted with dichloromethane. The organic extracts were concentrated and the residue was triturated with ether to yield the title compound (20 mg, 61%). NMR(CD₃OD) δ 8.52 (2H, d), 7.10 (14H, m), 6.87 (2H, s), 5.07 (2H, dd), 4.61 (1H, d), 3.80 (1H, m), 3.59 (1H, m), 2.77 (5H, m), 2.05 (1H, m), 1.83 (1H, m), 1.60 (1H, m), 0.84 (3H, d), 0.59 (3H, d).

MS m/e 570.5 [M+H]⁺.

Example 35

Preparation of (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-(4,4,4-trifluorobut-1-yl)hexanamide

a) (3R,5S,1'S)-(1'-t-butoxycarbonylamino-2'-phenyl)ethyl-3-(4,4,4-trifluorobut-1-yl)-tetrahydrofuran-2-one

To a solution of lithium diisopropyl amide (1.8 mL of a 1.5M solution, 2.2 eq) in tetrahydrofuran (10 mL) was added (5S, 1'S)-(1'-t-butoxycarbonylamino-2'-phenyl)ethyl-tetrahydrofuran-2-one (0.50 g; 1.0 eq) in anhydrous THF (2 mL) at -78°C. After stirring for 15 min at (-78°C), hexamethylphosphoramide (0.57 mL, 2.0 eq) was added to the solution. The solution was stirred for several min and 1,1,1-trifluoro-4-iodobutane (0.78 g, 2.0 eq) was added. After 2 h at -78°C, the reaction mixture was quenched with a 10% aqueous HCl and extracted with dichloromethane. The organic extracts were combined and evaporated to a clear oil. The oil was chromatographed (silica, 2% methanol/dichloromethane) to give the title compound as a white foam (0.248 g, 37%). NMR: (CDCl₃) δ 7.18 (5H, m), 4.57 (1H, d), 4.41 (1H, dd), 3.95 (1H, q), 2.82 (2H, d), 2.55 (2H, m), 2.49-1.49 (7H, m), 1.32 (9H, s); MS m/e 438.0 (M+Na)⁺.

b) (2R,4S,5S)-5-(t-butoxycarbonyl)amino-4-t-butyl-dimethyl-siloxy-6-phenyl-2-(4,4,4-trifluorobut-1-yl)hexanoic acid

Following the procedure of Example 12(b), except using the compound of Example 35(a) (245 mg), the title compound was prepared (215 mg, 67%). NMR(CDCl₃) δ 7.18 (5H, m), 4.70 (1H, d), 3.88 (1H, q), 3.69 (2H, m), 2.73 (1H, m), 2.38 (1H, m), 1.91 (2H, m), 1.45 (6H, m), 1.31 (9H, s) (rotamers observed), 0.90 (9H, s), 0.08 (6H, d); MS m/e 548.2 [M+H]⁺.

c) (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-t-butyl-dimethylsiloxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-(4,4,4-trifluorobut-1-yl)hexanamide

Following the procedure of Example 1(c), except using the compound of Example 35(b) (100 mg) and (1S)-1-imidazol-2-yl-2-methylpropylamine, the title compound was prepared (83 mg, 68%). NMR(CDCl₃) δ 7.22 (5H, m), 7.03 (1H, d), 6.89 (2H, s), 4.72 (1H, d), 4.51 (1H, t), 3.91 (1H, q), 3.65 (1H, m), 2.78 (2H, d), 2.33 (2H, m), 1.82 (4H, m), 1.48 (4H, m), 1.36 (9H,

two singlets; rotamers present), 0.99 (9H, s), 0.91 (3H, d), 0.79 (3H, d), 0.07, (6H, d); MS m/e 669.4 [M+H]⁺.

(2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-(4,4,4-trifluorobut-1-yl)hexanamide

Following the procedure of Example 9(d), except using the compound of Example 35(c) (83 mg), the title compound was prepared (40 mg, 58%). NMR(CD₃OD) δ 7.19 (5H, m), 6.92 (2H, s), 4.61 (1H, d), 3.64 (1H, q), 3.48 (1H, m), 2.79 (2H, m), 2.49 (1H, m), 2.13 (4H, m), 1.60 (5H, m), 1.36 (9H, s), 0.90 (3H, d), 0.71 (3H, d); MS m/e 555.2 [M+H]⁺.

Example 36

Preparation of (2R,4S,5S,1'S)-2-phenylmethyl-4-hydroxy-5-(t-butoxycarbonyl)amino-6-phenyl-N-(1'-isobutyl-1'-(imidazo-2-yl)methyl-hexanamide hydrochloride

20 (a) 2-(1'-carbobenzyloxyamino-1'-isobutyl)methyl-imidazole

Following the procedure of Example 1(a), except substituting Cbz-isoleucinal (1.83 g) for Cbz-valinal, the title compound was prepared (0.658 g, 31%). NMR(CDCl₃) δ 6.96 (2H, s), 5.31 (1H, d), 4.48 (1H, dd), 2.15 (1H, m), 1.44 (9H, s), 1.17 (2H, m), 0.92 (3H, t), 0.82 (3H, d); MS (DCI/NH₃) m/e 254.2 [M+H]⁺.

b) (2R,4S,5S,1'S)-2-phenylmethyl-4-hydroxy-5-(t-butoxycarbonyl)amino-6-phenyl-N-(1'-isobutyl-1'-(imidazo-2-yl)methyl-hexanamide hydrochloride

Following the procedure of Example 1(b)-1(d), except substituting the compound of Example 36(a) for (1'S)-1'-carbobenzyloxyamino-1'-isopropyl-1'-(imidazo-2-yl)methane, the title compound was prepared. NMR(DMSO-d₆) δ 7.90 (1H, d), 7.29-7.02 (10H, m), 6.89 (2H, s), 6.50 (1H, d), 4.81 (1H, m), 4.55 (1H, dd), 3.56 (1H, m), 2.69 (5H, m), 1.80 (1H, m), 1.59 (2H, m), 1.30 (9H, s), 1.17 (2H, m), 0.78 (3H, t), 0.63 (3H, d); MS (DCI/NH₃) m/e 549.7 [M+H]⁺.

Example 37

Preparation of (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-
hydroxy-N-(1'-isopropyl-1'-(4-((1RS)-1-hydroxyethyl)-
imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide

The t-butyldimethylsiloxy-protected alcohol from Example 30(e) (20 mg, 1.0 eq) was stirred in anhydrous THF under an argon atmosphere at room temperature. Tetrabutyl ammonium fluoride (0.33 mL of a 1.0M solution in THF, 6.0 eq) was added and the solution stirred for 16 h. The solution was diluted with water and extracted with dichloromethane. The combined organic extracts were washed with water and evaporated to a white solid. The solid was covered with diethyl ether and decanted twice to give the title compound as a white solid. (0.012 g, 72%). NMR(CDCl₃) δ 7.22-6.84 (10H, m), 6.61 (1H, s), 5.42 (1H, d), 4.69 (1H, m), 4.41 (1H, d), 3.58 (1H, m), 3.45 (1H, m), 2.78-2.40 (5H, m), 1.91 (1H, m), 1.59 (2H, m), 1.41 (3H, d), 1.26 (9H, s) (rotamers observed), 0.71 (3H, d), 0.59 (3H, dd); MS m/e 579.2 [M+H]⁺.

Example 38

Preparation of (2R,4S,5S,1'S)-5-(1,1-dimethyl-2-
hydroxyethoxycarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-
imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide

a) 2-t-butyldimethylsiloxy-1,1-dimethylethyl-(4-nitrophenyl) carbonate (0.996 g, 3.28 mmol), 2-t-butyldimethylsiloxy-1,1-dimethylethanol (0.67 g, 1 eq) and 4-dimethylaminopyridine (0.4 g, 1 eq) in dichloromethane (50 mL) was stirred at room temperature for 5 d. The mixture was diluted with dichloromethane and washed successively with H₂O and saturated aqueous NaCl, and dried over Na₂CO₃. The solvent was removed in vacuo, and the residue was purified by flash

- 79 -

chromatography (silica, 20% ethyl acetat /hexanes) to afford the title compound (35%). NMR(CDCl₃) δ 8.25 (2H, m), 7.35 (2H, m), 3.76 (2H, s), 1.53 (6H, s), 0.94 (9H, s), 0.09 (6H, s).

5

b) (2R,4S,5S,1'S)-5-(2-t-butyldimethylsiloxy-1,1-dimethylethoxycarbonyl)amino-4-t-butyldimethylsiloxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide

A solution of 2-t-butyldimethylsiloxy-1,1-dimethylethyl-4-nitrophenyl carbonate (137 mg, 0.372 mmol), (2R,4S,5S,1'S)-5-amino-4-t-butyldimethylsiloxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide (102 mg, 0.186 mmol) and DMAP (45 mg, 0.372 mmol) in methylene chloride was stirred at 20°C under Ar for 24 h. The solution was washed with aqueous Na₂CO₃, dried over solid Na₂CO₃ and concentrated.

Flash chromatography (4% methanol/dichloromethane) provided the intermediate (2R,4S,5S,1'S)-5-(2-t-butyldimethylsiloxy-1,1-dimethylethoxycarbonyl)amino-4-t-butyldimethylsiloxy-N-(1'-isopropyl-1'-(1-(2-t-butyldimethylsiloxy-1,1-dimethylethoxycarbonyl)imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide, which was dissolved in ether, washed with 10% NaOH, dried over Na₂CO₃, and concentrated to provide the title compound (110 mg, 78% overall). NMR(CDCl₃) δ 7.37-6.70 (13H, m), 6.39 (1H, d), 4.84 (1H, d), 4.55 (1H, t), 3.96 (1H, q), 3.69 (2H, s), 3.60-3.42 (2H, m), 2.94 (1H, s(br)), 2.85-2.44 (4H, m), 2.39 (1H, q), 1.90-1.60 (2H, m), 1.31 (6H, d), 1.02-0.85 (18H, m), 0.83 (6H, t), 0.98 (12H, m).

c) (2R,4S,5S,1'S)-5-(1,1-dimethyl-2-hydroxyethoxycarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide

A mixture containing the compound of Example 38(b) (110 mg) and tetra-n-butylammonium fluoride (6 eq of 1M solution in THF) under an argon atmosphere was allowed to stir at room temperature overnight. The solution was diluted with dichloromethane and washed with water, and the organic layer was concentrated. The residue was purified by flash chromatography (4% methanol/dichloromethane) to afford the

- 80 -

title compound (0.05 g, 66%). NMR(CDCl₃, CD₃OD) δ 7.30-6.78 (12H, m), 4.42 (1H, d), 3.75-3.38 (4H, m), 2.97-2.50 (5H, m), 2.08 (1H, m), 1.70-1.56 (2H, m), 1.30 (6H, s), 0.90-0.55 (6H, dd).

5

Example 39

Preparation of (2R,4S,5S,1'S)-5-(1,1-dimethyl-2-hydroxy-ethoxycarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide hydrochloride

10

A 1M solution of HCl in ether (63.5 mL) was added to a solution of the compound of Example 38(c) (35 mg, 0.064 mmol) in methanol (5 mL). The solvent was removed by rotary evaporation at 20°C, and the solid residue was triturated with ether and dried to afford the title compound as the hydrochloride salt (35 mg, 95%). NMR(CD₃OD) δ 7.37-6.85 (12H, m), 4.56 (1H, d), 3.59 (1H, m), 3.48-3.33 (3H, m), 2.85-2.48 (6H, m), 2.04 (1H, septet), 1.72-1.49 (2H, m), 1.22 (6H, d), 0.88 (3H, d), 0.61 (3H, dd).

15

Example 40

Preparation of (2R,4S,5S,1'S)-5-(2-hydroxyethoxycarbonyl)-amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethylhexanamide

25

a) benzyloxyethyl-(4-nitro)phenylcarbonate

To a solution of 2-benzyloxyethanol (2.5 g, 16.4 mmol) and bis(4-nitrophenyl)carbonate (5.0 g, 1 eq) in dichloromethane (200 mL), N-methylmorpholine (1.81 mL, 1 eq) was added. The resulting mixture was allowed to stir at room temperature for 3 d. The reaction mixture was washed successively with H₂O and saturated aqueous NaCl and dried over Na₂SO₄. The solvent was removed in vacuo, and the residue was purified by flash chromatography (silica, 20% ethyl acetate/hexanes) to afford the title compound (4.38 g,

30

35

- 81 -

(m) (b) 84%). ¹H NMR(CDCl₃) δ 8.26 (2H, m), 7.34 (7H, m), 4.62 (2H, s), 4.49 (2H, t), 3.70 (2H, t).

b) (2R,4S,5S,1'S)-5-(2-benzyloxyethoxycarbonyl)amino-4-t-butyl-
5 butyldimethylsiloxy-N-[1'-isopropyl-1'-(N'-2-benzyloxyethoxy) carbonyl)imidazol-2-yl]methyl-6-phenyl-2-phenylmethyl-hexanamide

To a solution of (2R,4S,5S,1'S)-5-amino-4-t-butyl-
butyldimethylsiloxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-
10 6-phenyl-2-phenylmethyl-hexanamide (134.5 mg, 0.24 mmol) in dichloromethane (40 mL) under an argon atmosphere, benzyloxyethyl 4-nitrophenyl carbonate (160 mg, 2 eq) and 4-dimethylaminopyridine (60 mg, 2 eq) were added. The
resulting mixture was allowed to stir at room temperature
15 overnight, and was diluted with dichloromethane. The organic extract was washed successively with aqueous Na₂CO₃, H₂O, aqueous Na₂CO₃ and H₂O, and dried over Na₂CO₃. The solvent was removed in vacuo, and the residue was purified by flash chromatography (silica, 4% methanol/dichloromethane) to
20 afford the title compound (180 mg, 82%). NMR(CDCl₃) δ 7.45-6.80 (22H, m), 6.62 (1H, d), 5.60 (1H, t), 5.06 (1H, d), 4.60 (2H, s), 4.52 (2H, s), 4.50 (2H, m), 4.31 (1H, m), 4.07 (2H, m), 3.80 (2H, t), 3.68 (1H, q), 3.57 (1H, q), 2.85 (1H, m), 2.77-2.41 (4H, m), 2.09 (1H, m), 1.90 (1H, m), 1.73 (1H, m),
25 0.95 (9H, s), 0.81 (6H, dd), 0.11 (6H, d).

c) (2R,4S,5S,1'S)-5-(2-hydroxyethoxycarbonyl)amino-4-t-butyl-
dimethylsiloxy-N-[1'-isopropyl-1'-(N'-2-benzyloxyethoxy-
carbonyl)imidazol-2-yl]methyl-6-phenyl-2-phenylmethyl-
30 hexanamide

The compound of Example 40(b) (68 mg, 0.44 mmol) was stirred as a solution in methanol (50 mL) with Pd(0) (10 mg) under 1 atm hydrogen for 12 h. The mixture was filtered, the solvent was removed in vacuo, and the residue was purified by
35 flash chromatography (silica, 4% methanol/dichloromethane) to afford the title compound (44 mg, 74%). NMR(CDCl₃) δ 7.36-6.72 (12H, m), 5.03 (1H, d), 4.80 (1H, dd), 4.50-4.32 (2H,

- 82 -

m), 4.07-3.52 (5H, m), 2.96-2.32 (6H, m), 1.98-1.85 (2H, m), 0.95 (9H, s), 0.90-0.75 (6H, dd), 0.05 (6H, d).

d) (2R,4S,5S,1'S)-5-(2-hydroxyethoxycarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide

To a solution of the compound of of Example 40(c) in methanol, excess aqueous HCl (approx. 5 equiv.) was added. The resulting solution was stirred at room temperature overnight, and concentrated under reduced pressure. The residue was diluted with H₂O, and made basic with aqueous Na₂CO₃. The mixture was extracted with dichloromethane, and the combined organic extracts were dried over Na₂CO₃. The solvent was removed in vacuo, and the residue was purified by flash chromatography to afford the title compound. NMR(CD₃OD) δ 7.28-6.85 (12H, m), 4.55 (1H, d), 3.95 (1H, m), 3.73-3.40 (4H, m), 2.86-2.47 (5H, m), 1.99 (1H, m), 1.71 (1H, m), 1.22 (1H, m), 0.84 (3H, d), 0.62 (3H, d).

Example 41

Preparation of (2R,4S,5S,1'S)-5-((1RS)-1-methyl-2-hydroxyethoxycarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide

a) 2-t-butyldimethylsiloxy-1-methylethyl-(4-nitrophenyl)-carbonate

A mixture containing bis(4-nitrophenyl)carbonate (3.20 g, 10.5 mmol), 2-t-butyldimethylsiloxy-1-methylethanol (2.0 g, 10.5 mmol) and 4-dimethylaminopyridine (1.30 g, 10.5 mmol) in dichloromethane (200 mL) was stirred at room temperature for 5 d. The mixture was then diluted with dichloromethane and washed successively with H₂O and saturated aqueous NaCl and dried over Na₂CO₃. The solvent was removed in vacuo, and the residue was purified by flash chromatography (silica, 10% ethyl acetate/hexane) to afford the title compound (88%). NMR(CDCl₃) δ 8.28 (2H, m), 7.39

- 83 -

1.38 (3H, s), 0.92 (9H, s), 0.11 (6H, s).

(b) (2R,4S,5S,1'S)-5-(2-*t*-butyldimethylsiloxy-1-methyl-ethoxycarbonyl)amino-4-*t*-butyldimethylsiloxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide

Following the procedure of Example 38(b), except substituting the compound of Example 4(a) for 2-*t*-butyldimethylsiloxy-1,1-dimethylethyl-4-nitrophenyl carbonate, the title compound was prepared. NMR(CDCl₃) δ 7.40-7.00 (10H, m), 6.90 (1/2H, s), 6.72 (1/2H, s), 6.45 (1H, dd), 4.92 (1H, dd), 4.84-4.61 (2H, m), 4.10 (1H, m), 3.76 (1H, m), 3.58 (1H, m), 2.92-2.73 (3H, m), 2.70-2.45 (3H, m), 1.78 (2H, m), 1.22-1.08 (3H, m), 1.04-0.81 (24H, m), 0.17-0.09 (12H, m).

(c) (2R,4S,5S,1'S)-5-((1R)-1-methyl-2-hydroxyethoxycarbonyl)-amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide

20 Following the procedure of Example 38(c), except using the compound of Example 4(b), the title compound is prepared. NMR(CD₃OD) δ 7.15-6.68 (12H, m), 5.72-5.60 (1H, dd), 4.58 (1H, m), 4.38 (1H, dd), 4.06 (1H, m), 3.62 (1H, m), 3.41 (1H, m), 2.79-2.55 (5H, m), 2.49 (1H, dd), 1.92 (1H, m), 1.67 (1H, m), 1.08-0.98 (3H, dd), 0.69 (3H, dd), 0.58 (3H, dd).

Example 42

Preparation of (2R,4S,5S,1'S)-5-(2-hydroxy-1-cyclopentylloxycarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethylhexanamide

(a) (trans)-2-(*t*-butyldimethylsiloxy)-cyclopentanol
To a mixture of *t*-butyldimethylsilyl chloride (5.08 g, 33.7 mmol) and imidazole (2.30 g, 33.7 mmol) in DMF (10 mL), a solution of trans-1,2-cyclopentanediol in DMF (4 mL) was added. The reaction mixture was stirred overnight at 25°C. The reaction mixture was diluted with ice water and extracted

with ether. The ether extract was washed with water and brine, dried over magnesium sulfate, filtered and the solvent removed in vacuo. The residue was purified by flash chromatography (silica, 9:1 hexane:ethyl acetate) to the title compound as an oil (3.44 g, 49%).

b) ((trans)-2-(t-butyldimethysiloxy)-cyclopentyl)-(4-nitrophenyl)carbonate

To a solution of the compound of Example 42(a) (1.08 g, 5 mmol) and DMAP (0.611 g, 5 mmol) in dichloromethane (12 mL), bis(4-nitrophenyl)carbonate (1.52 g, 5 mmol) was added. The solution was stirred overnight at 25°C. The reaction mixture was diluted with dichloromethane (15 mL) and washed with water and brine. The organic extract was dried over magnesium sulfate, filtered, and the solvent was removed at reduced pressure. The residue was triturated with hexane:ethyl acetate (1:1) and filtered. The filtrate was evaporated to an oil and purified by flash chromatography (silica, 9:1 hexane:ethyl acetate) to yield the title compound as an oil (1.75 g, 92%).

c) 5-((trans)-2-t-butyldimethylsiloxy-cyclopentyloxy-carbonyl)amino-4-t-butyldimethylsiloxy-N-[1'-isopropyl-1'-(1-(2-t-butyldimethylsiloxy-cyclopentyloxy-carbonyl))imidazol-2-yl]methyl-6-phenyl-2-phenylmethyl-hexanamide

A solution of 5-amino-4-t-butyldimethylsiloxy-N-[1'-isopropyl-1'-imidazol-2-yl]methyl-6-phenyl-2-phenylmethyl-hexanamide (171 mg, 0.311 mmol), DMAP (76.1 mg, 0.623 mmol) and the compound of Example 42(b) (238 mg, 0.623 mmol) in dichloromethane (9 mL) was stirred overnight at 25°C. The reaction mixture was diluted with dichloromethane, washed with water and saturated sodium bicarbonate solution, and dried with magnesium sulfate. The organic extract was filtered and the solvent was removed in vacuo. The residue was purified by flash chromatography (silica, 4:1 hexane:ethyl acetate) to yield the title compound as an oil (150 mg, 47%).

- 85 -

d) 5-((trans)-2-hydroxy-cyclopentylloxycarbonyl)amino-4-hydroxy-N-[1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide

To a solution of the compound of Example 42(c) (150 mg, 0.145 mmol) in methanol (5 mL), 3N HCl (3 mL) was added. The solution was stirred overnight at 25°C. The methanol was evaporated in vacuo, and the residue was diluted with water and extracted with ether. The aqueous solution was neutralized with 5% sodium carbonate (~pH 7) and a solid precipitated. The solid was filtered, washed with water and dried in vacuo to yield the title compound (51.5 mg, 63%).
 NMR(CD₃OD, 400 MHz): δ 7.0-7.3 (m, 10H), 6.87 (s, 2H), 4.63 (m, 2H), 3.88 (m, 1H), 3.55 (d, 1H), 2.5-2.9 m, 5H), 1.4-2.1 (br, 9H), 0.88 (d, 3H), 0.71 (d, 3H); TLC R_f 0.27 (silica, 8% methanol/chloroform).

Example 43

Preparation of (2R,4S,5S,1'S)-5-(4-hydroxybutanoyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethylhexanamide

a) t-butyldimethylsilyl 4-(t-butyldimethylsiloxy)-butanoate

To a suspension of t-butyldimethylsilyl chloride (29.9 g, 198 mmol) in dry DMF (20 mL), 4-hydroxybutyric acid, sodium salt (5.0 g, 397 mmol) and imidazole (27.0 g, 0.397 mol) were added. The reaction mixture was stirred overnight at 25°C. The solvent was removed under reduced pressure and the residue was diluted with 10% aqueous citric acid (200 mL). The residue was extracted with ether. The ether solution was dried with magnesium sulfate, filtered and evaporated to yield the title compound as an oil.

b) 4-t-butyldimethylsiloxy-butanoic acid

A solution of the compound of Example 43(a) (5.0 g) was dissolved in acetic acid:tetrahydrofuran:water (2:2:1, 50 mL) solution and stirred for 2.5 h. The solution was diluted with water and extracted with ether. The ether solution was

dried with magnesium sulfate, filtered and evaporated to an oil. The oil was purified by flash chromatography (silica, hexane-ethyl acetate, 9:1) to yield the title compound as an oil (180 mg).

c) (2R,4S,5S,1'S)-5-(4-t-butyldimethylsiloxy-butanoyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethylhexanamide

A solution of (2R,4S,5S,1'S)-5-amino-4-t-butyldimethylsiloxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide (175 mg, 0.319 mmol), 4-t-butyldimethylsiloxy-butanoic acid (84 mg, 0.41 mmol), BOP reagent (148, 0.335 mmol), triethylamine (46 μ L, 0.335 mmol) and dichloromethane (4 mL) were stirred at 20°C under Ar for 24 h. The reaction mixture was diluted with dichloromethane, washed with aqueous Na₂CO₃, water and brine, and dried over solid magnesium sulfate. The organic phase was filtered, and concentrated in vacuo. The residue was purified by flash chromatography (silica, 2% methanol/chloroform) to provide the title compound.

d) (2R,4S,5S,1'S)-5-(4-hydroxybutanoyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethylhexanamide

A solution of the compound of Example 43(c) (177 mg, 0.236 mmol) and tetra-n-butylammonium fluoride (2.84 mL, 2.84 mmol, 1M solution in THF) was stirred under an argon atmosphere at room temperature overnight. The solution was diluted with ethyl acetate, washed with saturated sodium bicarbonate solution, and water, and the organic layer was concentrated. The residue was precipitated from the ethyl acetate solution to afford the title compound. NMR δ (CD₃OD, 400 MHz) 7.0-7.3 (m, 10H), 6.86 (s, 2H), 4.62 (d, 1H), 4.05 (m, 1H), 3.43 (t, 2H), 2.55-2.90 (m, 4H), 2.60 (m, 1H), 2.17 (m, 2H), 2.05 (m, 1H), 1.76 (m, 1H), 1.67 (m, 2H), 1.55 (m, 1H), .88 (d, 3H), .72 (d, 3H); TLC R_f 0.40 (silica, 10% methanol/chloroform).

Example 44Preparation of (2R,4S,5S,1'S)-2-phenylmethyl-4-hydroxy-5-(benzyloxycarbonyl)valylamino-6-phenyl-N-(1'-isobutyl-1'-imidazo-2-yl)methyl-hexanamide

(a) (2R,4S,5S,1'S)-2-phenylmethyl-4-butyldimethylsiloxy-5-(benzyloxycarbonyl)valylamino-6-phenyl-N-(1'-isobutyl-1'-imidazo-2-yl)methyl-hexanamide.

10 A solution of carbobenzyloxy-(L)-valine (50.4 mg, 0.20 mmol), the product of Example 13(a) (110 mg, 0.20 mmol), BOP reagent (88.7 mg, 0.20 mmol) and triethylamine (28 μ l, 0.20 mmol) in methylene chloride (4 mL) was stirred at 25°C for 4 d. The reaction mixture was diluted with methylene chloride, washed with saturated sodium bicarbonate and the organic layer was concentrated. The product was purified by flash chromatography (silica gel, 4% CH₂Cl₂/MeOH) to give the title compound (104 mg, 67%).

20 (b) (2R,4S,5S,1'S)-2-phenylmethyl-4-hydroxy-5-(benzyloxycarbonyl-valyl)amino-6-phenyl-N-(1'-isobutyl-1'-imidazo-2-yl)methyl-hexanamide.

To a solution of the compound of Example 44(a) (104 mg, 0.133 mmol) in MeOH (8 mL), 3N HCl (2 mL) was added. The solution was stirred for 16 hrs at 25°C. The methanol was removed at reduced pressure and 10% sodium carbonate was added to pH ~7.5. Ether (10 mL) was added and the solid product was filtered and dried in vacuo to provide the title compound (58 mg, 65%). NMR(CDCl₃) δ : 0.62 (d, 3H), 0.78 (d, 3H), 0.82 (d, 3H), 0.90 (d, 3H), 1.62 (m, 2H), 1.96 (m, 1H), 2.06 (m, 1H), 2.55 (m, 1H), 2.77 (m, 4H), 3.38 (s, 1H), 3.53 (m, 1H), 3.91 (m, 1H), 3.99 (m, 1H), 4.47 (d, 1H), 5.11 (s, 2H), 5.78 (d, 1H), 6.85 (s, 2H), 6.92-7.34 (m, 15H); MS m/e 667 [M+H]⁺.

Example 45

Preparation of (2R,4S,5S,1'S)-2-phenylmethyl-4-hydroxy-5-(N-
acetylvalyl)amino-6-phenyl-N-(1'-isobutyl-1'-imidazo-2-
5 yl)methyl-hexanamide

(a) (2R,4S,5S,1'S)-2-phenylmethyl-4-*t*-butyldimethylsilyloxy-5-(N-acetyl-valyl)amino-6-phenyl-N-(1'-isobutyl-1'-imidazo-2-yl)methyl-hexanamide

10 To a solution of N-acetyl-(L)-valine (40.3 mg, 0.253 mmol) in dry THF (8 mL) at -40°C was added *n*-methylmorpholine (55.7 µL, 0.506 mmol) followed by isobutyl chloroformate (33.5 µL, 0.253 mmol). The reaction mixture was stirred for 15 min, and the compound of Example 13(b) (139 mg, 0.253 mmol) in THF (3 mL) was added. The reaction mixture was allowed to warm to room temperature and stirred for 2 d. The reaction was diluted with ethyl acetate, and washed with water and brine. The organic solution was dried with sodium sulfate, filtered and the solvent removed under reduced
15 pressure. The residue was purified by flash chromatography (silica, 4% methanol/chloroform) to give the product as an oil (47 mg, 27%).

(b) (2R,4S,5S,1'S)-2-phenylmethyl-4-hydroxy-5-(N-acetylvalyl)amino-6-phenyl-N-(1'-isobutyl-1'-imidazo-2-yl)methyl-hexanamide.

To a solution of the compound of Example 45(a) (47 mg, 0.0681 mmol) in methanol (3 mL), 3N HCl (0.5 mL) was added. The reaction was stirred for 16 h at 25°C. The methanol was removed under reduced pressure and the solution was diluted with water and neutralized with 5% sodium carbonate. The solid product was filtered, washed with water and ether, and dried in vacuo to yield the title compound (29.5 mg, 75%).
20 NMR (CD₃OD) δ 0.70 (d, 3H), 0.88 (m, 9H), 1.57 (m, 1H), 1.70 (m, 1H), 1.92 (s, 3H), 2.05 (m, 1H), 2.55 (q, 1H), 2.77 (m, 4H), 3.57 (d, 1H), 4.03 (m, 2H), 4.60 (d, 1H), 6.87 (s, 2H), 6.95-6.20 (m, 10H); MS m/e 575 [M+H]⁺.

8. You require to show that when $\alpha = 1$, Example 46 reduces to Example 45.

business as usual to support the U.S. mission. The

#5: Preparation of (2R,4S,5S,1'S)-5-[imidazol-2-

(yl)methyloxycarbonyllamino-4-hydroxy-N-(1'-isopropyl

5 imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexa

[illegible]

a) 4-(1-(benzyloxymethyl)imidazol-2-yl)methyl-

nitrophenyl) carbonate

A mixture of bis(4-nitrophenyl)carbonate, (1-

10 benzyloxymethyl)imidazol-2-yl)methanol and 4-

dimethylaminopyridine was reacted according to the

of Example 14(a) to afford the title compound (58%):

NMR (CDCl₃, 400 MHz) δ 8.18 (d, 2 H, J=8.38 Hz), 7.

6.67 (s, 1H), 7.11 (s, 1H), 7.13 (s, 1H), 5.48 (s, 2H), 5.4

[illegible][illegible]

b) (2R,4S,5S,1'S)-5-((1-benzyloxymethyl)imidazol-2-

yl)methyloxycarbonyl) amino-4-t-butyl dimethylsiloxane

4. 1-(2-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenyl

20. hexanamide: CCCCCCC(=O)N

A mixture of the compound of Example 46(a),

(11) (b) (2R,4S,5S,1'S)-5-amino-4-hydroxy-N-(1'-isopropyl-1

26 (2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide, an

dimethylaminopyridine was reacted according to the

25 of Example 14(b) to afford the title compound (32%

NMR (CDCl₃) δ 7.50-6.60 (m, 19H), 5.25 (m, 2H), 5.1

J=11.03 Hz), 4.68 (m, 1H), 4.39 (m, 2H), 3.97 (m,

(m, 1H), 2.88 (m, 1H), 2.72-2.28 (m, 6H), 1.85 (m,

δ 7.12 (m, 1H), 6.92-6.81 (m, 15H), 6.80 (s, 3H), 0.06 (s,

MS (ES) $m/e = 793$ [M+H]⁺.

oblongness of the leaf, the leaf is not so long as the leaf of the other species.

c) (2R, 4S, 5S, 1'S)-5-(imidazol-2-yl-

methyloxycarbonyl) amino-4-t-butyl dimethylsiloxy-N-

isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenyl

-1) 35D-hexanamide (C₁₆H₃₃N) (molecular weight 255.44)

The compound of Example 46(b) (58 mg, 0.073 mmol) was dissolved in 10 mL of 1,2-dichloroethane and 10 mL of 10% aqueous sodium hydroxide solution. The mixture was stirred at room temperature for 24 hours. The organic layer was separated, washed with water, and dried over anhydrous sodium sulfate. The solvent was removed under reduced pressure to give a white solid (45 mg, 78% yield).

methanol (3 mL), and 10% Pd on carbon (50 mg) were

and stirred under 1 atm of H_2 for 24 h. Additiona

- 90 -

(50 mg) was added and stirring under H₂ was continued for 8 h. The reaction was filtered through Celite®, concentrated and flash chromatographed (silica, step gradient, 0-8% MeOH/CH₂Cl₂) to yield the title compound (28 mg, 57%).

5 NMR(CDCl₃) δ 7.29-6.83 (m, 14H); 5.05 (d, 1H, J=11.2 Hz), 4.91 (d, 1H, J=11.2 Hz), 4.71 (m, 1H), 3.92 (m, 1H), 3.61 (m, 1H), 3.02 (m, 1H), 2.81-2.54 (m, 4H), 2.36 (m, 1H), 1.93 (m, 1H), 1.59 (m, 1H), 0.91 (d, 3H, J=7.1 Hz), 0.89 (s, 9), 0.69 (d, 3H, J=7.1 Hz), 0.84-0.05 (m, 6H); MS(ES) m/e 673 [M+H]⁺.

10

d) (2R,4S,5S,1'S)-5-(imidazol-2-yl-methyloxycarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide

The compound of Example 46(c) (24 mg, 0.035 mmol), 95% aqueous EtOH (0.50 mL), and concentrated aqueous HCl (0.050 mL) were stirred at 23°C for 24 h. The solution was diluted with H₂O (5 mL), washed with EtOAc and then the aqueous phase was made basic by addition of solid K₂CO₃. Extraction with EtOAc, concentration of the organic extract and trituration

20 with CH₂Cl₂ afforded the title compound (14 mg, 72%). NMR (CDCl₃) δ 7.33-6.85 (m, 14H), 5.11 (d, 1H, J=10.8 Hz), 4.96 (d, 1H, J=10.8 Hz), 4.47 (m, 1H), 3.72 (m, 1H), 3.38 (m, 1H), 2.81 (m, 4H), 2.59 (m, 1H), 2.07 (m, 1H), 1.72 (m, 1H), 1.62 (m, 1H), 0.78 (d, 3H, J=6.63 Hz), 0.67 (d, 3H, J=6.63 Hz); 25 (m, 6H); MS(ES) m/e 559 [M+H]⁺.

Example 47

30 Preparation of (2R,4S,5S,1'S,1"RS)-5-((1"-((imidazol-2-yl)-2"-methyl)propyloxycarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide

a) (1RS)-1-((1-benzyloxymethylimidazol-2-yl)-2-methyl)propyl-(4-nitrophenyl)carbonate

35 A mixture of bis(4-nitrophenyl)carbonate, (1RS)-1-((1-benzyloxymethylimidazol-2-yl)-2-methyl)propanol and 4-dimethylaminopyridine was reacted according to the procedure of Example 14(a) to afford the title compound (61%). NMR

- 91 -

(CDCl₃) δ 8.18 (d, 2H, J=8.31 Hz), 7.38-7.21 (m, 7H), 7.13 (s, 1H), 6.94 (s, 1H), 5.74 (d, 1H, J=11.1 Hz), 5.47 (d, 1H, J=10.2 Hz), 5.28 (d, 1H, J=10.2 Hz), 4.53 (d, 1H, J=11.3 Hz), 4.41 (d, 1H, J=11.3 Hz), 2.64 (m, 1H), 1.18 (d, 3H, J=6.02 Hz), 0.87 (d, 3H, J=6.02 Hz); MS(ES) m/e 426 [M+H]⁺.

b) (2R,4S,5S,1'S,1"RS)-5-((1"-((1-benzyloxymethylimidazol-2-yl)-2"-methyl-propyl)oxycarbonyl)amino-4-t-butylidimethylsiloxy-N-(1'-isopropyl-1'-(1-(1"-((1-benzyloxymethylimidazol-2-yl)-2"-methylpropyl)oxycarbonyl)imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide

A mixture of the compound of Example 47(a) (145 mg, 0.33 mmol), the compound of Example 13(a) (75.9 mg, 0.14 mmol), 4-dimethylaminopyridine (41 mg, 0.33 mmol) and DMF (0.5 mL) was stirred under argon for 18 h. The DMF was evaporated in vacuo and the residue was combined with 10% aq K₂CO₃ (10 mL) and extracted with EtOAc. The combined extracts were washed with saturated aq NaHCO₃, dried (K₂CO₃), filtered and concentrated in vacuo. The residue was flash chromatographed (silica, step gradient, 0-4% MeOH/CH₂Cl₂) to afford the title compound (96.1 mg, 57%). NMR (CDCl₃) δ 7.38-6.78 (m, 26H), 5.67 (m, 1H), 5.61-5.00 (m, 6H), 4.58-4.27 (m, 5H), 3.97-3.61 (m, 3H), 2.78-2.10 (m, 8H), 1.95-1.51 (m, 2H), 1.10-0.55 (m, 27H), 0.50-0.05 (m, 6H).

c) (2R,4S,5S,1'S,1"RS)-5-[(1"-((1-benzyloxymethylimidazol-2-yl)-2"-methyl-propyl)oxycarbonyl]amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide

A solution of the compound of Example 47(b) (81 mg, 0.07 mmol), CH₃OH (0.75 mL), and 3N aqueous HCl (0.25 mL) was stirred at 23°C for 20 h. The reaction mixture was diluted with H₂O (10 mL) and washed with EtOAc (3 x 15 mL). Solid K₂CO₃ was added to give a basic solution (pH>12), which was extracted with EtOAc. The extracts were dried (K₂CO₃), filtered, concentrated and flash chromatographed (silica, step gradient, 0-8% CH₃OH/CH₂Cl₂) to give the title compound

- 92 -

(34.9 mg, 65%). ^1H NMR (CDCl_3) δ 7.43-6.79 (m, 9H), 5.87, 5.66 (2d, 1H, $J=10.66$, 10.85 Hz), 5.28 (m, 2H), 4.68 (m, 1H), 4.42 (m, 2H), 3.71 (m, 1H), 3.58 (m, 1H), 2.90-2.31 (m, 6H), 2.11 (m, 1H), 1.75, 1.51 (2m, 2H), 1.05, 0.97 (2d, 3H, $J=6.32$, 6.45), 0.68 (m, 9H).

d) (2R,4S,5S,1'S,1"RS)-5-((1"-(imidazol-2-yl)-2"-methyl)propyloxycarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide.

10 A mixture of the compound of Example 47(c) (34 mg, 0.047 mmol), CH_3OH (4 mL), and 10% Pd/C (34 mg), was stirred under H_2 (1 atm) for 26 h. The suspension was filtered through Celite®, concentrated, and triturated with CH_2Cl_2 to yield the title compound (4 mg, 14%). ^1H NMR ($\text{CDCl}_3/\text{CD}_3\text{OD}$) δ 7.732-6.71 (m, 14H), 5.38 (m, 1H), 4.55 (m, 1H), 3.72 (m, 1H), 3.55 (m, 1H), 2.78 (m, 4H), 2.55 (m, 1H), 2.15 (m, 2H), 1.60 (m, 2H), 1.03-0.61 (m, 12H).

Example 48

Preparation of (2R,4S,5S,1'S)-5-(*t*-butoxycarbonyl)amino-4-hydroxy-N-[1'-isopropyl-1'-(4-(imidazol-2-yl)imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide]

25 (a) (1'S)-1'-(carbobenzyloxy)amino-1'-isopropyl-1'-(4-(imidazol-2-yl)imidazol-2-yl)methane

Cbz-(L)-valinal (0.45 g, 1.4 mmol) was stirred in anhydrous methanol at 0°C under argon. Glyoxal (40% in water) (0.22 mL, 1.4 mmol) and ammonium hydroxide (29% NH_3) (0.88 mL, 14 mmol) were added and the mixture was allowed to stir at 0°C for 1 h. The cooling bath was removed and the solution stirred at room temperature for 16 h. The methanol was evaporated in vacuo and the residue was diluted with 5% aqueous HCl. After extracting with dichloromethane, the aqueous layer was made basic with solid sodium carbonate and extracted with dichloromethane. The combined organic extracts were dried over sodium carbonate, filtered, and evaporated to a solid which was chromatographed (silica, 4%

- 93 -

methanol/dichloromethane) to give the title compound (0.216 g, 43%) as a white solid. NMR (CDCl₃) δ 7.15 (6H, s(br)), 6.88 (2H, s), 6.30 (1H, d), 4.89 (2H, dd), 4.52 (1H, t), 2.05 (1H, m), 0.73 (3H, d), 0.62 (3H, d). MS m/e 340.2 [M+H]⁺

5

(b) (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-[1'-isopropyl-1'-(4-(imidazol-2-yl)imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide]

The compound of Example 48(a) (0.13 gm.) was dissolved in anhydrous methanol with 10% Pd on activated carbon (0.02 g). Hydrogen gas was bubbled through the solution via a balloon for 1 h and the solution was stirred overnight under a hydrogen atmosphere. The mixture was filtered through a pad of Celite® and evaporated to yield 1'-amino-1'-isopropyl-15 [4-(imidazol-2-yl)imidazol-2-yl]methane as a white solid (0.13 g, 100%).

This compound was combined with the compound of Example 13(a) (0.334 g, 0.63 mmol), BOP reagent (0.28 g, 0.63 mmol), and triethylamine (0.13 mL, 0.945 mmol) in DMF (1 mL) and allowed to stir under Ar for 3 d. The DMF was evaporated in vacuo and the residue was diluted with dichloromethane. The solution was washed with water and brine. The organic layer was dried over sodium carbonate, filtered, and evaporated to yield (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-t-butyl-25 dimethylsiloxy-N-[1'-isopropyl-1'-(4-(imidazol-2-yl)imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide as a white solid.

A portion of the solid (0.100 g, 0.14 mmol) was stirred in THF at room temperature under argon. Tetrabutylammonium fluoride (0.84 mL, 0.84 mmol) was added and the mixture was allowed to stir for 16 h. The solution was diluted with water and extracted twice with dichloromethane. The combined organic extracts were washed with water and evaporated to an oily residue. The residue was dissolved in THF and several drops of diethyl ether were added until a white precipitate formed. The precipitate was collected by filtration and dried in vacuo to yield the title compound as a white solid (76 mg, 90%). NMR (CD₃OD) δ 7.37-6.84 (13H, m), 4.61 (1H,

35

d), 3.69 (2H, m), 3.54 (1H, d), 2.84-2.52 (5H, m), 2.06 (1H, m), 1.83 (2H, m), 1.57 (1H, m), 1.30 (9H, s), 0.87 (3H, d), 0.69 (3H, d); MS m/e 601.2 [M+H]⁺

5

Example 49

Preparation of (2R,4S,5S,1'S)-5-[di(hydroxymethyl)-methoxycarbonyllamino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide

10

a) di(t-butyldimethylsiloxymethyl) methyl-(4-nitrophenyl) carbonate

A mixture containing bis(4-nitrophenyl) carbonate (1.89 g, 6.21 mmol), di(t-butyldimethylsiloxymethyl)methanol (2.00 g, 1 eq) and 4-dimethylaminopyridine (757 mg, 1 eq) in dichloromethane (100 mL) was stirred at room temperature for 2 d. The mixture was diluted with dichloromethane and washed with saturated aqueous Na₂CO₃, brine, and dried over Na₂SO₄.

The solvent was removed in vacuo, and the residue was purified by flash chromatography (silica, 10% ethyl acetate/hexanes) to afford the title compound (75%). NMR (CDCl₃) δ 8.29 (2H, m), 7.37 (2H, m), 3.96 (1H, m), 3.85 (2H, d), 3.82 (2H, d), 0.89 (18H, s), 0.09 (12H, s).

25 b) (2R,4S,5S,1'S)-5-(di(t-butyldimethylsiloxymethyl)methyloxycarbonyl)amino-4-t-butyldimethylsiloxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide

A solution of di(t-butyldimethylsiloxymethyl)-methyl 4-nitrophenyl carbonate (475 mg, 0.974 mmol), the compound of Example 13(a) (178 mg, 0.325 mmol) and dimethylaminopyridine (119 mg, 0.974 mmol) in methylene chloride was stirred at 20°C under Ar for 24 h. The solution was washed with aqueous Na₂CO₃, dried over solid Na₂CO₃ and concentrated in vacuo.

35 Flash chromatography (silica, 4% methanol/dichloromethane) of the residue provided the intermediate (2R,4S,5S,1'S)-5-(di(t-butyldimethylsiloxymethyl) methyloxycarbonyl)amino-4-t-butyldimethylsiloxy-N-(1'-isopropyl-1'-(1-(di(t-

butyldimethylsiloxymethyl) methyloxycarbonyl)imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide, which was dissolved in ether, washed with 10% NaOH, dried over Na₂CO₃, and concentrated to provide the title compound (197 mg, 71%).

5 NMR (CDCl₃) δ 7.43-7.05 (10H, m), 6.90 (2H, s), 6.65 (1H, bs), 5.09 (1H, d), 4.78 (1H, bd), 4.08 (1H, m), 3.89-3.50 (7H, m), 3.00-2.80 (4H, m), 2.65 (1H, m), 2.55 (2H, m), 1.90 (1H, m), 1.78 (1H, m), 1.10-0.85 (33H, m), 0.20-0.06 (18H, m).

10 c) (2R,4S,5S,1'S)-5-(di(hydroxymethyl)methoxycarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide

A mixture containing the compound of Example 49(b) (50 mg) and ethereal HCl (4 eq) was allowed to stir in methanol:water (9:1) at room temperature overnight. The solvent was removed in vacuo, and the residue was diluted with ethyl acetate and washed with saturated aqueous Na₂CO₃. The product was purified by flash chromatography (silica, 4% 20 methanol/dichloromethane) to afford the title compound (29 mg, 94%). NMR (CD₃OD) δ 7.20-6.80 (10H, m), 6.71 (2H, s), 4.50 (1H, d), 3.90 (1H, m), 3.65-3.34 (5H, m), 2.82-2.45 (6H, m), 1.99 (1H, m), 1.74 (1H, m), 1.52 (1H, m), 0.78 (3H, d), 0.60 (3H, d).

Example 50

Preparation of (2R,4S,5S,1'S)-5-(1-oxo-thian-4-yl)oxycarbonylamino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethylhexanamide

30 Reacting the compound of Example 32(b) (81 mg, .133mmol) with m-chloro perbenzoic acid (23 mg, 0.133mmol) in CH₂Cl₂ yielded the title compound. NMR (CD₃OD) δ 7.20-6.85 (10H, m), 6.78 (2H, s), 4.51 (1H, d), 3.66 (1H, m), 3.42 (1H, m), 2.95-2.41 (9H, m), 2.32-2.01 (2H, m), 1.99-1.63 (4H, m), 1.60-1.41 (2H, m), 0.78 (3H, d), 0.60 (3H, d); MS m/e 595.2 [M+H]⁺.

- 96 -

Example 51

Preparation of (2R,4S,5S,1'S)-5-((tetrahydrosulfonylpyran-4-yl)oxycarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethylhexanamide

Reacting the compound of Example 50 (31 mg, 49.2 μ mol) with m-chloro perbenzoic acid (10 mg, 59.2 μ mol) in methylene chloride yielded the title compound. NMR (CD₃OD) δ 7.20-6.85 (10H, m), 6.76 (2H, s), 4.48 (1H, d), 3.68 (1H, m), 3.44 (1H, m), 2.96-2.42 (9H, m), 2.32-2.04 (2H, m), 1.97-1.62 (4H, m), 1.61-1.43 (2H, m), 0.79 (3H, d), 0.60 (3H, d); MS m/e 611.2 [M+H]⁺.

Example 52

Preparation of (2R,4S,5S,1'S)-5-(1,1-dimethyl-2-(acetoxymethoxycarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide

(a) (2R,4S,5S,1'S)-5-(1,1-dimethyl-2-(acetoxymethoxycarbonyl)amino-4-(t-butyldimethylsilyl)oxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide

The compound of Example 38(b) (223 mg, 0.221 mmol) was dissolved in 10% aqueous methanol and combined with 1M HCl in ether (0.221 mL, 1 eq) at room temperature. After completion of the reaction the solvents were removed in vacuo. The residue was dissolved in dichloromethane and washed with aqueous saturated Na₂CO₃. The organic layer was concentrated and the residue was purified by flash chromatography (silica, 4% methanol/dichloromethane) to provide the title compound (138 mg, 94%). NMR (CDCl₃) δ 7.38-6.81 (12H, m), 4.93 + 4.65 (1H, d, rotamers), 4.81 + 4.48 (1H, t, rotamers), 4.15 + 4.08 (1H, d, rotamers), 3.90 (1H, q), 3.72 (2H, m), 3.50 + 3.38 (1H, d, rotamers), 2.98-2.48 (5H, m), 2.35 (1H, m), 1.98 (1H, m), 1.79 (1H, m), 1.60 (1H, m), 1.30 (3H, s), 1.29 (3H, s), 1.09 -0.85 (15H, m), 0.79 (3H, d), 0.11 (6H, m).

(b) (2R,4S,5S,1'S)-5-((1,1-dimethyl-2-
 20) (benzoacetoxyethoxycarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-
 imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide
 25 The compound of Example 52(a) (103 mg, 0.155 mmol) was
 stirred with acetic anhydride (30 mg, 0.309 mmol) and DMAP
 (40 mg, 0.309 mmol) in methylene chloride at room temperature
 under argon overnight. The solvent was removed in vacuo and
 the residue was flash chromatographed (silica, 4%
 methanol/dichloromethane).

10 The resulting 4-t-butyldimethylsiloxy intermediate (105
 mg, 0.140 mmol) was stirred in methanol:water (9:1) with 1M
 HCl in ether (0.14 mL, 1 eq). The solvents were removed in
 vacuo, the residue was diluted with dichloromethane, and the
 solution was washed with aqueous Na₂CO₃. The organic layer
 15 was concentrated and the residue was purified by flash
 chromatography (silica, 5% methanol/dichloromethane) to
 provide the title compound (82 mg, 91%). NMR (CD₃OD) δ 7.29-
 6.90 (10H, m), 6.81 (2H, s), 4.51 (1H, d), 4.05 (2H, s),
 3.59 (1H, m), 3.42 (1H, m), 2.80-2.45 (5H, m), 2.00 (1H, m),
 1.98 (3H, s), 1.72 (1H, m), 1.50 (1H, m), 1.34 (6H, d), 0.81
 (m) (3H, d), 0.60 (3H, d).

25 The compound of Example 52(a) (100 mg, 0.151 mmol) was

Example 53

25 Preparation of (2R,4S,5S,1'S)-5-((1,1-dimethyl-2-(benzyloxy-
 carbonyl)glycyloxy)ethoxycarbonyl)amino-4-hydroxy-N-(1'-
 isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-
 hexanamide hydrochloride salt.

30 a) (2R,4S,5S,1'S)-5-((1,1-dimethyl-2-(benzyloxy-
 carbonyl)glycyloxy)ethoxycarbonyl)amino-4-(t-
 butyldimethylsilyloxy)-N-(1'-isopropyl-1'-imidazol-2-
 yl)methyl-6-phenyl-2-phenylmethyl-hexanamide

The compound of Example 52(a) (100 mg, 0.151 mmol) was
 35 reacted with 2-chloro-1-methyl-pyridium iodide (92 mg, 0.36
 mmol), DMAP (75 mg, 0.60 mmol) and Cbz-glycine (63 mg, 0.30
 mmol) in methylene chloride (5 mL) under argon at reflux for
 3 h. Solvents were removed in vacuo and the product was

purified by flash chromatography (silica, 40% methanol/dichloromethane) to provide the title compound (95 mg, 73%). NMR (CDCl₃) δ 7.41-6.71 (17H, m), 6.62 (1H, bs), 6.00 (1H, m), 5.20 (1H, m), 5.15 (2H, us), 4.83+4.55 (1H, d, 5 rotamers), 4.65+4.48 (1H, t, rotamers), 4.81+4.38 (1H, q, rotamers), 4.03 (1H, q), 4.02 (2H, d), 3.85+3.68 (2H, d, rotamers), 2.85-2.48 (5H, m), 2.38 (1H, m), 1.90 (1H, m), 1.55 (1H, m), 1.38 (3H, s), 1.29 (3H, s), 0.90 (9H, m), 0.85 (3H, d), 0.70 (3H, d), 0.11 (6H, m).

10 b) (2R,4S,5S,1'S)-5-((1,1-dimethyl-2-(benzyloxycarbonyl)ethoxycarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide hydrochloride salt

15 The compound of Example 53(a) (12 mg, 0.014 mmol) was stirred in methanol:water (9:1) with 1M HCl (2 eq) in ether overnight. The solvents were removed in vacuo to give the title compound (8 mg, 73%). NMR (CD₃OD) δ 7.35 (2H, s), 7.31-6.85 (15H, m), 5.00 (2H, s), 4.59 (1H, d), 4.15 (1H, d, 20 rotamers), 4.65+4.48 (1H, t, rotamers), 4.81+4.38 (2H, dd), 3.80 (2H, d), 3.59 (1H, m), 3.40 (1H, d), 2.85-2.48 (5H, m), 2.00 (1H, m), 1.60 (1H, m), 1.55 (1H, m), 1.31 (3H, s), 1.29 (3H, s), 0.91 (3H, d), 0.60 (3H, d).

25 Example 54
Preparation of (2R,4S,5S,1'S)-5-((1,1-dimethyl-2-(glycyloxy)ethoxycarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide
dihydrochloride salt

30 a) (2R,4S,5S,1'S)-5-((1,1-dimethyl-2-(glycyloxyethoxycarbonyl)amino-4-(t-butyldimethylsilyl)oxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide

35 The compound of Example 53(a) (58 mg, 0.0678 mmol) was stirred in methanol with 10% Pd/C (50 mg) under 1 atm hydrogen overnight. The reaction mixture was filtered

- 99 -

through C lite® and the solvents were removed *in vacuo* to yield the title compound (48 mg, 98%). NMR(CD₃OD) δ 7.32-7.02 (10H, m), 6.99 (2H, s), 4.68 (1H, d), 4.40-4.28 (2H, dd), 3.81 (2H, d), 3.80-3.67 (2H, m), 2.90-2.49 (5H, m), 2.15 (1H, m), 1.97 (1H, m), 1.48 (1H, m), 1.40 (3H, s), 1.39 (3H, s), 1.15 (3H, d), 0.95 (9H, s), 0.70 (3H, d), 0.11 (6H, d).

b) (2R,4S,5S,1'S)-5-((1,1-dimethyl-2-glycyloxy)ethoxy-carbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide dihydrochloride salt

The compound of Example 54(a) (43.5 mg, 0.060 mmol) was stirred in methanol:water (9:1) with 1M HCl in ether (0.12 mL, 2 eq) for 2 d. The solvents were removed *in vacuo* and the product was triturated with ether:methanol (20:1) to yield the title compound (40 mg, 98%). NMR(CD₃OD) δ 7.35 (2H, s), 7.30-6.92 (10H, m), 4.60 (1H, d), 4.25 (2H, dd), 3.75 (2H, d), 3.59 (1H, m), 3.49 (1H, m), 2.90-2.51 (6H, m), 2.10 (1H, m), 1.65 (1H, m), 1.54 (1H, m), 1.30 (6H, s), 0.90 (3H, d), 0.60 (3H, d).

Example 55

Preparation of (2R,4S,5S,1'S)-5-((1,1-dimethyl-2-hydroxy)ethoxycarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-(4-isopropylcarbonylimidazol-2-yl))methyl-6-phenyl-2-phenylmethyl-hexanamide dihydrochloride salt

a) (2R,4S,5S,1'S)-5-amino-4-t-butyldimethylsiloxy-N-[1'-isopropyl-1'-(4-isopropylcarbonyl-imidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide

Using the procedure of Example 13(a), except substituting the compound of Example 28(d), the title compound was prepared.

b) (2R,4S,5S,1'S)-5-((1,1-dimethyl-2-hydroxy)ethoxycarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-(4-

isopropylcarbonylimidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide dihydrochloride salt

Following the procedures of Example 38(b)-38(c), except substituting the compound of Example 55(a) for

- 5 (2R,4S,5S,1'S)-5-amino-4-t-butyldimethylsiloxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide, the title compound was prepared. NMR (CDCl₃) δ 7.49 (1H, s), 7.13 (5H, m), 6.84 (5H, m), 5.53 (1H, d), 4.47 (1H, d), 3.79 (1H, m), 3.60 (1H, m), 3.44 (2H, m), 3.16 (1H, m), 2.81-2.50 (5H, m), 1.92 (1H, m), 1.62 (2H, m), 1.18 (14H, m), 0.72 (3H, d), 0.58 (3H, d); MS m/e 621.4 [M+H]⁺.

Example 56

- 15 Preparation of (2R,4S,5S,1'S)-5-((1S)-1-methyl-2-hydroxyethoxycarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethylhexanamide

Using the procedure of Example 41, except substituting

- 20 2(S)-t-butyldimethylsiloxy-1-methylethanol in 41(a) (prepared from 2(S)-1,2-propanediol), the title compound was prepared. NMR(CD₃OD) δ 7.38-6.90 (10H, m), 6.83 (2H, s), 4.58 (2H, m), 3.61 (1H, m), 3.34 (3H, m), 2.82-2.44 (5H, m), 2.00 (1H, m), 1.66 (1H, m), 1.52 (1H, m), 1.08 (3H, d), 0.85 (3H, d), 0.60 (3H, d).

Example 57

- 30 Preparation of (2R,4S,5S,1'S)-5-((1R)-1-methyl-2-hydroxyethoxycarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethylhexanamide

Using the procedure of Example 41, except substituting 2(R)-t-butyldimethylsiloxy-1-methylethanol in 41(a), the title compound was prepared. NMR(CD₃OD) δ 7.39-6.88 (10H,

- 35 m), 6.82 (2H, s), 4.56 (2H, m), 3.60 (1H, m), 3.36 (3H, m), 2.81-2.45 (5H, m), 1.99 (1H, m), 1.65 (1H, m), 1.51 (1H, m), 1.03 (3H, d), 0.84 (3H, d), 0.60 (3H, d).

Example 58

Preparation of (2R,4S,5S,1'S)-5-((1-acetyl-amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl)hexanamide

The title compound was prepared by the procedure of Example 13(a)-(c), except substituting acetic anhydride in place of isopropyl chloroformate. NMR(CD₃OD) δ 7.21-6.90 (10H, m), 6.81 (2H, s), 4.58 (1H, d), 3.98 (1H, m), 3.51 (1H, m), 2.85-2.49 (5H, m), 1.99 (1H, m), 1.68 (3H, s), 1.61 (3H, m), 1.50 (1H, m), 0.80 (3H, d), 0.60 (3H, d).

Example 59

Preparation of (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-(4-benzyloxyphenylmethyl) hexanamide

a) (3R,5S,1'S)-(1'-t-butoxycarbonylamino-2'-phenyl) ethyl-3-(20 (4-benzyloxy)phenylmethyl-tetrahydrofuran-2-one. Following the procedure of Example 12(a), except using (4-benzyloxy)benzyl bromide, the title compound was prepared (284 mg, 27%). NMR(CDCl₃) δ 7.48-6.72 (14H, m), 4.94 (2H, s), 4.43 (1H, d), 4.12 (1H, dd), 3.83 (1H, q), 2.97-2.62 (5H, 25 m), 2.12 (1H, m), 1.85 (1H, m), 1.27 (9H, s).

b) (2R,4S,5S)-5-(t-butoxycarbonyl)amino-4-t-butyl dimethyl-siloxy-6-phenyl-2-(4-benzyloxyphenylmethyl)hexanoic acid

Following the procedure of Evans et al., J. Org. Chem. 50, 4615 (1985), except substituting the compound of Example 59(a) for benzyl bromide, the title compound was prepared. NMR(CDCl₃) δ 7.42-6.76 (14H, m), 4.99 (2H, s), 4.69 (1H, d), 3.91 (1H, q), 3.66 (1H, m), 2.98-2.36 (5H, m), 1.85 (1H, m), 1.52 (1H, m), 1.30 (9H, s), 0.88 (9H, s), 0.04 (6H, m).

c) (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-t-butyl dimethylsiloxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-(4-benzyloxyphenylmethyl) hexanamide

- 102 -

Following the procedure of Example 12(c), except using (b), the title compound was prepared (284 mg, 92%).

NMR(CDCl₃) δ 7.42-6.74 (16H, m), 5.04 (2H, s), 4.99 (1H, d), 4.77 (1H, d), 4.51 (1H, dd), 3.93 (1H, q), 3.69 (1H, m), 2.80-2.39 (5H, m), 1.81 (1H, m), 1.62 (1H, m), 1.33 (9H, s), 0.92 (9H, s), 0.75 (6H, dd), 0.07 (6H, d).

(d) (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-(4-benzyloxyphenylmethyl)hexanamide

Following the procedure of 12(d), except using (c), the title compound was prepared (100 mg, 94%). NMR(CD₃OD) δ 7.41-7.09 (10H, m), 6.85 (2H, d), 6.79 (2H, s), 6.58 (2H, d), 5.41 (1H, d), 4.90 (2H, s), 4.47 (1H, d), 3.62 (1H, q), 3.48 (1H, d), 2.79-2.48 (6H, m), 2.02 (1H, m), 1.62 (2H, m), 1.33 (9H, s), 0.74 (3H, d), 0.61 (3H, d).

Example 60

20 Preparation of (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-(4-benzyloxyphenylmethyl)hexanamide

Following the procedure of Example 4(b), except using the compound of 59(d), the title compound was prepared (56 mg, 86%). NMR(CD₃OD) δ 7.18 (5H, m), 6.84 (2H, s), 6.73 (2H, d), 6.44 (2H, d), 5.32 (1H, d), 4.45 (1H, d), 3.61 (1H, q), 3.42 (1H, m), 2.80-2.42 (5H, m), 2.04 (1H, m), 1.61 (2H, m), 1.31 (9H, s), 0.70 (3H, d), 0.61 (3H, d).

30 Preparation of (2R,4S,5S)-5-(t-butoxycarbonyl)amino-4-hydroxy-2-phenylmethyl-6-phenyl-N-[1'-cyclopropyl-1'-imidazol-2-yl]methyl-hexanamide

35 Preparation of (2R,4S,5S)-5-(t-butoxycarbonyl)amino-4-hydroxy-2-phenylmethyl-6-phenyl-N-[1'-cyclopropyl-1'-imidazol-2-yl]methyl-hexanamide

a) α -(t-butoxycarbonyl)-amino- α -cyclopropylacetonitrile

To a solution of cyclopropylmethanol (10.2 g, 141 mmol) in methylene chloride (250 mL) sodium acetate (1 g) and 20 g of Celite® were added. Pyridinium chlorochromate (30 g, 140 mmol) was added in small portions over a period of 30 min. After 1 h the reaction mixture was diluted with ether (100 mL), filtered through Celite® and washed with ether. The combined organic extracts (1 L) were concentrated in vacuo at 15-18°C to yield formyl cyclopropane.

The crude aldehyde was dissolved in water (50 mL), and ammonium chloride (6.51 g), potassium cyanide (7.16 g) and aqueous ammonium hydroxide (100 mL, 28% w/w). The reaction mixture was stirred at room temperature overnight, extracted with ethyl acetate, and the combined organic extracts were dried over MgSO₄. Filtration and evaporation of the solvent in vacuo yielded α -amino- α -cyclopropyl acetonitrile as an oil.

To a solution of the crude aminonitrile (2 g) in THF (20 mL) di-tert-butyl dicarbonate (1.53 g, 7 mmol) was added. The reaction was stirred overnight. Removal of the solvent in vacuo followed by flash chromatography (silica, 1:8 ethyl acetate:hexane) yielded the title compound (2.8 g). ¹H NMR (CDCl₃, 200 MHz) δ 5.0 (bs, 1H), 4.4 (bs, 1H), 1.4 (s, 9H), 1.2 (m, 1H), 0.7 (m, 2H), 0.5 (m, 2H).

(b) α -(t-butoxycarbonyl)-amino- α -cyclopropylacetaldehyde

To a solution of the compound of Example 61(a) (1 g, 5.1 mmol) in THF (20 mL), diisobutylaluminum hydride (10.5 mL, 10.5 mmol, 1M in THF) was added at -78°C, over 5 min. The reaction mixture was allowed to warm to 0°C over a period of 30 min, and stirred at 0°C for an additional 1 h. The reaction mixture was quenched with MeOH (2 mL), and saturated potassium sodium tartrate solution (100 mL) was added.

Extraction with ether, drying over MgSO₄ and removal of solvents in vacuo yielded an oil. Flash chromatography (silica, 1:10 ethyl acetate:hexane) gave the title compound as a colorless solid (225 mg). ¹H NMR (CDCl₃, 400 MHz) δ 9.45 (bs, 1H), 4.95 (bs, 1H), 3.5 (bs, 1H), 1.3 (s, 9H), 0.7 (m, 1H), 0.3-0.6 (m, 4H).

c) 1-(t-butoxycarbonyl)amino-1-(imidazol-2-yl)-1-cyclopropyl-methane

A mixture of the compound of Example 61(b) (178 mg, 0.89 mmol), glyoxal (150 mL, 1 mmol, 40% aq), ammonium hydroxide (5 mL, 28% aq) and MeOH (5 mL) was stirred at room temperature for 10 h. The solvents were removed in vacuo and the residue was titrated with ether to yield a brown solid (53 mg). The solid was passed through Florisil® and eluted with 5% MeOH/methylene chloride. Removal of the solvent in vacuo followed by trituration provided the title compound as a colorless solid (19 mg). MS(CI/NH₃) m/e 238.3, [M+H]⁺. ¹H NMR(CD₃OD, 200 MHz) δ 6.9 (s, 2H), 4.1 (bd, 1H), 1.4 (s, 9H), 1.3 (m, 1H), 0.6 (m, 2H), 0.4 (m, 2H).

d) 1-amino-1-(imidazol-2-yl)-1-cyclopropyl-methane, trifluoroacetate

The compound of Example 61(c) (15 mg) was dissolved in 1 mL of TFA and stirred at room temperature for 20 min.

Solvents removed in vacuo to give the title compound as a semisolid residue. ¹H NMR(CD₃OD, 200 MHz) δ 7.1 (s, 2H), 3.8 (d, 1H, J=7 Hz), 1.5 (m, 1H), 0.5-0.8 (m, 4H).

e) (2R,4S,5S)-5-(t-butoxycarbonyl)amino-4-(t-butyldimethyl)siloxy-2-phenylmethyl-6-phenyl-N-[1'-cyclopropyl-1'-imidazol-2-yl]methyl-hexanamide

The compound of Example 61(d) was dissolved in DMF (2 mL) and NMM (26 mg, 0.25 mmol) was added and the solution was stirred at 0°C for 30 min. (2R,4S,5S)-2-phenylmethyl-4-(t-butyldimethyl)siloxy-5-(t-butoxycarbonyl)amino-6-phenyl hexanoic acid (38 mg, 0.07 mmol) and BOP reagent (30 mg, 0.07 mmol) were added and the reaction was stirred at room temperature for 24 h. The reaction was diluted with ethyl acetate (100 mL), washed with aqueous sodium bicarbonate and dried over anhydrous potassium carbonate. Removal of solvents in vacuo, followed by flash chromatography (silica, 5% methanol/methylene chloride) yielded the title compound as a mixture of diastereomers (25 mg).

f) (2R,4S,5S)-5-(t-butoxycarbonyl)amino-4-hydroxy-2-phenylmethyl-6-phenyl-N-[1'-cyclopropyl-1'-imidazol-2-yl]methyl-hexanamide

5 The compound of Example 61(c) was dissolved in THF (2 mL) and tetrabutyl ammonium fluoride (200mL, 1M in THF) was added. The reaction was stirred at room temperature overnight and methylene chloride (100 mL) and water (10 mL) were added. The organic layer was dried over potassium carbonate, and the solvent was removed in vacuo to give an oil. Flash chromatography (silica, 5% methanol/methylene chloride) gave a colorless solid which was a 1:1 diastereomeric mixture of the title compound.

15 Example 62

Preparation of (2R,4S,5S,1'R)-5-(t-butoxycarbonyl)amino-4-hydroxy-2-phenylmethyl-6-phenyl-N-[1'-cyclopropyl-1'-imidazol-2-yl]methyl-hexanamide; and
 20 (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-2-phenylmethyl-6-phenyl-N-[1'-cyclopropyl-1'-imidazol-2-yl]methyl-hexanamide

a) Purification of the 125 mg of the compound of Example
 25 61(e) by flash chromatography (silica, 3% methylene chloride/methanol), yielded 48 mg of isomer 1, 15 mg of isomer 2 and 20 mg of combined fractions. ¹H NMR for isomer 1 (CDCl₃, 400 MHz) δ 7.1-7.4 (m, 10H), 6.95 (s, 2H), 6.1 (d, 1H), 4.85 (d, 1H), 4.15 (dd, 1H), 3.75 (q, 1H), 3.6 (m, 1H), 2.9 (dd, 1H), 2.7 (d, 2H), 2.6 (dd, 1H), 2.3 (m, 1H), 2.0 (m, 1H), 1.6 (m, 1H), 1.4 (m, 1H), 1.35 (s, 9H), 0.95 (s, 9H), 0.7 (m, 1H), 0.4 (m, 1H), 0.25 (m, 1H), 0.1 (m, 1H), 0.2 (s, 3H), 0.1 (s, 3H). ¹H NMR for isomer 2 (CDCl₃, 400 MHz) δ 7.1-7.4 (m, 10H), 6.8 (s, 2H), 6.26 (d, 1H), 4.6 (d, 1H), 4.0 (m, 2H), 2.5-3.0 (m, 4H), 1.8 (m, 1H), 1.7 (m, 1H), 1.5 (m, 1H), 1.4 (s, 9H), 1.0 (s, 9H), 0.7 (m, 2H), 0.2 (m, 2H), 0.1 (2 overlapping singlets, 6H).

b) Following the procedure of Example 61(f), except substituting the compounds of Example 62(a) yielded the title compounds. ^1H NMR for isomer 1 (CD_3OD , 400 MHz) δ 7.1-7.3 (m, 10H), 6.95 (s, 2H), 4.25 (d, 1H), 3.5-3.7 (m, 2H), 2.5-3.0 (m, 5H), 1.7 (m, 2H), 1.4 (s, 9H), 1.1 (m, 1H), 0.6 (m, 1H), 0.25-0.4 (m, 2H), 0.05 (m, 1H); MS (ESMS) m/e 533.2 $[\text{M}+\text{H}]^+$; ^1H NMR for isomer 2 (CD_3OD) δ 7.1-7.4 (m, 10H), 6.85 (s, 2H), 4.25 (d, 1H), 3.5-3.7 (m, 2H), 2.5-2.9 (m, 5H), 1.5-1.8 (m, 2H), 1.4 (s, 9H), 1.1 (m, 1H), 0.2-0.6 (m, 4H); MS (ESMS) m/e 533.4 $[\text{M}+\text{H}]^+$.

Example 63

Preparation of (2R,4S,5S,1'S)-5-((isopropylthiol)carbonyl)-amino-4-hydroxy-2-phenylmethyl-6-phenyl-N-[1-isopropyl-1'-imidazol-2-yl]methyl-hexanamide

a) 5-((isopropylthiol)carbonyl)amino-4-(t-butyldimethylsiloxy)-2-phenylmethyl-6-phenyl-N-[1'-isopropyl-1'-(1-(isopropylthiol)carbonyl-imidazol-2-yl)]methyl-hexanamide

To a solution of (2R,4S,5S,1'S)-5-amino-4-t-butyldimethylsiloxy-2-phenylmethyl-6-phenyl-N-[1'-isopropyl-1'-imidazol-2-yl]methyl-hexanamide (81 mg, 148 μmol) and DMAP (37 mg, 303 μmol) in dichloromethane (8 mL), isopropylthiolchloroformate (42 mg, 303 μmol) in dichloromethane (1 mL) was added. The solution was stirred for 20 h and an additional equivalent of the chloroformate and DMAP were added. The reaction mixture was stirred for an additional 20 h, diluted with dichloromethane, and washed with saturated sodium bicarbonate. The organic extract was dried over magnesium sulfate, filtered and evaporated to an oil. The oil was dissolved in chloroform and purified by flash chromatography (silica, 1% methanol/chloroform) to give the title compound as an oil (79.5 mg).

b) (2R,4S,5S,1'S)-5-((isopropylthiol)carbonyl)amino-4-hydroxy-2-phenylmethyl-6-phenyl-N-[1-isopropyl-1'-imidazol-2-yl)methyl-hexanamide

To a solution of the compound of Example 63(a) (79 mg, 105 μ mol) in methanol (8 mL), 10% hydrochloric acid (3 mL) was added. The reaction mixture was stirred overnight at 25°C. The methanol was evaporated in vacuo, and the residue was diluted with water. The solution was neutralized with 5% aqueous sodium carbonate, and a solid precipitated. The solid was filtered, washed with water, and triturated with ether. The solid was dried at high vacuum to yield the title compound (27 mg, 48%). NMR(CDCl₃, 250 MHz) δ 6.9-7.3 (m, 10H), 6.85 (s, 2H), 6.20 (d, 1H), 4.42 (d, 1H), 4.22 (m, 1H), 4.0 (m, 1H), 3.55 (m, 3H), 2.5-3.0 (m, 6H), 1.65 (t, 2H), 1.27 (m, 7H), 1.71 (d of d, 6H); MS (FAB) m/e 537 [M+H]⁺; TLC R_f 0.30 (silica, 4% methanol/chloroform).

Example 64

20. Preparation of (2R,4S,5S,1'S)-5-(1-hydroxymethyl-cyclopentylloxycarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide

a) 1-(t-butyldimethylsiloxy)methyl-cyclopentanol
25 To a solution of 1-hydroxymethyl-1-cyclopentanol (4.07 g, 0.035 mole) in dichloromethane (30 mL) t-butyldimethylsilyl chloride (5.28 g, 0.035 mol) in dichloromethane (30 mL) was added. Triethylamine (5.37 mL, 0.0385 mol) and DMAP (0.171 g, 0.0014 mol) were added and the solution was stirred overnight at 25°C. The solution was diluted with dichloromethane (30 mL) and washed with water and saturated ammonium chloride solution. The organic solution was dried over sodium sulfate, filtered and the solvent removed at reduced pressure. The product was purified by flash 35 chromatography (silica, 19:1 hexane:ethyl acetate) to yield the title compound as a colorless oil (6.95 g, 86%).

b) 1-(t-butyldimethylsiloxy)methyl-cyclopentyl 4-nitrophenyl carbonate

A solution of the compound of 64(a) (1.15 g, 5 mmol), DMAP (0.611 g, 5 mmol) and bis(4-nitrophenyl) carbonate (1.52 g, 5 mmol) in dichloromethane (16 mL) was stirred overnight at 25°C. The reaction mixture was diluted with dichloromethane and washed with 5% sodium carbonate. The solvent was removed at reduced pressure and the residual oil was triturated with hexane:ethyl acetate (3:2) and filtered. The product was purified by flash chromatography (silica, 19:1 hexane:ethyl acetate) to give a colorless oil (0.599 g, 30%).

c) (2R,4S,5S,1'S)-5-[1-(t-butyldimethylsiloxy)methyl-15-cyclopentylloxycarbonyl]amino-4-t-butyldimethylsiloxy-N-[1'-isopropyl-1'-(t-butyldimethylsiloxy)methyl-cyclopentylloxy]imidazol-2-yl]-6-phenyl-2-phenylmethyl-hexanamide

A solution of the compound of Example 13(a) (173 mg, 0.316 mmol), DMAP (81 mg, 0.663 mmol) and the compound of Example 64(b) (262 mg, 0.663 mmol) in dichloromethane (10 mL) was stirred for 48 h at 25°C. The organic solution was diluted with dichloromethane, washed with 5% sodium carbonate solution and the solvent removed at reduced pressure. The product was purified by flash chromatography (silica, 4:1 hexane:ethyl acetate) to yield the title compound as an oil (200 mg, 60%).

d) (2R,4S,5S,1'S)-5-(1-hydroxymethyl-cyclopentylloxy-carbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide

A solution of the silylated derivative (200 mg, 0.188 mmol) in methanol (7 mL) and 3N HCl (2.5 mL) was stirred overnight at 25°C. The methanol was removed at reduced pressure and the solution was diluted with water (15 mL) and extracted with ether (25 mL). The aqueous solution was neutralized with 5% sodium carbonate solution to pH 7-7.5 and the product precipitated as a solid. The solid was filtered,

- 109 -

washed with water and dried in vacuo to yield the title compound (51 mg, 47%). NMR (CD₃OD, 400 MHz) δ 7.0-7.3 (m, 10H), 6.87 (s, 2H), 4.62 (d, 1H), 3.70 (m, 3H), 3.55 (d, 1H), 2.5-2.9 (m, 5H), 2.05 (m, 1H), 1.5-2.0 (br, 10H), 0.88 (d, 3H), 0.70 (d, 3H); TLC R_f 0.50 (silica, 8% methanol/chloroform).

Example 65

10 Preparation of (2R,4S,5S,1'S)-5-[3-(R)-(1H-imidazol-2-yl)-3-hydroxy-4-methylpentylamidol-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide; and
15 (2R,4S,5S,1'S)-5-[3-(S)-(1H-imidazol-2-yl)-3-hydroxy-4-methylpentylamidol-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide.

(35) a) 1-(1-benzyloxymethylimidazol-2-yl)-2-methyl-1-propanol
1-benzyloxymethylimidazole prepared according to the procedure of Ngochindo, R., J. Chem. Res. (S), 58 (1990) (3.76 g, 20 mmol), and THF (40 mL) at -40°C, was treated dropwise with n-BuLi (8.4 mL, 21 mmol, 2.5M in hexane). The resulting solution was stirred at -40°C for 15 min, and i-butylaldehyde (2.0 mL, 22 mmol) was added dropwise. The reaction was stirred at -40°C for 1.5 h, 0°C for 1 h, warmed to 23°C, poured into H₂O, and extracted with EtOAc. The combined extracts were washed with brine, dried (Na₂SO₄) and concentrated in vacuo. Trituration of the residue with Et₂O/hexane gave a white solid which was dried in vacuo overnight to afford of the title compound (3.57 g, 69%).
30 NMR(CDCl₃, 400 MHz) δ 7.28 (m, 5H), 6.97 (s, 1H), 6.92 (s, 1H), 5.23 (d, 1H, J=12 Hz), 5.42 (d, 1H, J=12 Hz), 4.48 (s, 2H), 4.44 (d, 1H, J=9 Hz), 2.21 (m, 1H), 1.02 (d, 3H, J=7 Hz), 0.83 (d, 3H, J=7 Hz).

(35) b) 1-(benzyloxymethylimidazol-2-yl)-2-methyl-propan-1-one
The compound of Example 65(a) (1.0 g, 3.88 mmol), MnO₂, (1.69 g, 19.4 mmol), and CH₂Cl₂ (75 mL) were combined and stirred for 1 d. Additional MnO₂ (1.69 g, 19.4 mmol) was

- 110 -

added and stirring was continued for an additional 2 d. Filtration through Celite®, concentration and flash chromatography (silica, 0-1% CH₃OH/CH₂Cl₂) afforded the title compound (0.773 g, 77%). ¹H NMR(CDCl₃, 400 MHz) δ 7.28 (m, 5 6H), 7.18 (s, 1H), 5.85 (s, 2H), 4.52 (s, 2H), 3.94 (m, 1H), 1.21 (d, 2H, J=5 Hz).

c) t-butyl 3-(1-benzyloxymethylimidazol-2-yl)-3-hydroxy-4-methyl-pentanoate

10 Diisopropylamine (83 μL, 0.59 mmol) and THF (1.5 mL) were cooled to -40°C and n-BuLi (188 μL, 0.47 mmol, 2.5M in hexane) was added. The reaction mixture was warmed to -10°C and stirred for 15 m, recooled to -70°C and t-butyl acetate (63 μL, 0.47 mmol) was added. The reaction was stirred for 5 15 m, and HMPA (254 μL, 1.41 mmol) was added. The reaction was stirred at -70°C for 5 m and 1-(benzyloxymethylimidazol-2-yl)-2-methyl-propan-1-one (100 mg, 0.39 mmol) in THF (1.5 mL) was added dropwise. The mixture was stirred at -70°C for 30 20 m, -40°C for 30 m, -10°C for 30 m, warmed to 23°C, poured into 10% aqueous K₂CO₃ and extracted with EtOAc. The combined organic extracts were washed with brine, dried (K₂CO₃), 25 concentrated and flash chromatographed (silica gel, step gradient, 0-20% EtOAc/hexanes) to afford the title compound (131 mg, 90%). ¹H NMR(CDCl₃, 400 MHz) δ 7.25 (m, 5H), 6.96 (s, 1H), 6.91 (s, 1H), 5.69 (d, 1H, J=10 Hz), 5.65 (d, 1H, J=10 Hz), 4.53 (d, 1H, J=11 Hz), 4.48 (d, 1H, J=11 Hz), 3.23 (d, 1H, J=6 Hz), 2.57 (d, 1H, J=6 Hz), 2.14 (m, 1H), 1.39 (s, 9H), 0.97 (d, 3H, J=7 Hz), 0.75 (d, 3H, J=7 Hz); MS(ES) m/e 375 [M+H]⁺.

30

d) 3-(1-benzyloxymethylimidazol-2-yl)-3-hydroxy-4-methyl pentanoic acid trifluoroacetate

The compound of Example 65(c) (93 mg, 0.24 mmol) was dissolved in TFA (1 mL) and stirred for 20 m. The TFA was 35 removed in vacuo to give the title compound (102 mg, 100%). ¹H NMR(CDCl₃, 400 MHz) 7.30 (m, 7H), 6.06 (d, 1H, J=9 Hz), 5.74 (d, 1H, J=1 Hz), 4.67 (d, 1H, J=9 Hz), 4.61 (d, 1H, J=9 Hz), 3.62 (d, 1H, J=12 Hz), 2.93 (d, 1H, J=12 Hz), 2.04 (m,

- 111 -

1H), 0.92 (d, 3H, J=12 Hz), 0.88 (d, 3H, J=12 Hz); MS(ES) m/e 319 [M+H]⁺.

(e) (2R,4S,5S,1'S)-5-[3-(RS)-(1-benzyloxymethylimidazol-2-yl)-3-hydroxy-4-methylpentanoyl]amino-4-t-butyl-dimethylsilyloxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide

A mixture of the compound of Example 65(d) (1.0 eq) (2R,4S,5S,1'S)-5-amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide (1.1 eq), BOP reagent (1.1 eq), and triethylamine (4 eq) were reacted according to the procedure of Example 1(c). The product was purified by flash chromatography to afford the title compound (57%) (silica, step gradient, 0-4% CH₃OH/CH₂Cl₂). ¹H

NMR(CDCl₃, 400 MHz) δ 7.36-6.76 (m, 19H), 5.65 (m, 2H), 4.66 (m, 1/2H), 4.51 (m, 2H), 4.39 (m, 1/2H), 4.30 (m, 1/2H), 4.02 (m, 1/2H), 3.68 (m, 1H), 3.28 (m, 1H), 2.90-2.35 (m, 6H), 2.13 (m, 1H), 1.76 (m, 1/2H), 1.68 (m, 1/2H), 1.40 (m, 1/2H), 1.00-0.70 (m, 21H), 0.10-0.00 (m, 6H); MS(ES) m/e 849 [M+H]⁺.

(f) (2R,4S,5S,1'S)-5-[3-(RS)-(1-benzyloxymethylimidazol-2-yl)-3-hydroxy-4-methylpentanoyl]amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide

The compound of Example 65(e) (100 mg, 0.12 mmol) was desilylated by the procedure of 47(c) to cleanly afford the title compound (78 mg, 89%). ¹H NMR(CDCl₃, 400 MHz) δ 7.40-6.80 (m, 19H), 5.75 (m, 2H), 4.97 (m, 1/2H), 4.78 (m, 1/2H), 4.51 (m, 2H), 3.94 (m, 1/2H), 3.85 (m, 1/2H), 3.51 (m, 1H), 3.21 (m, 1H), 2.97-2.43 (m, 6H), 2.00 (m, 1H), 1.60 (m, 1H), 1.43 (m, 1H), 0.97-0.49 (m, 12H); MS(ES) m/e 735 [M+H]⁺.

(g) (2R,4S,5S,1'S)-5-[3(R)-(imidazol-2-yl)-3-hydroxy-4-methylpentanoyl]amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide; and

(h) (2R,4S,5S,1'S)-5-[3-(S)-(imidazol-2-yl)-3-hydroxy-4-methylpentanoyl]amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide

Using the procedure of Example 47(d), the compound of Example 65(f) (72 mg, 0.98 mmol) was hydrogenated to afford a diastereomeric mixture of the title compounds. The mixture was purified by flash chromatography (silica, step gradient, 0-8% CH₃OH/CH₂Cl₂) to afford tail fractions containing the pure diastereomers (35 mg total, 58%).

Isomer 1, last eluting, (2R,4S,5S,1'S)-5-[3-(R)-(1H-Imidazol-2-yl)-3-hydroxy-4-methylpentylamido]-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide. ¹H NMR(CDCl₃, 400 MHz) δ 7.35-6.82 (m, 10H), 6.93 (s, 1H), 6.84 (s, 1H), 4.42 (d, 1H, J=9 Hz), 3.77 (m, 1H), 3.40 (m, 1H), 3.00-2.40 (m, 5H), 2.14 (m, 1H), 1.99 (m, 1H), 1.56 (m, 1H), 1.47 (m, 1H), 0.93-0.64 (m, 12H); MS(ES) m/e 615 [M+H]⁺.

Isomer 2, first eluting, (2R,4S,5S,1'S)-5-[3(S)-(1H-Imidazol-2-yl)-3-hydroxy-4-methylpentylamido]-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide. ¹H NMR(CDCl₃, 400 MHz) δ 7.35-6.81 (m, 10H), 6.83 (s, 1H), 6.81 (s, 1H), 4.46 (d, 1H, J=9 Hz), 3.93 (m, 1H), 3.40 (m, 1H), 3.00-2.40 (m, 5H), 2.13 (m, 1H), 1.91 (m, 1H), 1.41 (m, 1H), 1.10 (m, 1H), 0.93-0.64 (m, 12H); MS(ES) m/e 615 [M+H]⁺.

Example 66

Preparation of (2R,4S,5S,1'S)-5-[(4-methoxyphenoxy)carbonyl]amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide

(1a) (2R,4S,5S,1'S)-5-[(4-methoxyphenoxy)carbonyl]amino-4-t-butylldimethylsiloxo-N-[1'-isopropyl-1'-(N'-methoxycarbonyl)imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide

Following the procedure of Example 13(b), except using p-methoxyphenyl chloroformate and (2R,4S,5S,1'S)-5-amino-4-t-butylldimethylsiloxo-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide (114 mg, 0.21 mmol), the title compound was prepared (63%). NMR(CDCl₃), δ 7.44-6.76

(m, 20H), 5.66 (m, 1H), 5.18 (d, 1H), 4.40 (m, 1H), 3.83 (s, 3H), 3.76 (m, 1H), 3.73 (s, 3H), 2.96-2.50 (m, 5H), 2.05 (m, 5H), 1.60 (m, 1H), 0.94 (s, 9H), 0.79 (d, 3H, J=7 Hz), 0.74 (s, 3H), 0.12 (s, 3H), 0.11 (s, 3H).

b) (2R,4S,5S,1'S)-5-(methoxycarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide

Following the procedure of Example 13(c), except using the compound of Example 66(a), the title compound was prepared (32%). NMR(CDCl₃/CD₃OD), δ 7.36-6.84 (m, 16H), 4.49 (d, 1H, J=9 Hz), 3.79 (s, 3H), 3.37 (m, 1H), 2.92-2.60 (m, 5H), 2.10-1.70 (m, 3H), 0.78 (d, 3H, J=7 Hz), 0.67 (d, 3H, J=7 Hz); MS(ES) m/e 585 [M+H]⁺.

Example 67

Preparation of (2R,4S,5S,1'S)-5-(t-butylaminocarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenylmethyl-hexanamide

a) (2R,4S,5S,1'S)-5-(t-butylaminocarbonyl)amino-4-(t-butyltrimethylsiloxy)-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenylmethyl-hexanamide

The compound of Example 13(a) (0.13 g, 0.24 mmol) was dissolved in dichloromethane (3 mL) and t-butyl isocyanate (0.028 g, 0.29 mmol) was added. After stirring at 30°C for 18 h, the solvent was removed under reduced pressure and the residue was chromatographed (silica, 2:3 ethylacetate:hexane) to give the title compound as a white solid (0.12 g, 77%). NMR(CDCl₃), δ 7.35-7.05 (12H, m), 6.85 (2H, s), 4.69 (1H, d, J=9 Hz), 4.60 (1H, t, J=8 Hz), 4.38 (1H, br), 4.24 (1H, q, J=8 Hz), 3.66 (1H, dd, J=4 Hz, 10 Hz), 2.95 (1H, dd, J=9 Hz, 13 Hz), 2.73 (2H, m), 2.54 (1H, dd, J=5 Hz, 13 Hz), 2.42 (1H, m), 1.82 (1H, m), 1.67 (1H, m), 1.22 (9H, s), 0.93 (9H, s), 0.84 (3H, d, J=7 Hz), 0.79 (3H, d, J=7 Hz), 0.08 (3H, s), 0.07 (3H, s); MS(ES) m/e 648.4 [M+H]⁺.

b). (2R,4S,5S,1'S)-5-(t-butylaminocarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenylmethyl-hexanamide.

The compound of Example 67(a) (0.0338g, 0.05 mmol) was stirred in dry THF (0.25 mL) and tetrabutylammonium fluoride (0.25 mL, 0.25 mmol) in THF was added. After 18 h at 50°C the reaction was cooled, diluted with ethyl acetate (25 mL), washed with water (5 mL), and dried (MgSO₄). The combined organic extracts were filtered and concentrated in vacuo.

Chromatography (silica, 1:1 ethyl acetate:hexane) gave the title compound as a white solid (0.018 g, 66%). M.p 226°C (dec); NMR(CD₃OD) δ 7.37-6.90 (10H, m), 6.90 (2H, s), 4.58 (1H, d, J=9 Hz), 3.71 (1H, t, J=7 Hz), 3.52 (1H, d, J=9 Hz), 2.75 (4H, m), 2.53 (1H, dd, J=4 Hz, 12 Hz), 2.03 (1H, m), 1.76 (1H, m), 1.66 (1H, m), 1.22 (9H, s), 0.79 (3H, d, J=7 Hz), 0.67 (3H, d, J=7 Hz); MS(ES) m/e 534 [M+H]⁺.

Example 68

20. Preparation of (2R,4S,5S,1'S)-5-(methylaminocarbonyl)-amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenylmethyl-hexanamide.

Following the procedure of Examples 67(a)-67(b), except substituting methyl isocyanate for t-butylisocyanate, the title compound was prepared (0.075 mg, 51%). M.p 253°C (dec); NMR(DMSO-d₆) δ 7.78 (1H, d, J=9 Hz), 7.80-6.96 (11H, m), 6.88 (2H, s), 5.78 (1H, d, J=5 Hz), 5.72 (1H, d, J=9 Hz), 4.84 (1H, d, J=4 Hz), 4.65 (1H, m), 3.68 (1H, q, J=7 Hz), 3.44 (1H, br), 2.74 (3H, m), 2.58 (1H, dd, J=7 Hz, 13 Hz), 2.50 (3H, s), 2.41 (1H, d, J=8 Hz), 1.92 (1H, m), 1.46 (2H, m), 0.72 (3H, d, J=7 Hz), 0.63 (3H, d, J=7 Hz); MS(ES) m/e 492 [M+H]⁺.

Example 69

35. Preparation of (2R,4S,5S,1'S)-5-(phenylaminocarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenylmethyl-hexanamide.

- 115 -

Following the procedure of Examples 67(a)-67(b), except substituting phenyl isocyanate for t-butylisocyanate, the title compound was prepared (87 mg, 79%). Mp 273°C (dec); NMR(DMSO-d₆) δ 8.50 (1H, s), 7.81 (1H, d, J=9 Hz), 7.34-6.83 (18H, m), 6.07 (1H, d, J=9 Hz), 4.99 (1H, d, J=4 Hz), 4.65 (1H, t, J=8 Hz), 3.75 (1H, m), 3.52 (1H, br), 2.77 (3H, m), 2.66 (1H, m), 2.42 (1H, d, J=7 Hz), 1.89 (1H, m), 1.50 (2H, m), 0.68 (3H, d, J=7 Hz), 0.61 (3H, d, J=7 Hz); MS (DCI/NH₃) m/e 554.3 [M+H]⁺.

10

Example 70

(2R,4S,5S,1'S)-5-N-(propylaminocarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenylmethyl-15-hexamide.

Following the procedure of Examples 67(a), except substituting n-propyl isocyanate for t-butylisocyanate, the title compound was prepared (0.048 g, 54%). Mp 247-9°C (dec); NMR(DMSO-d₆) δ 7.75 (1H, d, J=8 Hz), 7.23-6.94 (11H, m), 6.85 (2H, s), 5.87 (1H, t, J=5 Hz), 5.65 (1H, d, J=9 Hz), 4.82 (1H, d, J=4 Hz), 4.64 (1H, t, J=8 Hz), 3.66 (1H, m), 3.38 (1H, br), 2.87 (2H, q, J=6 Hz), 2.74 (3H, m), 2.56 (1H, dd, J=7 Hz, 13 Hz), 2.39 (1H, d, J=7 Hz), 1.91 (1H, m), 1.43 (2H, m), 1.28 (2H, q, J=7 Hz), 0.77 (3H, t, J=7 Hz), 0.71 (3H, d, J=7 Hz), 0.62 (3H, d, J=7 Hz); MS (CI) m/e 520.2 [M+H]⁺.

Example 71

Preparation of (2R,4S,5S,1'S)-5-(n-propylaminothiono)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenylmethyl-hexamide.

Following the method of Example 67(a)-67(b), except using n-propyl thioisocyanate, the title compound was prepared (0.012 g, 21%). Mp 195-7°C (dec); NMR (CD₃OD) δ 7.32-6.86 (12H, m), 4.59 (1H, m), 3.64 (1H, br), 3.34 (2H, br), 2.79 (5H, m), 2.03 (1H, m), 1.73 (1H, m), 1.58 (3H, m),

- 116 -

0.92 (3H, t, J=7Hz), 0.83 (3H, d, J=7Hz), 0.68 (3H, d, J=7 Hz); MS (CI) m/e 536.2 [M+H]⁺.

Example 72

Preparation of (2R,4S,5S,1'S)-5-(isopropylaminocarbonyl)-amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenylmethyl-hexamide

Following the method of Example 67(a)-67(b), except substituting isopropyl isocyanate for t-butyl isocyanate, the title compound was prepared (0.034g, 46%). NMR(DMSO-d₆) δ 7.78 (1H, d, J=8 Hz), 7.24-6.97 (11H, m), 6.85 (2H, s), 5.74 (1H, d, J=8 Hz), 5.57 (1H, d, J=9 Hz), 4.83 (1H, d, J=4 Hz), 4.66 (1H, d, J=7 Hz), 3.62 (2H, m), 3.43 (1H, br), 2.73 (3H, m), 2.57 (1H, dd, J=7 Hz, 13.5 Hz), 2.41 (1H, d, J=7 Hz), 1.91 (1H, m), 1.45 (2H, m), 0.95 (3H, d, J=6.5 Hz), 0.93 (3H, d, J=6.5 Hz), 0.72 (3H, d, J=6.5 Hz), 0.63 (3H, d, J=6.5 Hz); MS (CI) m/e 520.2 [M+H]⁺.

Example 73

Preparation of (2R,4S,5S,1'S)-5-(aminocarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenylmethyl-hexamide

The compound of Example 67(a) (0.050 g, 0.094 mmol) was dissolved in trifluoroacetic acid (2 mL) and stirred at 50°C for 2 h. After cooling, the reaction mixture was poured into saturated sodium bicarbonate solution (50 mL) and was extracted into ethyl acetate (100 mL). The organic solution was washed with brine, dried (MgSO₄) and the solvent removed under reduced pressure. Chromatography of the residue (silica, 19:1 dichloromethane:methanol) gave the title compound as a white solid (0.036g, 80%). Mp 235°C (dec); NMR(DMSO) δ 7.82 (1H, d), 7.30-6.90 (11H, m), 6.85 (2H, d), 5.88 (1H, m), 4.86 (1H, d), 4.67 (1H, t), 3.67 (1H, m), 3.45 (1H, m), 2.75 (3H, m), 2.60 (1H, m), 2.43 (1H, m), 1.94 (1H, m), 1.49 (2H, m), 0.73 (3H, d), 0.62 (3H, d); MS (CI) m/e 478 [M+H]⁺.

- 117 -

Example 74

Preparation of (2R,4S,5S,1'S)-5-(6-quinolinylmethoxy-
carbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-
yl)methyl-6-phenylmethyl-hexanamide

Using the procedure of Example 34, except substituting
 (6-quinolinylmethyl)-(4-nitrophenyl) carbonate for (4-
 picolinyl)-(4-nitrophenyl) carbonate, the title compound was
 10 prepared.

Example 75

Preparation of (2R,4S,5S,1'S)-5-(benzoyl)amino-4-hydroxy-N-
(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenylmethyl-
hexanamide

(a) (2R,4S,5S,1'S)-5-benzoyl amino-4-t-
 (butyldimethylsiloxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-
 20 6-phenylmethyl-hexanamide.

The compound of Example 13(a) (0.11 g, 0.2 mmol),
 benzoyl chloride (0.025 g, 2.2 mmol) and

di(isopropyl)ethylamine (0.026 g, 0.2 mmol) were stirred
 together in dichloromethane (4 mL) at ambient temperature for
 25 48hr. The solvent was removed under reduced pressure and the
 residue chromatographed (silica, 1:1 ethyl acetate:hexane) to
 yield the title compound as a white solid (0.080 g, 61%).

¹H NMR (CDCl₃) 7.53 (2H, d), 7.40-7.04 (11H, m), 6.93 (2H, d),
 6.69 (2H, s), 6.59 (1H, d), 6.37 (1H, d), 4.54 (2H, m), 3.68
 30 (1H, t), 2.78 (2H, m), 2.66 (2H, m), 2.39 (1H, dd), 2.13 (1H,
 m), 1.62 (2H, t), 0.87 (9H, s), 0.53 (3H, d), 0.48 (3H, d),
 0.02 (3H, s), 0.00 (3H, s).

(b) (2R,4S,5S,1'S)-5-(benzoyl)amino-4-hydroxy-N-(1'-
 35 isopropyl-1'-imidazol-2-yl)methyl-6-phenylmethyl-hexanamide

The compound of Example 75(a) (0.080 g, 0.12 mmol) was
 dissolved in THF (1 mL) and to this was added tetrabutyl-
 ammonium fluoride, 0.16 mL, 0.16 mmol, 1M solution in THF).

- 118 -

After stirring at 40°C for 24 hr, the solvent was removed under reduced pressure and the residue was chromatographed (silica, step gradient, 1:1 ethyl acetate:hexane, 9:9:2 ethyl acetate:hexane:methanol) to give the title compound as a white solid (0.051 g, 79%). Mp 253-6°C; NMR(DMSO-d₆) δ 7.99 (1H, d), 7.91 (1H, d), 7.72 (2H, d), 7.50-7.02 (13H, m), 6.94 (2H, s), 4.83 (1H, br), 4.68 (1H, d), 4.14 (1H, m), 3.58 (1H, d), 2.82 (4H, m), 2.49 (1H, m), 1.92 (1H, m), 1.73 (1H, t), 1.40 (1H, m), 0.73 (3H, d), 0.63 (3H, d); MS(ES) m/e 539.2 [M+H]⁺.

Example 76

Preparation of (2R,4S,5S,1'S)-5-(2-furylcarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenylmethyl-hexanamide

Following the procedure of Example 75(a), except using furoyl chloride in place of benzoyl chloride, the title compound was prepared as a white solid (0.019 g, 18%). Mp 212-3°C (dec); NMR(CDCl₃/CD₃OD) δ 7.46 (1H, s), 7.30-6.88 (12H, m), 6.85 (2H, s), 6.49 (1H, m), 4.48 (1H, d), 4.20 (1H, m), 3.67 (1H, m), 2.96 (4H, m), 2.77 (2H, m), 2.58 (1H, d), 2.07 (1H, m), 1.71 (2H, m), 0.74 (3H, d), 0.65 (3H, d); MS(ES) m/e 528.32 [M+H]⁺.

Example 77

Preparation of (2R,4S,5S,1'S)-5-(4-methoxybenzoyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenylmethyl-hexanamide

Following the procedure of Example 75(a), except using 4-methoxybenzoyl chloride in place of benzoyl chloride, the title compound was prepared as a white solid (32%). Mp 235-7°C (dec); NMR(CDCl₃/CD₃OD) δ 7.64 (2H, d), 7.22-6.87 (14H, m), 6.80 (2H, m), 4.52 (1H, d), 4.16 (1H, m), 3.81 (3H, s), 3.62 (1H, d), 2.92 (2H, d), 2.72 (2H, m), 2.53 (1H, dd), 1.98 (1H, m), 1.73 (1H, m), 1.63 (1H, m), 0.71 (3H, d), 0.62 (3H, d); MS(ES) m/e 569.4 [M+H]⁺.

Example 78Preparation of (2R,4S,5S,1'S)-5-benzylcarbonylamino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenylmethyl-hexanamide.

a) (2R,4S,5S,1'S)-5-benzylcarbonylamino-4-t-butyldimethylsiloxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenylmethyl-hexanamide.

Following the procedure of Example 75(a), except using phenylacetyl chloride in place of benzoyl chloride and triethylamine in place of di(isopropyl)ethylamine, the title compound was prepared as a white solid (20%). NMR(CDCl₃) δ 7.40-6.75 (19H, m), 5.40 (1H, d), 4.73 (1H, t), 4.41 (1H, q), 3.68 (1H, m), 3.48 (2H, s), 2.96 (1H, dd), 2.69 (1H, m), 2.49 (4H, m), 1.61 (2H, m), 0.92 (6H, t), 0.77 (9H, s), 0.04 (3H, s), 0.00 (3H, s).

b) (2R,4S,5S,1'S)-5-benzylcarbonylamino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenylmethyl-hexanamide

The product of Example 78 (a) (0.018 g, 0.03 mmol) was dissolved in methanol (5 mL) and 2N hydrochloric acid (0.027 mL, 0.06 mmol) was added. After stirring at ambient temperature for 18 h the solvent was removed under reduced pressure and the residue was chromatographed (silica, gradient, dichloromethane/methanol) to yield the title compound (0.011 g, 66%). Mp 240-2°C; NMR(CDCl₃/CD₃OD) δ 7.38-7.06 (18H, m), 6.98 (2H, s), 4.72 (1H, d), 4.14 (1H, m), 3.67 (1H, m), 3.54 (2H, s), 2.99 (4H, m), 2.67 (1H, m), 2.14 (1H, m), 1.87 (1H, m), 1.63 (1H, m), 0.94 (3H, d), 0.79 (3H, d), 0.0 (d); MS (ES) m/e 553.2 [M+H]⁺.

Example 79

Preparation of (2R,4S,5S,1'S)-5-(4-hydroxybenzoyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenylmethyl-hexanamide

- 120 -

a) (2R,4S,5S,1'S)-5-(4-acetoxyphenyl)-4-t-butyl dimethylsiloxyl-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenylmethyl-hexanamide.

5 The compound of Example 13(a) (0.11 g, 0.2 mmol) was dissolved in dichloromethane (2 mL), and BOP reagent (0.089 g, 0.2 mmol), triethylamine (0.028 mL, 0.2 mmol) and 4-acetoxybenzoic acid (0.043 g, 0.24 mmol) were added. After stirring at ambient temperature overnight the solvent was removed under reduced pressure. The residue was chromatographed (silica, 49:1 dichloromethane:methanol) to give the title compound as a white solid (0.11 g, 78%).

10 NMR(CDCl₃) δ 7.53 (2H, d), 7.28-6.97 (13H, m), 6.83 (1H, d), 6.78 (2H, s), 6.44 (1H, d), 4.54 (2H, m), 3.72 (1H, dd), 2.79 (4H, m), 2.49 (1H, dd), 2.24 (3H, s), 2.20 (1H, m), 1.70 (2H, m), 0.91 (9H, s), 0.66 (3H, d), 0.57 (3H, d), 0.07 (3H, s), 0.02 (3H, s).

b) (2R,4S,5S,1'S)-5-(4-hydroxybenzoyl)amino-4-t-butyl dimethylsiloxyl-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenylmethyl-hexanamide

20 The product from reaction 79(a) (0.11 g, 0.15 mmol) was dissolved in methanol (5 mL) and powdered potassium carbonate (0.12 g, 0.9 mmol) was added. After stirring the suspension vigorously for 2 h, the mixture was filtered and the solvent removed from the filtrate at reduced pressure.

25 Chromatography of the residue (silica, 19:19:2 ethyl acetate:hexane:methanol) gave the title compound as a white solid (0.066 g, 66%). NMR(CDCl₃) δ 7.35 (2H, d), 7.24-6.98 (12H, m), 6.67 (4H, m), 6.32 (1H, d), 4.63 (2H, m), 3.76 (1H, dd), 2.78 (4H, m), 2.44 (1H, d), 2.12 (1H, m), 1.64 (2H, m), 0.88 (9H, s), 0.44 (3H, d), 0.32 (3H, d), 0.05 (3H, s), 0.01 (3H, s).

35 c) (2R,4S,5S,1'S)-5-(4-hydroxybenzoyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenylmethyl-hexanamide

Following the procedure of Example 75(b), except using the compound of Example 79(b) in place of the compound of

- 121 -

Example 75(a), the title compound was prepared as a white solid (57%). Mp 267-8°C (dec); NMR(CDCl₃/CD₃OD) δ 7.57 (2H, d), 7.33-6.75 (17H, m), 4.48 (1H, d), 4.14 (1H, m), 3.58 (1H, d), 2.90 (2H, m), 2.82 (1H, m), 2.73 (1H, m), 2.53 (1H, dd), 2.04 (1H, m), 1.65 (2H, m), 0.73 (3H, d), 0.58 (3H, d); MS (ES) m/e 555.2 [M+H]⁺.

Example 80

10 Preparation of (2R,4S,5S,1'S)-5-(cinnamoyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenylmethyl-hexanamide

Following the procedure of Example 75(a), except using cinnamoyl chloride in place of benzoyl chloride, the title compound was prepared as a white solid (25%). Mp 273°C; NMR(CDCl₃/CD₃OD) δ 7.55-6.91 (19H, m), 6.86 (2H, s), 6.53 (1H, d), 4.37 (1H, d), 4.15 (1H, dt), 3.62 (1H, d), 2.91 (2H, m), 2.78 (2H, m), 2.59 (1H, dd), 2.04 (1H, m), 1.76 (1H, m), 1.65 (1H, m), 0.79 (3H, d), 0.69 (3H, d); MS (ES) m/e 565.2 [M+H]⁺.

Example 81

25 Preparation of (2R,4S,5S,1'S)-5-(2-hydroxybenzoyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenylmethyl-hexanamide

Following the procedure of Example 79(a), except using (2-acetoxybenzoic acid) in place of 4-acetoxybenzoic acid, the title compound was prepared (50%). Mp 197°C; NMR(CD₃OD) δ 7.77 (1H, d), 7.42-6.78 (17H, m), 4.62 (1H, d), 4.32 (1H, dt), 3.71 (1H, m), 2.94 (2H, m), 2.78 (2H, m), 2.57 (1H, m), 2.03 (1H, m), 1.84 (1H, m), 1.67 (1H, m), 0.82 (3H, d), 0.68 (3H, d); MS (ES) m/e 555.2 [M+H]⁺.

Example 82

Preparation of (2R,4S,5S,1'S)-5-(imidazol-4-yl-acetyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenylmethyl-hexanamide

Following the procedure of Example 79(a)-79(c), except using (imidazol-4-yl)acetic acid in place of 4-acetoxy benzoic acid, the title compound was prepared.

Example 83

Preparation of (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-[1'-isopropyl-1'-(4-carbomethoxyethylimidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide

a) (1S) 1-carbobenzyloxyamino-1-isopropyl-1-[(4-(E-carbomethoxyethylene)imidazol-2-yl)]methane

The compound of Example 27(b) (100 mg, 0.33 mmol), lithium chloride (28 mg, 0.66 mmol) and

trimethylphosphonoacetate (61 mg, 0.33 mmol) were dissolved in anhydrous acetonitrile (2 mL). 1,8-Diazabicyclo[5.4.0]-undec-7-ene (55 mg, 0.36 mmol) was added and the reaction mixture was stirred at room temperature overnight. The solvent was removed under reduced pressure and the residue

was purified by flash chromatography (silica, 2% methanol/dichloromethane to afford the title compound (72 mg, 61%). NMR(CDCl₃) δ 7.60-7.10 (6H, m), 6.50 (1H, br s), 6.10 (1H, br s), 5.15-4.95 (2H, m), 4.50 (1H, br m), 3.75 (3H, s), 2.30 (1H, br m), 1.10-0.80 (6H, m); MS m/e 358.2 [M+H]⁺.

b) (1S)-1-amino-1-isopropyl-1-(4-carbomethoxyethylimidazol-2-yl)methane

Following the procedure of Example 1(b), except substituting the compound of Example 82(a) for the compound of Example 1(a), the title compound was prepared. NMR(CDCl₃) δ 6.65 (1H, s), 4.40 (2H, br s), 3.82 (1H, d, J=3 Hz), 3.65 (3H, s), 2.90-2.55 (4H, m), 2.05 (1H, m), 0.90 (6H, d, J=3Hz).

c) (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-t-butyl-di-methylsiloxy-N-[1'-isopropyl-1'-(4-carbomethoxyethylimidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide

5 Following the procedure of Example 1(c) except using the compound of Example 82(b), the title compound was prepared. NMR(CDCl₃) δ 7.35-6.90 (12H, m), 6.55 (1H, s), 4.75 (1H, d, J=4 Hz), 4.45 (1H, m), 3.95 (1H, m), 3.70 (3H, s), 2.90-2.40 (9H, m), 1.90-1.60 (2H, m), 1.38 (9H, s), 10 0.90-0.70 (15H, m), 0.10 (6H, d, J=2 Hz).

d) (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-[1'-isopropyl-1'-(4-carbomethoxyethylimidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide.

15 Following the procedure of Example of 9(d) except using the compound of Example 83(c), the title compound was prepared. NMR(CDCl₃) δ 7.30-6.90 (10H, m), 6.55 (1H, s), 5.00 (1H, d, J=4 Hz), 4.45 (1H, m), 3.70 (3H, s), 2.95-2.50 (9H, m), 2.25 (1H, m), 1.80-1.60 (2H, m), 0.85 (9H, s), 0.70 20 (6H, d, J=3 Hz); MS m/e 621.4 [M+H]⁺.

Example 84

Preparation of (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-[1'-isopropyl-1'-(4-carboxamidoimidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide

25 (a) (1S)-1-carbobenzyloxyamino-1-isopropyl-1-[(4-(hydrazinocarbonyl)imidazol-2-yl)]methane
30 Anhydrous hydrazine (47 μ L, 1.5 mmol) was added to a solution of the compound of Example 26(b) (100 mg, 0.30 mmol) in anhydrous methanol. The resulting mixture was stirred overnight at room temperature and concentrated under reduced pressure. The residue was partitioned between ethyl acetate and 10% aqueous Na₂CO₃ and the organic extract was dried over Na₂CO₃ and vaporated under reduced pressure. The residue was purified by flash chromatography (silica, 4% methanol/dichloromethane) to afford the title compound (52

- 124 -

mg, 52%). NMR(CD₃OD) δ 7.50 (1H, s), 7.30-7.20 (5H, m), 5.00-4.90 (2H, m), 4.45 (1H, d, J=6 Hz), 2.10 (1H, br m), 0.95-0.75 (6H, m); MS m/e 332.2 [M+H]⁺

5 b) (1S)-1-carbobenzyloxyamino-1-isopropyl-1-[(4-azidocarbonyl)imidazol-2-yl]methane

The compound of Example 83(a) was dissolved in 2N HCl (1 mL) and glacial acetic acid (0.2 mL) and cooled in an ice bath. A solution of sodium nitrite (11 mg, 0.16 mmol) in H₂O (200 μ L) was added dropwise. The reaction mixture was stirred for 0.5 h, neutralized with cold concentrated ammonium hydroxide and extracted with ethyl acetate. The organic extract was dried over Na₂CO₃ and the solvent removed in vacuo to yield the title compound (54mg, 100%).

15 NMR(CDCl₃) δ 7.75 (1H, s), 7.35-7.20 (5H, m), 5.20-5.00 (2H, m), 4.62 (1H, br m), 2.60 (1H br m), 1.10-0.80 (6H, m); IR 2123cm⁻¹ (CON₃).

20 c) (1S)-1-carbobenzyloxyamino-1-isopropyl-1-(4-carboxamidoimidazol-2-yl)methane

The compound of Example 83(b) was dissolved in 2 mL of ethyl acetate and stirred with of concentrated ammonium hydroxide (1 mL) at 0°C for 0.5 h, then at room temperature overnight. The reaction mixture was diluted with H₂O, extracted with ethyl acetate, and dried over Na₂CO₃. The solvent was removed in vacuo and the residue was purified by flash chromatography (silica, 4% methanol/ dichloromethane) to afford the title compound (50mg, 100%). NMR(CDCl₃) δ 7.45 (1H, s), 7.25-7.10 (5H, m), 5.00-4.85 (2H, m), 4.35 (1H, d, J=3 Hz), 2.00 (1H, br m), 0.90-0.70 (6H, m); MS m/e 317.2 [M+H]⁺.

35 d) (1S)-1-amino-1-isopropyl-1-(4-carboxamidoimidazol-2-yl)methane.

Following the procedure of Example 1(b), except substituting the compound of Example 83(c) for the compound of Example 1(a), the title comp und was prepared. NMR(CDCl₃)

Example 85

δ 7.45 (1H, s), 3.47 (1H, d, J=3 Hz), 1.80 (1H, br m), 0.75-0.60 (6H, m).

e) (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-t-butyl-
5 butyldimethylsiloxy-N-[1'-isopropyl-1'-(4-carboxamidoimidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide

Following the procedure of Example 1(c), except using the compound of Example 83(d), the title compound was prepared. NMR(CDCl₃) δ 7.50 (1H, s), 7.45-6.90 (11H, m), 6.25 (1H, d, J=4 Hz), 4.50 (1H, d, J=6Hz), 4.10 (1H, br m), 3.60 (1H, m), 2.90-2.40 (5H, m), 1.90 (1H, br m), 1.70-1.50 (2H, br m), 1.35 (9H, s), 0.90 (9H, s), 0.70-0.60 (6H, m), 0.10 (6H, m).

f) (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-[1'-isopropyl-1'-(4-carboxamidoimidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide

Following the procedure of Example 9(d) except using the compound of Example 83(e) the title compound was prepared. NMR(CD₃OD) δ 7.45 (1H, s), 7.25-6.85 (10H, m), 4.50 (1H, d, J=6 Hz), 4.10 (1H, m), 3.60 (1H, m), 2.85-2.50 (5H, m), 2.00 (1H, br m), 1.80-1.50 (2H, m), 1.30 (9H, s), 0.80-0.65 (6H, m); MS m/e 578.2 [M+H]⁺.

Preparation of (2R,4S,5S,1'S)-2-phenylmethyl-4-hydroxy-5-(5-(1-oxopropyl)-2-thiazolyl)amino)-6-phenyl-N-(1'-isopropyl-1'-(imidazo-2-yl))methyl-hexanamide

a) (2R,4S,5S,1'S)-2-phenylmethyl-4-t-butyl-5-thioureido-6-phenyl-N-(1'-isopropyl-1'-(imidazo-2-yl))methyl-hexanamide

A solution of benzoyl isothiocyanate (prepared from ammonium thiocyanate (147 mg, 1.93 mmol) and benzoyl chloride (257 mg, 1.84 mmol) in of acetone (10 mL) according to the

procedure of *J. Amer. Chem. Soc.*, 56, 1408 (1934)) was treated with a solution of (2R,4S,5S,1'S)-2-phenylmethyl-4-t-butylldimethylsiloxy-5-amino-6-phenyl-N-(1'-isopropyl-1'-(imidazo-2-yl))methyl-hexanamide (1.0 g, 1.83 mmol) in acetone. After 20 min at 23°C, the solvent was evaporated, and the residue was dissolved in diethyl ether. The ether extract was washed with water, dried, and the solvent was evaporated. This residue was dissolved in MeOH (25 mL), treated with 2.5N NaOH (0.1 mL) and heated to 50°C for 30 min. The solvent was evaporated, and the residue was dissolved in EtOAc. The organic solution was washed with water, dried, and the solvent evaporated. The residue was chromatographed (silica, 5% MeOH/CHCl₃) to yield the title compound (520 mg, 47%). NMR(DMSO) δ 7.80 (1H, d), 7.35 (1H, d), 6.70-7.20 (15H, m), 4.69 (1H, t), 4.54 (1H, m), 3.78 (1H, m), 2.72-2.86 (3H, m), 2.54 (1H, dd), 2.42 (1H, dd), 2.04 (1H, m), 1.82 (1H, m), 1.30 (1H, m), 0.92 (9H, s), 0.86 (3H, d), 0.74 (3H, d), 0.15 (6H, d).

b) dimethylformamidino derivative of (2R,4S,5S,1'S)-2-phenylmethyl-4-dimethyl-t-butyl silyloxy-5-thioureido-6-phenyl-N-(1'-isopropyl-1'-(imidazo-2-yl))methyl-hexanamide
A solution of the compound of Example 85(a) (122 mg, 0.2 mmol) and dimethylformamide dimethylacetal (26 mg, 0.22 mmol) in CHCl₃ (2 mL) was stirred at 23°C for 16 h. The solvent and excess reactant was removed under high vacuum, and the residue was chromatographed (Florisil®, 2% MeOH/CHCl₃) to yield the title compound (100 mg, 76%). NMR(CDCl₃) δ 8.82 (1H, s), 7.05-7.40 (12H, m), 6.76 (1H, br s), 6.60 (1H, d), 5.32 (1H, m), 4.66 (1H, dd), 3.88 (1H, dd), 3.14 (3H, s), 3.05 (3H, s), 2.70-3.04 (4H, m), 2.40 (2H, m), 1.68 (2H, m), 1.00 (9H, s), 0.80 (6H, dd), 0.14 (6H, d).

c). (2R,4S, 5S, 1'S)-2-phenylmethyl-4-dimethyl-t-butyl silyloxy-5-(5-(1-oxopropyl)-2-thiazolyl)amino)-6-phenyl-N-(1'-isopropyl-1'-(imidazo-2-yl))methyl-hexanamide

A solution of the compound of Example 85(b) (100 mg, 0.15 mmol), 1-bromo-2-butanone (25 mg, 0.165 mmol), and

triethylamine (33 mg, 0.165 mmol) in acetonitrile (10 mL) was heated at 80°C for 3.5 h. The solvent was evaporated, and the residue shaken with a mixture of diethyl ether and aqueous NaHCO₃. The ether was separated, washed with water, dried, and the solvent was evaporated. The residue was recrystallized from a mixture of CHCl₃ and hexane to yield the title compound (59 mg, 57%). NMR(CDCl₃) δ 7.75 (1H, s), 7.02-7.38 (10H, m), 6.88 (2H, m), 6.80 (1H, br s), 6.70 (1H, d), 6.60 (1H, d), 4.62 (1H, t), 3.96 (1H, m), 3.78 (1H, t), 2.82 (3H, m), 2.72 (2H, q), 2.54 (2H, m), 2.20 (1H, m), 2.04 (1H, m), 1.66 (1H, m), 1.15 (3H, t), 0.96 (9H, s), 0.72 (6H, t), 0.10 (6H, d).

d). (2R,4S, 5S, 1'S)-2-phenylmethyl-4-hydroxy-5-(5-(1-oxopropyl)-2-thiazolyl)amino)-6-phenyl-N-(1'-isopropyl-1'-(imidazo-2-yl))methyl-hexanamide

A solution of the compound of Example 85(c) (50 mg, 0.07 mmol) in (2 mL) of THF was treated with) of tetrabutylammonium fluoride (0.2 mL, 1N solution in THF) 58°C for 1 h. The solvents were evaporated, and the residue dissolved in ether. The ether was washed with water, dried, and the solvent evaporated. The residue was chromatographed (neutral alumina, Activity V, impurities removed with 2% MeOH/EtOAc, product eluted with 5% MeOH/CHCl₃) to yield the title compound (22 mg, 55%). NMR(DMSO) δ 7.75 (1H, s), 7.66 (1H, d), 6.80-7.30 (13H, m), 4.93 (1H, br s), 4.78 (1H, t), 3.78 (1H, m), 3.68 (1H, dd), 3.00 (1H, dd), 2.92 (1H, dd), 2.86 (1H, m), 2.80-2.90 (1H, br), 2.76 (2H, q), 2.56 (2H, m), 2.12 (1H, m), 1.74 (1H, m), 1.69 (1H, m), 1.20 (3H, t), 0.80 (3H, d), 0.73 (3H, d).

Example 86

Preparation of (2R,4S,5S,1'S)-2-phenylmethyl-4-hydroxy-5-(5-(1-oxopropyl)-2-thiazolyl)amino)-6-phenyl-N-(1'-isopropyl-1'-(imidazo-2-yl))methyl-hexanamide

a) (2R,4S,5S,1'S)-2-phenylmethyl-4-dimethyl-t-butyl silyloxy-5-(2-thiazolylamino)-6-phenyl-N-(1'-isopropyl-1'-(imidazo-2-yl))methyl-hexanamide.

The compound of Example 85(a) (50 mg, 0.08 mmol) in 5 CHCl_3 (2 mL) was treated with chloroacetaldehyde (50 mg, 0.64 mmol). After 20 min the solvent and excess reagent were evaporated. The residue was dissolved in EtOAc, washed with aqueous NaHCO_3 , dried and the solvent evaporated. The residue was chromatographed (Florisil®, 60% EtOAc/hexane) to 10 yield the title compound (42 mg, 83%). NMR(CDCl_3) δ 7.12-7.30 (10H, m), 7.02 (1H, d), 6.92 (2H, m), 6.82 (1H, br), 6.62 (1H, br), 6.38 (1H, d), 5.86 (1H, br), 4.58 (1H, t), 4.00 (1H, m), 3.86 (1H, m), 2.85 (3H, m), 2.52 (2H, m), 2.26 (1H, m), 2.16 (1H, m), 1.68 (1H, m), 0.98 (9H, s), 0.70 (6H, 15 t), 0.12 (6H, d).

b) (2R,4S,5S,1'S)-2-phenylmethyl-4-hydroxy-5-(5-(1-oxopropyl)-2-thiazolyl)amino)-6-phenyl-N-(1'-isopropyl-1'-(imidazo-2-yl))methyl-hexanamide.

20 Following the procedure of Example 85(d), except substituting the compound of Example 86(a) for the compound of Example 85(c), the title compound was prepared. NMR($\text{CDCl}_3/\text{DMSO}$) δ 6.80-7.42 (14H, m), 6.40 (2H, m), 5.18 (1H, br), 4.74 (1H, t), 3.70 (1H, m), 3.62 (1H, m), 3.00 (2H, m), 25 2.88 (2H, m), 2.58 (1H, m), 2.18 (1H, m), 1.80 (2H, m), 1.72 (6H, dd).

Example 87

30 Preparation of (2R,4S,5S,1'S)-2-phenylmethyl-4-hydroxy-5-(5-propyl-2-thiazolyl)amino)-6-phenyl-N-(1'-isopropyl-1'-(imidazo-2-yl))methyl-hexanamide

a) (2R,4S,5S,1'S)-2-phenylmethyl-4-t-butyl dimethylsilyloxy-5-(5-propyl-2-thiazolyl)amino)-6-phenyl-N-(1'-isopropyl-1'-(imidazo-2-yl))methyl-hexanamide.

A solution of the compound of Example 85(a) (120 mg, 0.2 mmol) in CHCl_3 (5 mL) was treated with 2-bromovaleraldehyde

- 129 -

100 mg, 0.6 mmol) and warmed to 60°C for 30 min and 80°C for 5 min. The solvent and excess reagent were removed under reduced pressure. The residue was dissolved in EtOAc, washed with aqueous K₂CO₃, dried, and the solvent evaporated. The residue was chromatographed (silica, 3% MeOH/CHCl₃) to yield the title compound (55 mg, 41%). NMR(CDCl₃) δ 7.10-7.30 (10H, m), 6.88 (2H, m), 6.72 (1H, br), 6.68 (1H, s), 6.60 (1H, br), 5.60 (1H, br), 4.62 (1H, t), 3.94 (1H, m), 3.78 (1H, t), 2.82 (3H, m), 2.50 (4H, m), 2.26 (1H, m), 2.04 (1H, m), 1.66 (1H, m), 1.55 (2H, sextet), 0.94 (9H, s), 0.92 (3H, t), 0.70 (6H, dd), 0.08 (6H, d).

b). (2R,4S, 5S, 1'S)-2-phenylmethyl-4-hydroxy-5-(5-propyl-2-thiazolyl)amino)-6-phenyl-N-(1'-isopropyl-1'-(imidazo-2-yl)methyl-hexanamide.

Following the procedure of Example 85(d), except substituting the compound of Example 87(a) for the compound of Example 85(c), the title compound was prepared. NMR(CDCl₃) δ 7.50 (1H, br), 6.90-7.24 (10H, m), 6.78 (2H, s), 6.60 (1H, s), 6.18 (1H, br), 5.76 (1H, br), 4.60 (1H, t), 3.68 (1H, m), 3.52 (1H, m), 3.05 (1H, dd), 2.95 (2H, m), 2.82 (1H, dd), 2.62 (1H, m), 2.58 (2H, t), 2.32 (1H, m), 1.86 (2H, m), 1.60 (2H, sextet), 0.96 (6H, t), 0.75 (6H, dd).

25

Example 88Preparation of (2R,4S,5S,1'S)-5-(nicotinyl)amino-4-hydroxy-N-(1'-isopropyl-1'-(imidazol-2-yl)methyl-6-phenylmethyl-hexanamide

Following the procedure of Example 75(a), except using nicotinoyl chloride in place of benzoyl chloride, the title compound was prepared as a white solid (43%). Mp 233-4°C (dec); NMR(CDCl₃/CD₃OD) δ 8.81 (1H, d), 8.59 (1H, dd), 7.99 (1H, m), 7.35-6.86 (14H, m), 6.79 (2H, s), 4.44 (1H, d), 4.19 (1H, dt), 3.59 (1H, m), 2.90 (2H, d), 2.68 (2H, m), 2.52 (2H, m), 1.96 (1H, m), 1.71 (1H, m), 1.58 (1H, m), 0.70 (3H, d), 0.58 (3H, d); MS(ES) m/e 540.2 [M+H]⁺.

5 the scope of the following claims:

[illegible]

(b) (7)(C), (b) (7)(D), (b) (7)(F), (b) (7)(G), (b) (7)(H), (b) (7)(I), (b) (7)(J), (b) (7)(K), (b) (7)(L), (b) (7)(M), (b) (7)(N), (b) (7)(O), (b) (7)(P), (b) (7)(Q), (b) (7)(R), (b) (7)(S), (b) (7)(T), (b) (7)(U), (b) (7)(V), (b) (7)(W), (b) (7)(X), (b) (7)(Y), (b) (7)(Z)


Following the procedure of the following

[illegible]

1. The first part of the document is a list of names and dates, which appears to be a roster or a list of personnel. The names are written in a cursive script, and the dates are written in a more formal, printed style. The list includes names such as "John A. Smith", "John B. Smith", "John C. Smith", "John D. Smith", "John E. Smith", "John F. Smith", "John G. Smith", "John H. Smith", "John I. Smith", "John J. Smith", "John K. Smith", "John L. Smith", "John M. Smith", "John N. Smith", "John O. Smith", "John P. Smith", "John Q. Smith", "John R. Smith", "John S. Smith", "John T. Smith", "John U. Smith", "John V. Smith", "John W. Smith", "John X. Smith", "John Y. Smith", "John Z. Smith", "John A. Smith", "John B. Smith", "John C. Smith", "John D. Smith", "John E. Smith", "John F. Smith", "John G. Smith", "John H. Smith", "John I. Smith", "John J. Smith", "John K. Smith", "John L. Smith", "John M. Smith", "John N. Smith", "John O. Smith", "John P. Smith", "John Q. Smith", "John R. Smith", "John S. Smith", "John T. Smith", "John U. Smith", "John V. Smith", "John W. Smith", "John X. Smith", "John Y. Smith", "John Z. Smith".

What is claimed is:

1. A compound of the formula (I):



(I)

wherein:


R¹ and R³ are each independently Q, Q-C₁₋₆alkyl, Q-C₂₋₆alkenyl, Q-C₂₋₆alkynyl or C₁₋₆alkyl substituted by one to five fluorine atoms, each optionally substituted by R²³;

Q is H, C₃₋₆cycloalkyl, C₅₋₆cycloalkenyl, Ar or Het

R² is H or OH;

R⁴ is R⁶-NR¹¹ or CONR¹¹CHR⁶R⁷;

R⁵ is R⁶-NR¹¹ or R¹⁰-NR¹¹;



X is NR¹¹, O or S;

R⁷ is Q, Q-C₁₋₆alkyl or Q-C₂₋₆alkenyl;

R⁸ and R⁹ are each independently H, OH, halo, NO₂, COR¹², CF₃, Ar, C₁₋₆alkyl-R¹⁵, or R¹⁷(R¹⁸R¹⁹C)_m, or together form a fused C₂₋₄alkylene, aryl or heteroaryl moiety;

R¹⁰ is A-(B)_n;

R¹¹ is H or C₁₋₄alkyl;

R¹² is R⁷, OR⁷, NR⁷R¹¹ or an amino acid or amino alcohol;

B is an amino acid;

A is H, Ar, Het, R¹⁷(R¹⁸R¹⁹C)_m, Ar-W, Het-W or

R¹⁷(R¹⁸R¹⁹C)_m-W, or phthaloyl each optionally substituted by one to three groups chosen from R¹⁵ or C₁₋₆alkyl-R¹⁵;

W is C=O, OC(=O), NR¹¹C(=O), SC(=O), NR¹¹C(=S), SO₂,

NR¹¹SO₂ or P(=O)(OR²²);

R¹⁵ is H, nitro, C₁₋₆alkoxy, C₁₋₆alkylthio, O(C=O)R¹⁶, C=OR²², CO₂R²², CON(R¹⁶)₂, N(R²²)₂, NHC(=N)NH-A, I, Br, Cl, F, OR¹⁰, or OH, provided that when R¹⁵ is a substituent of the carbon adjacent to W, R¹⁵ is not halogen or OH when W is OC(=O) or NHCO;

R¹⁶ is H or C₁₋₆alkyl;

- 132 -

R¹⁷, R¹⁸ and R¹⁹ are independently: i) H, R¹⁵ or C₁₋₄alkyl, C₂₋₆alkenyl, phenyl, naphthyl, C₃₋₆cycloalkyl or Het, each optionally substituted by one to three R¹⁵ or R¹⁵-C₁₋₆alkyl groups, or ii) R¹⁷ is as above and (R¹⁸R¹⁹C) are joined together to form a phenyl, naphthyl, C₃₋₆cycloalkyl or Het ring, or iii) R¹⁷ is as above and R¹⁸ and R¹⁹ together are =O;

R²² is H, C₁₋₆alkyl, phenyl or phenyl-C₁₋₄alkyl;

R²³ is -X'-(CH₂)_qNR²⁴R²⁵, X''[((CH₂)_rO)_s]R²⁶,
 10 CH₂X''[((CH₂)_rO)_s]R²⁶, or benzofuryl, indolyl, azacycloalkyl, azabicyclo C₇₋₁₁cycloalkyl or benzopiperidinyl, optionally substituted with C₁₋₄alkyl;

q is 2-5;

s is 1-6 and r is 1-3 within each repeating unit s;

15 X' is CH₂, O, S or NH;

X'' is CH₂, NR', O, S, SO or SO₂;

R²⁴ and R²⁵ are i) C₁₋₆alkyl, optionally substituted by OH, C₁₋₃alkoxy, or N(R')₂, ii) the same or different and joined together to form a 5-7 member heterocycle containing
 20 up to two additional heteroatoms selected from NR, O, S, SO, SO₂, said heterocycle optionally substituted with C₁₋₄alkyl, iii) aromatic heterocycle, optionally substituted with C₁₋₄alkyl or N(R')₂;

R' is H or C₁₋₄alkyl;

25 R²⁶ is H, C₁₋₄alkyl, C(=O)R²⁷, C(=O)U[(CH₂)_mO]_nNR', P(=O)(OM)₂, CO₂R²⁷, C(=O)NR²⁷R²⁸, where M is a mono or divalent metal ion, and U is NR' or O;

R²⁷ is C₁₋₆alkyl or Ar, optionally substituted with one or more hydroxy, carboxy, halo, C₁₋₃alkoxy, CONR'₂, NR'₂,
 30 CO₂R', SO₂NR'₂, CH₂NR₂, NR'COR', NR'SO₂R', X''[((CH₂)_rO)_s]R' or CH₂X''[((CH₂)_rO)_s]R';

R²⁸ is H, C₁₋₆alkyl or together with R²⁷ forms a 5-7 membered heterocycle or a 6 membered heterocycle containing a heteroatom selected from N, O and S;

35 m is 1-4; and

n is 0 or 1;

or a pharmaceutically acceptable salt thereof.

2. A compound according to claim 1 wherein:

R^1 and R^3 are C_1 -6alkyl, Ar- C_1 -6alkyl, Ar- C_2 -6alkenyl, Ar- C_2 -6alkynyl, or C_1 -6alkyl optionally substituted by one to five fluorine atoms;

5 X is N- R^{11} ;

R^4 is CONR¹¹CHR⁶R⁷;

R^5 is R^{10} -NR¹¹;

R^7 is H, C_1 -6alkyl, C_3 -6cycloalkyl, phenyl or benzyl;

R^8 is H, C_1 -6alkyl, COR¹², NO₂ or Br;

R^9 is H, NO₂, Br, COR¹², CF₃, Ar, C_1 -6alkyl or

C_1 -6alkyl- R^{15} , wherein R^{12} is H, C_1 -6alkyl, Ar, OC $_1$ -6alkyl,

NH₂, and R^{15} is OH;

A is H, Het, $R^{17}(R^{18}R^{19}C)_m$ -W or Het-W;

B is absent or Val;

15 R^{17} , R^{18} and R^{19} are H, or C_1 -4alkyl, Het or Ar, each

optionally substituted by one or two R^{15} or $R^{15}C_1$ -6alkyl

groups, or $(R^{18}R^{19}C)$ are joined together to form a phenyl,

C_3 -6cycloalkyl or Het ring; and

W is C=O, OC(=O), NHC(=O), NHC(=S) or SC(=O).

20

3. A compound according to claim 1 wherein R^4 is

CONR¹¹CHR⁶R⁷ and X is N-H.

4. A compound according to claim 3 wherein R^8 is H and R^9

25 is H or COR¹².

5. A compound according to claim 4 wherein R^7 is C_1 -6alkyl.

6. A compound according to claim 3 wherein R^1 is benzyl and

30 R^3 is benzyl, 4-hydroxy-benzyl or phenylpropenyl.

7. A compound according to claim 3 wherein A is

$R^{17}(R^{18}R^{19}C)_m$ -W, and R^{17} , R^{18} and R^{19} are H, or C_1 -4alkyl, Het

or Ar.

35

8. A compound according to claim 3 wherein B is absent and

A is C_1 -6alkylOC(=O).

9. A compound according to claim 3 wherein W is C=O.
10. A compound according to claim 1 wherein the compound is:
- 5 (2R,4S,5S,1'S)-2-phenylmethyl-4-hydroxy-5-(t-butoxycarbonyl)-amino-6-phenyl-N-(1'-isopropyl-1'-(imidazo-2-yl))methyl-hexanamide hydrochloride;
- (2R,4S,5S,1'S)-2-phenylmethyl-4-hydroxy-5-(t-butoxycarbonyl)-amino-6-phenyl-N-[1'-isopropyl-1'-(4-aminocarbonyl-thiazo-2-yl)]methyl-hexanamide;
- 10 (2R,4S,5S,1'S)-2-phenylmethyl-4-hydroxy-5-(t-butoxycarbonyl)-amino-6-phenyl-N-[1'-isopropyl-1'-(thiazo-2-yl)]methyl-hexanamide;
- (2R,4S,5S,1'S)-2-phenylmethyl-4-hydroxy-5-(t-butoxycarbonyl)-amino-6-phenyl-N-(1'-imidazo-2-yl)methyl-hexanamide
- 15 hydrochloride;
- (2R,4S,5S,1'S)-2-phenylmethyl-4-hydroxy-5-(t-butoxycarbonyl)-amino-6-phenyl-N-[1'-methyl-1'-(imidazo-2-yl)]methyl-hexanamide hydrochloride;
- (2R,4S,5S,1'S)-2-phenylmethyl-4-hydroxy-5-(t-butoxycarbonyl)-amino-6-phenyl-N-[1'-benzyl-1'-(imidazo-2-yl)]methyl-hexanamide hydrochloride;
- 20 (2R,4S,5S,1'S)-5-(carbobenzyloxy)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide;
- 25 (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-[1'-isopropyl-1'-(4,5-dimethyl)imidazol-2-yl]methyl-6-phenyl-2-phenylmethyl-hexanamide;
- (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-[1'-isopropyl-1'-(N'-methyl)imidazol-2-yl]methyl-6-phenyl-2-phenylmethyl-hexanamide;
- 30 (2R,4S,5S,1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-(3-phenylpropargyl)hexanamide;
- (2R,4S,5S,1'S)-5-(isopropoxycarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide;
- 35

- (2R,4S,5S,1'S)-5-(benzyloxyethoxycarbonyl) amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide;
- (2R,4S,5S,1'S)-5-(methoxycarbonyl) amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide;
- (2R,4S,5S,1'S)-5-(ethoxycarbonyl) amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide;
- 10 (2R,4S,5S,1'S)-5-(t-butoxycarbonyl) amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-(3-phenyl-2-propenyl)hexanamide;
- (2R,4S,5S,1'S)-5-(t-butoxycarbonyl) amino-4-hydroxy-N-[1'-isopropyl-1'-(4-nitroimidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide;
- 15 (2R,4S,5S,1'S)-5-(t-butoxycarbonyl) amino-4-hydroxy-N-(1'-methyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide;
- (2R,4S,5S,1'S)-5-(t-butoxycarbonyl) amino-4-hydroxy-N-(1'-propyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide;
- 20 (2R,4S,5S,1'S)-5-(t-butoxycarbonyl) amino-4-hydroxy-N-[1'-isopropyl-1'-(4-bromoimidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide;
- 25 (2R,4S,5S,1'S)-5-(t-butoxycarbonyl) amino-4-hydroxy-N-[1'-isopropyl-1'-(4,5-dibromoimidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide;
- (2R,4S,5S,1'S)-5-(t-butoxycarbonyl) amino-4-hydroxy-N-[1'-isopropyl-1'-(4-methylimidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide;
- 30 (2R,4S,5S,1'S)-5-(t-butoxycarbonyl) amino-4-hydroxy-N-[1'-isopropyl-1'-(4-trifluoromethylimidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide;
- (2R,4S,5S,1'S)-5-(t-butoxycarbonyl) amino-4-hydroxy-N-methyl-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide;
- 35

- (2R, 4S, 5S, 1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-[1'-isopropyl-1'-(4-carbomethoxyimidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide;
- (2R, 4S, 5S, 1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-[1'-isopropyl-1'-(4-methylcarbonylimidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide;
- (2R, 4S, 5S, 1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-[1'-isopropyl-1'-(4-isopropylcarbonyl-imidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide;
- 10 (2R, 4S, 5S, 1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-[1'-isopropyl-1'-(4-phenylcarbonyl-imidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide;
- (2R, 4S, 5S, 1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-[1'-isopropyl-1'-(4-formylimidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide;
- 15 (2R, 4S, 5S, 1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-[1'-isopropyl-1'-(4-(hydroxymethyl)-imidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide;
- (2R, 4S, 5S, 1'S)-5-((tetrahydrothiopyran-4-yl)oxycarbonyl)-amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide;
- 20 (2R, 4S, 5S, 1'S)-5-((tetrahydro-4H-pyran-4-yl)oxycarbonyl)-amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide;
- (2R, 4S, 5S, 1'S)-5-(4-picolinyloxy)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide;
- 25 (2R, 4S, 5S, 1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-(4,4,4-trifluorobut-1-yl)hexanamide ;
- (2R, 4S, 5S, 1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-[1'-isopropyl-1'-(4-((1RS)-1-hydroxyethyl)-imidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide;
- (2R, 4S, 5S, 1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-[1'-(1-methyl)propyl-1'-(imidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide;
- 35

- (2R, 4S, 5S, 1'S)-5-(propylaminocarbonyl) amino-4-hydroxy-N-[1'-isopropyl-1'-(imidazol-2-yl)]methyl-6-phenyl-2-phenylmethylhexanamide;
- (2R, 4S, 5S, 1'S)-5-(4-hydroxybutanoyl) amino-4-hydroxy-N-(1'-isopropyl-1'-(imidazol-2-yl)methyl-6-phenyl-2-phenylmethylhexanamide;
- (2R, 4S, 5S, 1'S)-2-phenylmethyl-4-hydroxy-5-(benzyloxy-carbonyl)valylamino-6-phenyl-N-(1'-isobutyl-1'-(imidazo-2-yl)methyl-hexanamide;
- (2R, 4S, 5S, 1'S)-2-phenylmethyl-4-hydroxy-5-(N-acetylvalyl)-amino-6-phenyl-N-(1'-isobutyl-1'-(imidazo-2-yl)methylhexanamide;
- (2R, 4S, 5S, 1'S)-5-[(imidazol-2-yl)methyloxycarbonyl]amino-4-hydroxy-N-(1'-isopropyl-1'-(imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide;
- (2R, 4S, 5S, 1'S, 1"RS)-5-((1"-(imidazol-2-yl)-2"-methyl)-propyloxycarbonyl) amino-4-hydroxy-N-(1'-isopropyl-1'-(imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide;
- (2R, 4S, 5S, 1'S)-5-(t-butoxycarbonyl) amino-4-hydroxy-N-[1'-isopropyl-1'-(4-(imidazol-2-yl)imidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide;
- (2R, 4S, 5S, 1'S)-5-(1-oxo-thian-4-yl) oxycarbonyl) amino-4-hydroxy-N-(1'-isopropyl-1'-(imidazol-2-yl)methyl-6-phenyl-2-phenylmethylhexanamide;
- (2R, 4S, 5S, 1'S)-5-((tetrahydrosulfonylpyran-4-yl) oxycarbonyl) amino-4-hydroxy-N-(1'-isopropyl-1'-(imidazol-2-yl)methyl-6-phenyl-2-phenylmethylhexanamide;
- (2R, 4S, 5S, 1'S)-5-((1,1-dimethyl-2-(benzyloxycarbonyl)-glycyloxy)ethoxycarbonyl) amino-4-hydroxy-N-(1'-isopropyl-1'-(imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide hydrochloride salt;
- (2R, 4S, 5S, 1'S)-5-((1,1-dimethyl-2-glycyloxy)ethoxycarbonyl)-amino-4-hydroxy-N-(1'-isopropyl-1'-(imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamedihydrochloridesalt;
- (2R, 4S, 5S, 1'S)-5-((1-acetyl) amino-4-hydroxy-N-(1'-isopropyl-1'-(imidazol-2-yl)methyl-6-phenyl-2-phenylmethylhexanamide;

- (2R, 4S, 5S, 1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-(4-(benzyloxyphenylmethyl)hexanamide;
- (2R, 4S, 5S, 1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-(4-hydroxyphenylmethyl)hexanamide;
- 5 (2R, 4S, 5S)-5-(t-butoxycarbonyl)amino-4-hydroxy-2-phenylmethyl-6-phenyl-N-[1'-cyclopropyl-1'-imidazol-2-yl)methyl-hexanamide;
- 10 (2R, 4S, 5S, 1'S)-5-((isopropylthiol)carbonyl)-amino-4-hydroxy-2-phenylmethyl-6-phenyl-N-[1-isopropyl-1'-imidazol-2-yl)methyl-hexanamide;
- (2R, 4S, 5S, 1'S)-5-[3-(1H-imidazol-2-yl)-3-hydroxy-4-methylpentylamido]-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide;
- 15 (2R, 4S, 5S, 1'S)-5-[(4-methoxyphenoxy)carbonyl]amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide;
- (2R, 4S, 5S, 1'S)-5-(t-butylaminocarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenylmethyl-hexanamide;
- 20 (2R, 4S, 5S, 1'S)-5-(methylaminocarbonyl)-amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenylmethyl-hexanamide;
- (2R, 4S, 5S, 1'S)-5-phenylaminocarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenylmethyl-hexanamide;
- 25 (2R, 4S, 5S, 1'S)-5-N-(propylaminocarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenylmethyl-hexanamide;
- (2R, 4S, 5S, 1'S)-5-(n-propylaminothiono)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenylmethyl-hexanamide;
- 30 (2R, 4S, 5S, 1'S)-5-(isopropylaminocarbonyl)-amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenylmethyl-hexanamide;
- (2R, 4S, 5S, 1'S)-5-(aminocarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenylmethyl-hexanamide;
- 35 (2R, 4S, 5S, 1'S)-5-(6-quinolinylmethyloxy-carbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenylmethyl-hexanamide;

- 139 -

- (2R, 4S, 5S, 1'S)-5-(benzoyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenylmethyl-hexanamide;
- (2R, 4S, 5S, 1'S)-5-(2-furylcarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenylmethyl-hexanamide;
- (2R, 4S, 5S, 1'S)-5-(4-methoxybenzoyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenylmethyl-hexanamide;
- (2R, 4S, 5S, 1'S)-5-benzylcarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenylmethyl-hexanamide;
- (2R, 4S, 5S, 1'S)-5-(4-hydroxybenzoyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenylmethyl-hexanamide;
- (2R, 4S, 5S, 1'S)-5-(cinnamoyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenylmethyl-hexanamide;
- (2R, 4S, 5S, 1'S)-5-(2-hydroxybenzoyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenylmethyl-hexanamide;
- (2R, 4S, 5S, 1'S)-5-(imidazol-4-yl-acetyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenylmethyl-hexanamide;
- (2R, 4S, 5S, 1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-(4-carbomethoxyethylimidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide;
- (2R, 4S, 5S, 1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-(4-carboxamidoimidazol-2-yl)]methyl-6-phenyl-2-phenylmethyl-hexanamide;
- (2R, 4S, 5S, 1'S)-2-phenylmethyl-4-hydroxy-5-(5-(1-oxopropyl)-2-thiazolyl)amino)-6-phenyl-N-(1'-isopropyl-1'-(imidazo-2-yl))methyl-hexanamide;
- (2R, 4S, 5S, 1'S)-2-phenylmethyl-4-hydroxy-5-(5-(1-oxopropyl)-2-thiazolyl)amino)-6-phenyl-N-(1'-isopropyl-1'-(imidazo-2-yl))methyl-hexanamide;
- (2R, 4S, 5S, 1'S)-2-phenylmethyl-4-hydroxy-5-(5-propyl-2-thiazolyl)amino)-6-phenyl-N-(1'-isopropyl-1'-(imidazo-2-yl))methyl-hexanamide; and
- (2R, 4S, 5S, 1'S)-5-(nicotinyl)amino-4-hydroxy-N-(1'-isopropyl-1'-imidazol-2-yl)methyl-6-phenylmethyl-hexanamide.

35

11. A compound according to claim 1 which is (2R, 4S, 5S, 1'S)-5-(t-butoxycarbonyl)amino-4-hydroxy-N-(1'-isopropyl-1'-(4-

- 140 -

isopropylcarbonylimidazol-2-yl)methyl-6-phenyl-2-phenylmethyl-hexanamide

12. A compound according to claim 1 which is (2R,4S,5S,1'S)-2-phenylmethyl-4-hydroxy-5-(t-butoxycarbonyl)-amino-6-phenyl-N-(1'-isopropyl-1'-(imidazo-2-yl))methyl-hexanamide.


13. A pharmaceutical composition comprising a compound according to Claim 1 and a pharmaceutically acceptable carrier.

14. A pharmaceutical formulation comprising a compound according to Claim 1 and an oil.

15. A method of treating disease states associated with HIV infection comprising administering an effective amount of a compound according to Claim 1.

16. The use of a compound according to Claim 1 in the manufacture of a medicament.


17. A compound of the formula:



(VIII)

wherein Pr^2 is an amino protecting group, and R^7 , R^8 and R^9 are as defined in Claim 1 with any reactive groups protected.

18. A compound of formula:




wherein:


- 141 -

R_1 and R_3 are each independently C_{1-6} alkyl, $Ar-C_{1-6}$ alkyl, H or $t-C_{1-6}$ alkyl, C_{2-6} alkenyl, $Ar-C_{2-6}$ alkenyl, $Het-C_{2-6}$ alkenyl, C_{3-6} cycloalkyl- C_{1-6} alkyl or C_{3-6} cycloalkenyl- C_{1-6} alkyl;

5 R_2 is H or OH ;

R_4 is R_6-NH- , or  R_6 ;

R_5 is R_6-NH- or $R_{10}-NH-$;


R_6 is  R_8 and R_9 are each independently H , OH , halo, acyl, or substituted alkyl;

wherein:

10 X is NR_{11} , O , or S ;

R_{11} is H or C_{1-3} alkyl;

R_8 and R_9 are each independently H , OH , halo, acyl, or substituted alkyl;

or R_6 is  X and Y are each independently H , OH , halo, acyl, or substituted alkyl;

15 wherein:

X is NH , O , or S ;

Y is a fused C_{2-4} alkylene, aryl or heteroaryl moiety;

R_7 is C_{1-6} alkyl, $Ar-C_{1-6}$ alkyl, $Het-C_{1-6}$ alkyl, C_{2-6} alkenyl, $Ar-C_{2-6}$ alkenyl, $Het-C_{2-6}$ alkenyl, C_{3-6} cycloalkyl- C_{1-6} alkyl or C_{3-6} cycloalkenyl- C_{1-6} alkyl;

20 R_{10} is a moiety $A-(B)_n-$, where $n = 0$ or 1 ; and B is, independently, an α -amino acid chosen from the group: Ala, Asn, Cys, Trp, Gly, Gln, Ile, Leu, Met, Phe, Pro, Ser, Thr, Tyr, Val, His, or trifluoroalanine, wherein the amino group of B is bonded to A and the carboxy group of B is bonded to the structure;

A is covalently attached to the amino group of the adjacent residue B or to the amino group of the structure if $n = 0$ and is:

- 1) trityl,
- 2) hydrogen,
- 3) C_{1-6} alkyl,
- 4) $R_{14}-CO-$ wherein R_{14} is:

- 142 -

- a) hydrogen,
- b) C₁₋₆alkyl, unsubstituted or substituted with one or more hydroxyl groups, chlorine atoms, or fluorine atoms,
- c) phenyl or naphthyl unsubstituted or substituted with one or more substituents R₁₅ wherein R₁₅ is:
- i) C₁₋₄alkyl,
 - ii) halogen, where halogen is F, Cl, Br or I,
 - iii) hydroxyl,
 - iv) nitro,
 - v) C₁₋₃alkoxy, or
 - vi) -CO-N(R₁₆)₂ wherein R₁₆ is, independently, H or C₁₋₄alkyl; or
- d) a 5-7 member heterocycle such as pyridyl, furyl, or benzisoxazolyl;
- 5) phthaloyl wherein the aromatic ring is unsubstituted or substituted with one or more substituents R₁₅;
- 6) R₁₇(R₁₈R₁₉C) m-CO- wherein m = 1-3 and R₁₇, R₁₈, and R₁₉ are independently:
- a) hydrogen,
 - b) chlorine or fluorine,
 - c) C₁₋₃alkyl unsubstituted or substituted with one or more (chlorine or fluorine atoms or hydroxyl groups,
 - d) hydroxyl,
 - e) phenyl or naphthyl unsubstituted or substituted with one or more substituents R₁₅,
 - f) C₁₋₃alkoxy,
 - g) a 5-7 member heterocycle, or
 - h) R₁₇, R₁₈, and R₁₉ may be independently joined to form a monocyclic, bicyclic, or tricycle ring system each ring of which is C₃₋₆ cycloalkyl;
- 7) R₁₇(R₁₈R₁₉C)_m-W- wherein m = 1-3 and W is OCO or SO₂ and R₁₇, R₁₈, and R₁₉ are as defined above,

- 143 -

except R17, R18, and R19 are not chlorine, fluorine or hydroxyl if they are adjacent to W;

- 8) R20-W-wherein R20 is a 5-7 member heterocycle such as pyridyl, furyl, or benzisoxazolyl;

- 9) R21-W- wherein R21 is phenyl or naphthyl unsubstituted or substituted with one or more substituents R15;

- 10) $R_{17}-(R_{18}R_{19}C)_m-P(0)(OR_{22})-$ wherein R_{22} is C_{1-4} alkyl or phenyl;

- 11) $R_{20-P} \cap (OR_{22})$; or

- 12) $R_{21}-P(O)(OR_{22})-$;

or pharmaceutically acceptable salt thereof.

INTERNATIONAL SEARCH REPORT

PCT/US92/06047

A. CLASSIFICATION OF SUBJECT MATTER

IPC(5) : C07D 233/64, 263/32, 277/30; A61K 31/415, 31/42, 31/425

US CL : Please See Extra Sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : Please See Extra Sheet.

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
STN, structure searched

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US, A, 4,431,653 (WEI ET AL.) 14 February 1984 (See Ex. 38)	17

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be part of particular relevance	X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*G* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

12 AUGUST 1992

Date of mailing of the international search report

21 DEC 1992

Name and mailing address of the ISA/
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile N . NOT APPLICABLE

Authorized officer

ROBERT GERSTL

Telephone N . (703) 308-1235

INTERNATIONAL SEARCH REPORT

International Application No.
PCT/US93/07173

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	The Journal of Antibiotics, Volume 40, issued June 1987, K. Nemoto, "Enhancement of colony Formation of mouse bone marrow cells by ubenimex", pages 894-898, see entire document.	1-26
Y	Cancer Immunology Immunotherapy, Volume 29, Issued April 1989, F. Abe et al., "Chemoimmunotherapy with cyclophosphamide and bestatin in experimental metastasis in mice", pages 231-236, see entire document.	1-26

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US92/06047

A. CLASSIFICATION OF SUBJECT MATTER:

US CL :

514/19, 365, 370, 377, 392, 397, 398, 400; 546/175, 278; 548/193, 194, 200, 204, 233, 236, 312.7, 315.1, 328.5, 332.5, 335.5, 338.1, 338.5

B. FIELDS SEARCHED

Minimum documentation searched

Classification System: U.S.

514/19, 365, 370, 377, 392, 397, 398, 400; 546/175, 278; 548/193, 194, 200, 204, 233, 236, 312.7, 315.1, 328.5, 332.5, 335.5, 338.1, 338.5